International Journal of Astrobiology

cambridge.org/ija

Spaceflight from Super-Earths is difficult

Michael Hippke

Sonneberg Observatory, Sternwartestr. 32, 9 6515 Sonneberg, Germany

Abstract

Many rocky exoplanets are heavier and larger than the Earth and have higher surface gravity. This makes space-flight on these worlds very challenging because the required fuel mass for a given payload is an exponential function of planetary surface gravity, $\exp(g_0)$. We find that chemical rockets still allow for escape velocities on Super-Earths up to 10× Earth mass. More massive rocky worlds, if they exist, would require other means to leave the planet, such as nuclear propulsion. This is relevant for space colonization and the search for extraterrestrial intelligence.

Research Article

Cite this article: Hippke M (2019). Spaceflight from Super-Earths is difficult. *International Journal of Astrobiology* **18**, 393–395. https://doi.org/10.1017/S1473550418000198

Received: 7 April 2018
Revised: 10 May 2018
Accepted: 7 June 2018
First published online: 2 August 2018

Key words:
Spaceflight; SETI; Super-Earth

Author for correspondence: Michael Hippke, E-mail: michael@jaekle.info

Introduction

Do we inhabit the best of all possible worlds (Leibnitz 1710)? From a variety of habitable worlds that may exist, Earth might well turn out as one that is marginally habitable. Other, more habitable ('superhabitable') worlds might exist (Heller and Armstrong 2014). Planets more massive than Earth can have a higher surface gravity, which can hold a thicker atmosphere and thus better shielding for life on the surface against harmful cosmic rays. Increased surface erosion and flatter topography could result in an 'archipelago planet' of shallow oceans ideally suited for biodiversity. There is apparently no limit for habitability as a function of surface gravity as such (Dorn *et al.* 2017). Size limits arise from the transition between Terran and Neptunian worlds about $2 \pm 0.6\,R_\oplus$ (Chen and Kipping, 2017). The largest rocky planets known so far are $\sim 1.87\,R_\oplus$, $\sim 9.7\,M_\oplus$ (Kepler-20 b, Buchhave *et al.*, 2016). When such planets are in the habitable zone, they may be inhabited. Can'Super-Earthlings' still use chemical rockets to leave their planet? This question is relevant for the search for extraterrestrial intelligence (SETI) and space colonization (Lingam, 2016; Forgan, 2016, 2017 and Dutil and Dumas, 2010).

Method

At our current technological level, spaceflight requires a rocket launch to provide the thrust needed to overcome Earth's force of gravity. Chemical rockets are powered by exothermic reactions of the propellant, such as hydrogen and oxygen. Other propulsion technologies with high specific impulses exist, such as nuclear thermal rockets (e.g., NERVA, Arnold and Rice, 1969), but have been abandoned due to political issues. Rockets suffer from the Tsiolkovsky equation (Tsiolkovsky, 1903): if a rocket carries its own fuel, the ratio of total rocket mass versus final velocity is an exponential function, making high speeds (or heavy payloads) increasingly expensive (Plastino and Muzzio, 1992).

The achievable maximum velocity change of a chemical rocket is

$$\Delta v = v_{\mathrm{ex}} \ln \frac{m_0}{m_{\mathrm{f}}} \tag{1}$$

where m_0 is the initial total mass (including fuel), m_{f} is the final total mass without fuel (the dry mass) and v_{ex} is the exhaust velocity. We can substitute $v_{\mathrm{ex}} = g_0\, I_{\mathrm{sp}}$ where $g_0 = \mathrm{GM}_\oplus / R_\oplus^2 \sim 9.81\,\mathrm{m\,s^{-1}}$ is the standard gravity and I_{sp} is the specific impulse (total impulse per unit of propellant), typically ~350…450 s for hydrogen/oxygen.

To leave Earth's gravitational influence, a rocket needs to achieve at minimum the escape velocity

$$v_{\mathrm{esc}} = \sqrt{\frac{2\mathrm{GM}_\oplus}{R_\oplus}} \sim 11.2\,\mathrm{km\,s^{-1}} \tag{2}$$

for Earth and v_{esc} ~27.1 km s^{-1} for a $10\,M_\oplus$, $1.7\,R_\oplus$ Super-Earth is similar to Kepler-20 b.

Results

We consider a single-stage rocket with $I_{\mathrm{sp}} = 350$ s and wish to achieve $\Delta v > v_{\mathrm{esc}}$. The mass ratio of the vehicle becomes

$$\frac{m_0}{m_{\mathrm{f}}} > \exp\left(\frac{v_{\mathrm{esc}}}{v_{\mathrm{ex}}}\right). \tag{3}$$

© Cambridge University Press 2018

CAMBRIDGE UNIVERSITY PRESS

which evaluates to a mass ratio of ∼26 on Earth and ∼2, 700 on Kepler-20 b. Consequently, a single-stage rocket on Kepler-20 b must burn 104× as much fuel for the same payload (∼2, 700 t of fuel for each t of payload).

This example neglects the weight of the rocket structure itself and, is therefore, a never achievable lower limit. In reality, rockets are multistage and have typical mass ratios (to Earth escape velocity) of 50 … 150. For example, the Saturn V had a total weight of 3,050 t for a lunar payload of 45 t, so that the ratio is 68. The Falcon Heavy has a total weight of 1,400 t and a payload of 16.8 t, so that the ratio is 83 (i.e., the payload fraction is ∼1%).

For a mass ratio of 83, the minimum rocket (1 t to v_{esc}) would carry 9, 000 t of fuel on Kepler-20 b, which is 3× larger than a Saturn V (which lifted 45 t). To lift a more useful payload of 6.2 t as required for the James Webb Space Telescope on Kepler-20 b, the fuel mass would increase to 55,000 t, about the mass of the largest ocean battleships. For a classical Apollo moon mission (45 t), the rocket would need to be considerably larger, ∼400, 000 t. This is of the order the mass of the Pyramid of Cheops and is probably a realistic limit for chemical rockets regarding cost constraints.

Discussion

Launching from a mountain top

Rockets work better in space than in an atmosphere. One might consider launching the rocket from high mountains on Super-Earths. The rocket thrust is given by

$$F = \dot{m}\, v_{ex} + A_e(P_1 - P_2) \quad (4)$$

where $\dot{m}$ is the mass flow rate, A_e is the cross-sectional area of the exhaust jet, P_1 is the static pressure inside the engine and P_2 is the atmospheric pressure. The exhaust velocity is maximized for zero atmospheric pressure, i.e. in a vacuum. Unfortunately, the effect is not very large in practice. For the Space Shuttle's main engine, the difference between sea level and vacuum is ∼25 % (Boeing, Rocketdyne Propulsion & Power, 1998). Atmospheric pressure below 0.4 bar (Earth altitude 6,000 m) is not survivable long term for humans, and presumably neither for 'Super-Earthlings'. Such low pressures are reached in lower heights on Super-Earths because the gravity pulls the air down.

One disadvantage is that the bigger something is, the less it can deviate from being smooth. Tall mountains will crush under their own weight (the 'potato radius' is ∼238 km, Caplan, 2015). The largest mountains in our Solar System are on less massive bodies, such as the Rheasilvia central peak on Vesta (22 km) or Olympus Mons on Mars (21.9 km). Therefore, we expect more massive planets to have smaller mountains. This will be detectable through transit observations in future telescopes (McTier and Kipping, 2018). One option would be to build artificial mountains as launch platforms.

Launching rockets from water-worlds

Many habitable (and presumably, inhabited) planets might be waterworlds (Simpson, 2017) and intelligent life in water and subsurface is plausible (Lingam and Loeb, 2018). Can rockets be launched from such planets? We here neglect how chemical fuels and whole rockets, are assembled on such worlds.

Rockets on waterworlds could either be launched from floating pontoon-based structures, or directly out of the water. Underwater submarine rocket launches use classical explosives to flash-vaporize water into steam. The pressure of the expanding gas drives the missile upwards in a tube. This works well for ICBMs launched from submerged submarines.

These aquatic launch complications make the theory of oceanic rocket launches appear at first quite alien; presumably land-based launches seem equally human to alien rocket scientists.

Relevance of spaceflight for SETI

It is not trivial to estimate the longevity of civilizations and the 'doomsday' argument limits the remaining lifetime of our species to between 12 and 8×10^6 yrs at 95 % confidence (Gott, 1993). If this argument is correct (Haussler, 2016), it would also apply to other civilizations and it appears reasonable to assume that population growth is intertwined with technological and philosophical progress. Indeed, the risk of catastrophic extinction has been estimated of order 0.2 % per year (Matheny, 2007; Simpson, 2016; Turchin and Denkenberger, 2018) at our current technological level and might be applicable to other civilizations within less than a few orders of magnitude (Gerig, 2012; Gerig *et al.*, 2013). The possession of ever more powerful technology in the hands of (many) individuals is increasingly dangerous (Cooper, 2013).

The risk of extinction could be reduced by colonization of other planets in a stellar system, such as Mars. If space-flight is much more difficult due to high surface gravity, colonization would be temporally delayed and the chances of survival reduced. Then, Drake's L would be (on average) smaller for civilizations on Super-Earths (Drake, 2013). Inversely, planets with lower surface gravity would make colonization cheaper and thus increase L.

Conclusion

For a payload to escape velocity, the required amount of chemical fuel scales as $\exp(g_0)$. Chemical rocket launches are still plausible for Super-Earths $\lesssim 10\,M_\oplus$, but become unrealistic for more massive planets. On worlds with a surface gravity of $\gtrsim 10\,g_0$, a sizable fraction of the planet would need to be used up as chemical fuel per launch, limiting the total number of flights. On such worlds, alternative launch methods such as nuclear-powered rockets or space elevators are required.

References

Arnold WH and Rice CM (1969) Recent NERVA technology development. *Journal of Spacecraft and Rockets* **6**, 565–569.

Boeing, Rocketdyne Propulsion & Power (1998) Space Shuttle Main Engine Orientation. *Technical Report, June 1998*, Published online: http://www.lpre.de/p_and_w/SSME/SSME_PRESENTATION.pdf

Buchhave LA, Dressing CD, Dumusque X, Rice K, Vanderburg A, Mortier A, Lopez-Morales M, Lopez E, Lundkvist MS, Kjeldsen H, Affer L, Bonomo AS, Charbonneau D, Collier Cameron A, Cosentino R, Figueira P, Fiorenzano AFM, Harutyunyan A, Haywood RD, Johnson JA, Latham DW, Lovis C, Malavolta L, Mayor M, Micela G, Molinari E, Motalebi F, Nascimbeni V, Pepe F, Phillips DF, Piotto G, Pollacco D, Queloz D, Sasselov D, Ségransan D, Sozzetti A, Udry S and Watson C. (2016) A 1.9 Earth Radius Rocky Planet and the Discovery of a Non-transiting Planet in the Kepler-20 System. *Astronomical Journal* **152**, 160.

Caplan ME (2015) Calculating the Potato Radius of Asteroids using the Height of Mt. Everest. ArXiv e-prints, arXiv:1511.04297 [physics.ed-ph]. http://arxiv.org/abs/1511.04297

https://doi.org/10.1017/S1473550418000198 Published online by Cambridge University Press

Chen J and Kipping D (2017) Probabilistic forecasting of the masses and radii of other worlds. *The Astrophysical Journal* **834**, 17.

Cooper J (2013) Bioterrorism and the Fermi paradox. *International Journal of Astrobiology* **12**, 144–148.

Dorn C, Bower DJ and Rozel AB (2017) Assessing the interior structure of terrestrial exoplanets with implications for habitability. ArXiv e-prints, arXiv:1710.05605 [astro-ph.EP]. http://arxiv.org/abs/1710.05605

Drake F (2013) Reflections on the equation. *International Journal of Astrobiology* **12**, 173–176.

Dutil Y and Dumas S (2010) Cost analysis of space exploration for extraterrestrial civilisations. In *Astrobiology Science Conference 2010: Evolution and Life: Surviving Catastrophes and Extremes on Earth and Beyond, held April 26–20, 2010 in League City, Texas*. LPI Contribution No. 1538, p. 5173. https://ui.adsabs.harvard.edu/#abs/2010LPICo1538.5173D/abstract

Forgan DH (2016) The Galactic Club, or Galactic Cliques? Exploring the limits of interstellar hegemony and the Zoo Hypothesis. ArXiv e-prints, arXiv:1608.08770 [physics.pop-ph]. http://arxiv.org/abs/1608.08770

Forgan DH (2017) Exoplanet transits as the foundation of an interstellar communications network. ArXiv e-prints, arXiv:1707.03730 [astro-ph.IM]. http://arxiv.org/abs/1707.03730

Gerig A (2012) The Doomsday argument in many worlds. ArXiv e-prints, arXiv:1209.6251. http://arxiv.org/abs/1209.6251

Gerig A, Olum KD and Vilenkin A (2013) Universal doomsday: analyzing our prospects for survival. *Journal of Clinical & Anatomic Pathology* **5**, 013.

Gott JR III (1993) Implications of the Copernican principle for our future prospects. *Nature* **363**, 315–319.

Haussler D (2016) Odds for an enlightened rather than barren future. ArXiv e-prints, arXiv:1608.05776. http://arxiv.org/abs/1608.05776

Heller R and Armstrong J (2014) Superhabitable worlds. *Astrobiology* **14**, 50–66.

Leibnitz GW (1710) *Essais de théodicée*. Paris: Flammarion.

Lingam M (2016) Interstellar Travel and Galactic Colonization: insights from Percolation Theory and the Yule Process. *Astrobiology* **16**, 418–426.

Lingam M and Loeb A (2018) Subsurface exolife. *International Journal of Astrobiology*, 1–30. doi: 10.1017/S1473550418000083.

Matheny JG (2007) Reducing the risk of human extinction. *Risk Analysis* **27** (5), 1335–1344.

McTier MA and Kipping DM (2018) Finding mountains with molehills: the detectability of exotopography. *Monthly Notices of the Royal Astronomical Society* **475**, 4978–4985.

Plastino AR and Muzzio JC (1992) On the use and abuse of Newton's second law for variable mass problems. *Celestial Mechanics and Dynamical Astronomy* **53**, 227–232.

Simpson F (2016) Apocalypse now? Reviving the Doomsday argument. *Monthly Notices of the Royal Astronomical Society* **468**, 2803–2815.

Simpson F (2017) Bayesian evidence for the prevalence of waterworlds. *Monthly Notices of the Royal Astronomical Society* **468**, 2803–2815.

Tsiolkovsky KE (1903) Issledovanie mirovykh prostransty reaktivnymi priborami (exploration of space with rocket devices). *Naootchnoye Obozreniye (Scientific Review)*. May, 12–13.

Turchin A and Denkenberger D (2018) Global catastrophic and existential risks communication scale. *Futures*. https://doi.org/10.1016/j.futures.2018.01.003

https://doi.org/10.1017/S1473550418000198 Published online by Cambridge University Press