Zygote

cambridge.org/zyg

Effects of multi-gradient equilibration during vitrification on oocyte survival and embryo development in mice

Yan Zhu[1], Zhen Zhang[1], Guang-Li Zhang[2] and Man-Xi Jiang[2]

[1]Medical Experimental Center, Guangdong Second Provincial General Hospital, Guangzhou 510317, People's Republic of China and [2]Center for Reproductive Medicine, Guangdong Second Provincial General Hospital, Guangzhou 510317, People's Republic of China

Research Article

Cite this article: Zhu Y *et al*. (2023) Effects of multi-gradient equilibration during vitrification on oocyte survival and embryo development in mice. *Zygote*. **31**: 612–619. doi: 10.1017/S0967199423000540

Received: 30 April 2023
Revised: 19 September 2023
Accepted: 29 September 2023
First published online: 24 November 2023

Keywords:
Embryo development; Fertilization; Multi-gradient equilibration; Oocyte survival; Oocyte vitrification

Corresponding author:
Man-Xi Jiang; Email: manxijiang@yahoo.com

Summary

Vitrification has been widely used for oocyte cryopreservation, but there is still a need for optimization to improve clinical outcomes. In this study, we compared the routine droplet merge protocol with modified multi-gradient equilibration vitrification for cryopreservation of mouse oocytes at metaphase II. Subsequently, the oocytes were thawed and subjected to intracytoplasmic sperm injection (ICSI). Oocyte survival and spindle status were evaluated by morphology and immunofluorescence staining. Moreover, the fertilization rates and blastocyst development were examined *in vitro*. The results showed that multi-gradient equilibration vitrification outperformed droplet merge vitrification in terms of oocyte survival, spindle morphology, blastocyst formation, and embryo quality. In contrast, droplet merge vitrification exhibited decreasing survival rates, a reduced proportion of oocytes with normal spindle morphology, and lower blastocyst rates as the number of loaded oocytes increased. Notably, when more than six oocytes were loaded, reduced oocyte survival rates, abnormal oocyte spindle morphology, and poor embryo quality were observed. These findings highlight that the vitrification of mouse metaphase II oocytes by the modified multi-gradient equilibration vitrification has the advantage of maintaining oocyte survival, spindle morphology, and subsequent embryonic development.

Introduction

Cryopreservation of gametes has become a routine technique for germplasm preservation and oocyte cryopreservation is becoming indispensable for the preservation of fertility. Oocyte vitrification is a simpler and more convenient strategy that has marked improved cryopreservation outcomes in several species, including mice (Choi *et al*., 2015), rabbits (Jiménez-Trigos *et al*., 2014), cows (Ezoe *et al*., 2015), and humans (Shanshan *et al*., 2015). However, the adverse effects of oocyte vitrification, including spindle confusion, zona pellucida (ZP) hardening, aneuploidy formation, and cytoplasm degeneration, are well known. Adverse effects are related to the duration of equilibration in the cryoprotective agent (CPA) and the speed of temperature decrease to −196°C in liquid nitrogen. In particular, the large size of oocytes (−80 to 120 μm) renders them more susceptible to damage of cellular genetic material in response to temperature changes and chemical reagents. Vitrification alters the spindle structure (Tamura *et al*., 2013), including microfilaments (Ci *et al*., 2014), but this damage can be repaired within 3 h after thawing (Ci *et al*., 2014). A recent study (Zhou *et al*., 2016) systematically investigated the effects of duration and temperature of oocyte exposure to vitrification and thawing solutions on the oocyte survival rate and fertilization rates and demonstrated that the presence of cumulus cells surrounding the cryopreserved oocytes could be beneficial to the *in vitro* fertilization of these oocytes. These observations indicated that vitrification could be improved by modifying the procedure.

Therefore, in this study we conducted two different parallel vitrification protocols (droplet merge and multi-gradient equilibration) each with five types of oocyte loading (2, 4, 6, 8 or 10 oocytes each carrier once). After thawing, the oocyte survival rates, fertilization rates, and blastocyst development were assessed, and Oct-positive cells in the blastocyst were compared between the two protocols. Additionally, we also evaluated spindle integrity and chromosomes of vitrified–thawed oocytes.

Materials and methods

Humane care and use of animals

Here, 8-week-old female and 10-week-old male BDF1 mice were used for the experiments. All mice were bred and maintained under controlled temperature (~22°C) with a 14-h light and

© The Author(s), 2023. Published by Cambridge University Press.




Figure 1. Schematic diagram of droplet merge and multi-gradient equilibration vitrification protocols. Mature mouse oocytes were transferred into holding medium (H) for a duration of 1 min and equilibrated through two different methods in equilibration solution (ES) or ES/H mixture, and treated with vitrification solution (VS) for 90 s. After the oocytes were loaded onto a Cryotop, they were immersed in liquid nitrogen. (A) Routine droplet merge and vitrification. (B) Modified multi-gradient equilibration and vitrification.

10-h dark cycle at the Animal Center, Guangdong Second Provincial General Hospital, Guangzhou, Guangdong. All animal procedures were approved by the Animal Center. All animal procedures in this study were performed in compliance with the Guide for Care and Use of Laboratory Animals of Guangdong Second Provincial General Hospital (No. 2014-KYLLM-065).

Collection of metaphase II (MII) oocytes in mice

Mice were administered intraperitoneal injections of 7.5 IU pregnant mare serum gonadotrophin (PMSG; Ningbo Sansheng Hormone Factory, Ninbo, China) followed by intraperitoneal injection of 7.5 IU human chorionic gonadotrophin (hCG; Ningbo Sansheng) 48 h later. The mice were euthanized by cervical dislocation 13–14 h after the hCG injection, and the oviducts were excised and placed in 1 ml of M2 medium (Sigma-Aldrich Chemical Co. St. Louis, MO, USA). Cumulus–oocyte complexes (COCs) were released by tearing the ampullae of the oviducts. The cumulus cells were enzymatically removed by treatment with 80 IU/ml hyaluronidase (Sigma-Aldrich) and mechanically dissociated with a glass pipette. Only morphologically normal mature MII oocytes, as determined by the presence of a first polar body, were used.

Routine droplet merge equilibration and vitrification of oocytes

In total, 2, 4, 6, 8 or 10 MII oocytes were classified to DM2, DM4. DM6, DM8 or DM10 groups, respectively, were placed into the holding medium (H) and left undisturbed for 1 min. The holding medium was mixed with equilibration solution 1 (ES1) using the tip of a transfer pipette, and two droplets were spontaneously mixed for 2 min. Then ES2 was merged with the holding medium + ES1, and the two droplets were spontaneously mixed for 2 min. The oocytes from the merged drop were transferred to ES3 and exposed undisturbed for 6–10 min. Then, oocyte(s) from ES3 were transferred to a vitrification solution (VS) for 90 s before they were loaded. The oocytes were gently but thoroughly pipetted in a VS drop to ensure a complete rinse. Different numbers of oocytes were loaded onto the tips of a Cryotop (Kuwayama *et al*., 2005a, 2005b; Kitazato Co., Minato-ku, Japan) within 90 s, taking care not to exceed 110 s following the initial exposure to the VS. Finally, once the vitrification was complete, the Cryotop was immediately capped and dropped into liquid nitrogen for storage (Figure 1A). The vitrification and thawing kits were purchased from Irvine Scientific Inc. (Santa Ana, CA). Procedures were performed strictly according to the manufacturer's instruction manual.

Modified multi-gradient equilibration and vitrification of oocytes

In total, 2, 4, 6, 8 or 10 oocytes were classified into MG2, MG4. MG6, MG8 or MG10 groups, respectively, and were suspended in holding medium (H), 4H1ES [80% H + 20% equilibration solution (ES)], 3H2ES (60% H + 40% ES), 2H3ES (40% H + 60% ES), and 1H4ES (20% H + 80% ES), respectively, for 1 min each, then transferred to ES and incubated for 6–10 min. The oocytes were then transferred to VS and incubated for 90 s (<110 s) at room temperature. The oocytes were loaded onto the tip of a Cryotop (Kitazato, Japan), capped, and immediately dropped into liquid nitrogen for storage (Figure 1B). All the oocytes were

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press

cryopreserved until they were thawed for analysis. Each vitrification via this protocol was repeated 25 times, similar to the droplet merge vitrification method.

Thawing of oocytes

Oocytes were thawed by direct immersion of the Cryotop in pre-warmed TS1 solution at 37°C for 1 min. The warmed oocytes were sequentially transferred to TS2 for 3 min each, washed twice with TS3 medium, and finally transferred to human tubal fluid (HTF) medium (Millipore Co., Billerica, MA, USA). They were incubated at 37°C in 5% CO_2 in humidified air for 2 h. The survival of oocytes was assessed by morphological analysis, which involved examining the integrity of the plasma membrane and the discoloration of ooplasm after the thawed oocytes were recovered. The oocytes that survived were inseminated by intracytoplasmic injection of spermatozoa.

Intracytoplasmic spermatozoa injection (ICSI) of oocytes

Epididymal spermatozoa were collected from the cauda epididymis of 8-week-old to 10-week-old BDF1 mice and incubated in HTF medium for 30 min at 37°C in air containing 5% CO_2. Then, 1 μl of spermatozoa suspension was mixed with 10 μl of 10% polyvinyl pyrrolidone (PVP)–HEPES-buffered CZB medium in a ICSI manipulation chamber. ICSI was performed as described previously (Jiang *et al.*, 2005) with a minor modification. Briefly, the sperm head was separated from the tail by piezo pulses at the neck region, and the head was immediately injected into a thawed oocyte. After 10 min of recovery at room temperature, the oocytes were washed at least three times and then placed into KSOMaa medium (Millipore Co.). Fertilization was evaluated by observing the formation of two pronuclei (2PN) 5 h after ICSI. The rates of embryos developed to the blastocyst were assessed on day 3.5 after insemination.

Oct4 staining and cell counting of blastocysts

The thawed blastocysts from both vitrification protocols were fixed and treated as previously described (Kong *et al.*, 2013; Zhu *et al.*, 2015). Briefly, blastocysts were fixed in 4% (w/v) paraformaldehyde in phosphate-buffered saline (PBS) for 40 min at room temperature. Fixed samples were stored in 0.3% (w/v) bovine serum albumin (BSA)/PBS for 1 week at 4°C for further analysis. For immunofluorescence, samples were permeabilized in 0.1% Triton X-100 and 0.3% BSA in PBS (w/v) at 37°C for 30 min. The samples were washed at least three times with 0.01% Triton X-100/ PBS, and then were incubated in blocking solution (150 mM glycine and 0.3% BSA in PBS) for ~30 min at 37°C. Then, the samples were treated with primary rabbit anti-Oct4 antibodies (H-134, 1:50, Santa Cruz Biotechnology, Santa Cruz, CA, USA) and FITC-labelled chicken anti-rabbit antibodies (1:100; Invitrogen, Grand Island, NY, USA), and diluted in blocking solution for 30–40 min each at 37°C or overnight at 4°C. After three washes (each time for 5 min), the nucleus was stained with 10 μg/ml 4′-6-diamidino-2-phenylindole (DAPI). The samples were then mounted on slides with anti-fade mounting medium, and examined under a laser-scanning confocal microscope (Zeiss LSM 510 Meta; Carl Zeiss AG, Oberkochen, Germany). Blastomeres expressing positive Oct4 showed green fluorescence and intact nuclei were blue. The ratio of blastomeres expressing green fluorescence to the total blastomeres with blue nuclei was considered the Oct4-positive rate (Figure 4A). The images shown in the Results section are representative of at least 30 samples from more than 25 experimental replicates.

Based on the results of the number of Oct4-positive blastomeres, blastocysts were graded as Grade A, B, or C. Blastocysts with more than 15 Oct4-positive blastomeres were classified as Grade A blastocysts; those with 8–15 Oct-positive blastomeres were classified as Grade B blastocysts; and those with 0–8 blastomeres were classified as Grade C blastocysts. Blastocysts of grades A and B were considered to be good-quality embryos.

Meiotic spindle and chromosome evaluation of oocytes

Microtubules of oocytes were analyzed as described previously with minor modifications (Zhu *et al.*, 2003, 2015) in two vitrification protocols. In brief, after vitrification and thawing, the oocytes were incubated in HTF medium at 37°C for 2 h in an atmosphere containing 5.0% CO_2 and then fixed in 4% paraformaldehyde for 10 min. Then the oocytes were washed in PBS and transferred to 0.25% Triton X-100/ PBS (w/v) for 10 min at room temperature. The oocytes were washed twice for 5 min each and blocked for 1 h at room temperature with 2% BSA in PBS. After washing twice with PBS, the oocytes were incubated with fluorescein isothiocyanate (FITC)-labelled mouse monoclonal antibodies against α-tubulin (Sigma-Aldrich Co.), and diluted with blocking solution (1:100) for 30–40 min at 37°C or overnight at 4°C. After washing three times for 5 min each time, the nuclei were stained with 10 μg/ml DAPI (Sigma-Aldrich Co.). Finally, the samples were mounted in an antifluorescence-fade medium (DABCO, Sigma-Aldrich Co.) on slides. They were then observed under a laser-scanning confocal microscope (Zeiss LSM 510 Meta). Typical barrel-shaped microtubules traversing the two poles and centrally aligned chromosomes were considered normal. Images shown in the Results are representatives of at least 20 samples from each group.

Statistical analysis

Data were analyzed using Prism 6.0 (GraphPad Software Inc., La Jolla, CA, USA). The survival rates and fertilization rates were compared using one-way analysis of variance (ANOVA) from Prism 6.0. The proportions of oocytes with normal spindle, blastocyst rates and proportions of good-quality blastocyst were compared using chi-squared test. A P-value of <0.05 was considered statistically significant.

Results

Effects of different vitrification protocols on oocyte survival, fertilization, and blastocyst formation

The multi-gradient equilibration vitrification protocol displayed higher survival (Figure 2, Table S1), fertilization rates (Figure 2, Table S2), and blastocyst rates (Table 1) compared with the droplet merge vitrification protocol, regardless of the number of oocytes loaded onto each Cryotop ($P > 0.05$).

However, the droplet merge vitrification protocol showed a reduced survival rate, fertilization rate and blastocyst formation rate (Figure 2 and Table 1). Lower survival rates (Figure 2) were observed when the loading number exceeded six oocytes, and the fertilization rates (Fig 2) and blastocyst formation (Table 1) rates decreased when the loading number exceeded eight. There were significant differences in the oocyte survival rates between the two protocols when more than six oocytes were loaded ($P < 0.001$).

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press




Figure 2. Survival rates of oocytes and fertilization rates of droplet merge and multi-gradient equilibration vitrification protocols. (A) Oocyte survival rates achieved using the droplet merge and multi-gradient equilibration vitrification protocols. (B) Fertilization rates achieved using the droplet merge and multi-gradient equilibration vitrification protocols. (C) Comparison of survival and fertilization rates between droplet merge and multi-gradient equilibration vitrification protocols when the number of loaded oocytes is the same. Different superscript letters on the bars in (A) and (B) indicate $P < 0.05$. Asterisks *** and **** on the columns in (C) indicate $P < 0.05$, 0.001 and 0.0001. NS: No significant difference.

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press

Table 1. Blastocyst rates between droplet merge and multi-gradient equilibration vitrification protocols

	Blastocyst rates (%)				
Number of loaded oocytes	2	4	6	8	10
Droplet merge	23/38 (60.53)[a]	48/80 (60.00)[a]	43/82 (52.44)[a,b]	30/69 (43.48)[b]	19/58 (32.76)[b]
Multi-gradient equilibration	24/39 (61.54)[a]	48/79 (60.76)[a]	75/121 (61.98)[a]	100/164 (60.98)[a]	124/206 (60.19)[a]

Values with different superscripts within columns are significantly different at $P < 0.05$ (chi-squared test).




Figure 3. Spindle morphology of oocytes after thawing of droplet merge and multi-gradient equilibration vitrification protocols. (A) Representative images of oocyte spindles from droplet merge or multi-gradient equilibration and vitrification protocols. Green: TUBB (tubulin); Blue: Nuclei; Merge: overlay of TUBB and nuclei. Scale bar, 10 μm. (B) Differences in the percentage of oocytes exhibiting normal spindle morphology between droplet merge and multi-gradient equilibration vitrification protocols when the number of loaded oocytes was the same. Different letters on the bars in (B) indicate $P < 0.05$.

Additionally, the fertilization rates ($P < 0.05$) and blastocyst formation rates ($P < 0.05$) did significantly differ between the two protocols when eight oocytes were loaded onto the Cryotops (Figure 2, Table 1).

In addition, we also found significant differences in oocyte survival rates for 6–10 oocytes loaded onto each Cryotop ($P < 0.001$, Figure 2), but not for 2–4 oocytes. Furthermore, there were differences in oocyte fertilization rates when the loading number exceeded six ($P < 0.05$, Figure 2).

Modified vitrification maintains oocytes with normal spindle morphology

After thawing and incubation in HTF medium for 2 h, the proportion of oocytes with normal spindles significantly decreased with an increase in the number of oocytes loaded onto a Cryotop in the droplet merge vitrification protocol. Significant differences were observed when the number of loaded oocytes ranged from 8–10 ($P < 0.05$; Figure 3B; Table S3). Conversely, no similar results were observed with the multi-gradient equilibration vitrification protocol, regardless of the number of oocytes loaded onto the Cryotop ($P > 0.05$; Figure 3B; Table S). These results indicate that the modified multi-gradient equilibration vitrification protocol is a more stable procedure, resulting in a higher percentage of oocytes with normal spindles compared with the routine droplet merge vitrification.

Additionally, we found that the spindle morphology was significantly altered with 8–10 oocytes per Cryotop ($P < 0.05$, Figure 3B), but not with 2–6 oocytes per Cryotop ($P > 0.$ 05; Figure 3B).

Effects of different vitrification protocols and oocyte loading number on Oct4 expression in expanded blastocysts

When 2, 4, 6, 8 or 10 oocytes were loaded in the droplet merge vitrification protocol, the percentages of good-quality blastocysts (grade A+B) obtained were 75.00%, 73.91%, 71.43%, 39.13%, and 11.54%, respectively. Conversely, for the multi-gradient equilibration vitrification protocol, the percentages of good-quality blastocysts obtained were 77.27%, 76.92%, 77.78%, 76.92%, and 72.73% when 2, 4, 6, 8, and 10 oocytes were loaded onto the Cryotop, respectively (Figure 4; Table S4). These results indicated that there was a significant difference in the percentage of good-quality blastocysts (grades A and B) between the droplet merge and multi-gradient equilibration vitrification protocols ($P < 0.05$; Figure 4B) when the number of oocytes loaded onto the

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press




Figure 4. Oct4 expression and its positive percentage in expanded blastocysts of droplet merge and multi-gradient equilibration vitrification protocols. (A) Representative images of Oct4 expression in expanded mouse blastocysts derived from two vitrification protocols. Upper panels show an example of a Grade A blastocyst with more than 15 Oct4-positive blastomeres. Middle panels show an example of Grade B blastocysts with 8–15 Oct4-positive blastomeres, and lower panels show an example of Grade C blastocysts with <8 Oct4-positive blastomeres. Oct4 expression is shown in green and 4′-6-diamidino-2-phenylindole (DAPI)-stained nuclei are shown in blue. Scale bar: 20 μm. (B) Proportions of Grade A+B blastocysts are shown for each protocol. Different letters on the bars indicate $P < 0.05$.

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press

Cryotop was between 8 and 10. It is evident that multi-gradient equilibration vitrification is more beneficial for embryonic development.

Discussion

Oocyte cryopreservation is an important fertility preservation technique in humans and various animal species. Cryopreservation of human oocytes holds great promise for preserving fertility in cancer patients who must undergo radiation or chemotherapy and can also be used for oocyte donation programmes (Huang *et al.*, 2007), and it is a promising technique for preserving female fertility in failed testicular sperm extraction cycles (Song *et al.*, 2010). Mouse oocyte cryopreservation is important as it provides preliminary data that may indirectly be applicable to human oocyte cryopreservation, and these oocyte cryopreservation techniques can be used to preserve invaluable genetic resources or endangered species.

Vitrification is an alternative to traditional cryopreservation methods (slow freezing) to avoid chilling injury and ice crystal formation (Rall and Fahy, 1985; Gupta *et al.*, 2007). Some studies have shown the success of cryopreserving mature bovine oocytes using very rapid cooling methods and brief exposure to vitrification solutions (Martino *et al.*, 1996; Vajta *et al.*, 1998). Kuleshova *et al.* (1999) recently reported the first case of birth from vitrified human oocytes. At present, oocytes are mainly vitrified using single-step (ultra-rapid) and stepwise protocols in animal studies and for human clinical applications, but these protocols were unsatisfactory and appeared to need more improvements.

In our present study, mouse MII oocytes were cryopreserved using the routine droplet merge vitrification protocol according to the manual of oocyte vitrification kits (Figure 1A), and multi-gradient equilibration vitrification that was modified from the droplet merge vitrification protocol (Figure 1B).

Our results from the routine droplet merge vitrification showed that the survival rates, fertilization rates, and blastocyst rates of thawed oocytes significantly decreased with an increase in the number of oocytes loaded per Cryotop. Additionally, morphological analysis of the oocyte spindle revealed an increased proportion of oocytes with abnormal spindle morphology as the number of oocytes loaded increased. In contrast, the multi-gradient equilibration vitrification exhibited higher survival rates (Figure 2A) and blastocyst rates (Table 1), regardless of the oocyte loading number. These findings suggest that the droplet merge vitrification is not suitable for simultaneous cryopreservation of a large number of oocytes. Therefore, the multi-gradient equilibration vitrification method is more suitable for the simultaneous cryopreservation of a greater number of oocytes. This suggests that, if our intention is to cryopreserve a larger quantity of oocytes at once, the multi-gradient equilibration vitrification approach may be a preferable option.

Oocyte cryopreservation could lead to ZP hardening (Johnson *et al.*, 1988); therefore, conventional IVF is unsuccessful due to ZP modifications, necessitating ICSI for fertilization. Moreover, the cytoskeletal structure is very sensitive to temperature changes (Tamura *et al.*, 2013). Both are not conducive to sperm penetration into oocytes and fertilization. Many studies have demonstrated that incubating oocytes in a fertilization medium for 15 min to several hours after freezing could rescue cytoskeletal disruption, which benefits fertilization and embryo development (Eroglu *et al.*, 1998). In our study, oocytes that were inseminated by ICSI after incubation in HTF for 2 h showed higher rates of fertilization and embryo development. Another study (Chang *et al.*, 2011) showed that phase transition and low temperature caused little effect on the mouse oocyte spindle morphology during the vitrification and warming process, as a result of which the oocyte spindle was able to recover immediately after warming. However, in our study, we found some oocytes with abnormal spindle morphology even after the oocytes were allowed to recover for 2 h; therefore, we proposed that recovery in fertilization medium for adequate time before fertilization allows the reorganization of tubulin and microfilament components inside oocytes, which in turn leads to improved fertilization and embryo development. Similar to our results, some studies in humans reported that oocyte spindles that were allowed to recover for a period of time after vitrification resulted in improved embryonic development (Rienzi *et al.*, 2004; Gao *et al.*, 2009) and this recovery may result in comparable cleavage time, cell number, and DNA methylation patterns between fresh and vitrified oocytes (De Munck *et al.*, 2015).

Oct4 is a specific gene marker for the inner cell mass (ICM) at the expanded blastocyst stage. The expanded blastocysts were categorized into three grades based on the number of Oct4-positive cells to assess blastocyst quality (Figure 4A). To evaluate the quality of embryos derived from cryopreserved-thawed oocytes, we specifically examined the relationship between vitrification protocols and Oct4 expression in expanded blastocysts. Our results showed that embryos derived from the modified protocol had a higher percentage of pluripotent cells in the ICM, as evidenced by Oct4 expression, and a higher blastocyst development rate. Blastocysts from the multi-gradient equilibration vitrification protocol exhibited superior quality compared with those from the droplet merge vitrification method. These findings provide clear evidence that the utilization of the multi-gradient vitrification protocol results in significantly higher quality blastocysts, further supporting its suitability for oocyte cryopreservation.

In summary, routine droplet merge vitrification offers certain advantages for human oocyte cryopreservation, including high survival and fertilization rates. However, it is limited to vitrifying only 2–4 oocytes at a time. In particular, this procedure is time-consuming for animal oocyte cryopreservation, results in cryocarrier wastage, and requires significant storage space. Consequently, the droplet merge vitrification has clear drawbacks and is unsuitable for freezing a larger number of oocytes at one time or establishing an oocyte bank. Conversely, multi-gradient equilibration vitrification can be used to freeze ~10 oocytes at a time, and the oocyte survival rates, proportions with normal spindle morphology, fertilization rates and high-quality blastocyst rates obtained using this protocol are markedly high. Additionally, this multi-gradient equilibration vitrification has some advantages in reducing the total time of vitrification, minimizing the number of cryocarriers, and saving storage space. Therefore, it is recommended for use in establishing human oocyte banks and preserving valuable animal resources.

Supplementary material. To view supplementary material for this article, please visit https://doi.org/10.1017/S0967199423000540

Acknowledgements. This work was supported by grants from the Guangdong Science and Technology Project funds (2017A020214019), the Guangzhou Science and Technology Project fund (Grant number 201904010058) and the introduction of talent scientific research start-up fund of Guangdong Second Provincial General Hospital (grant number no. YY2017–003).

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press

Competing financial interests. The authors declare no competing financial interests.

References

Chang, C. C., Lin, C. J., Sung, L. Y., Kort, H. I., Tian, X. C. and Nagy, Z. P. (2011). Impact of phase transition on the mouse oocyte spindle during vitrification. *Reproductive Biomedicine Online*, **22**(2), 184–191. doi: 10.1016/j.rbmo.2010.10.009

Choi, J. K., Huang, H. and He, X. (2015). Improved low-CPA vitrification of mouse oocytes using quartz microcapillary. *Cryobiology*, **70**(3), 269–272. doi: 10.1016/j.cryobiol.2015.04.003

Ci, Q., Li, M., Zhang, Y., Ma, S., Gao, Q. and Shi, Y. (2014). Confocal microscopic analysis of the microfilament configurations from human vitrification-thawed oocytes matured *in vitro*. *Cryo Letters*, **35**(6), 544–548.

De Munck, N., Petrussa, L., Verheyen, G., Staessen, C., Vandeskelde, Y., Sterckx, J., Bocken, G., Jacobs, K., Stoop, D., De Rycke, M. and Van de Velde, H. (2015). Chromosomal meiotic segregation, embryonic developmental kinetics and DNA (hydroxy)methylation analysis consolidate the safety of human oocyte vitrification. *Molecular Human Reproduction*, **21**(6), 535–544. doi: 10.1093/molehr/gav013

Eroglu, A., Toner, M., Leykin, L. and Toth, T. L. (1998). Cytoskeleton and polyploidy after maturation and fertilization of cryopreserved germinal vesicle-stage mouse oocytes. *Journal of Assisted Reproduction and Genetics*, **15**(7), 447–454. doi: 10.1007/BF02744940

Ezoe, K., Yabuuchi, A., Tani, T., Mori, C., Miki, T., Takayama, Y., Beyhan, Z., Kato, Y., Okuno, T., Kobayashi, T. and Kato, K. (2015). Developmental competence of vitrified-warmed bovine oocytes at the germinal-vesicle stage is improved by cyclic adenosine monophosphate modulators during *in vitro* maturation. *PLOS ONE*, **10**(5), e0126801. doi: 10.1371/journal.pone.0126801

Gao, S., Li, Y., Gao, X., Hu, J., Yang, H. and Chen, Z. J. (2009). Spindle and chromosome changes of human MII oocytes during incubation after slow freezing/fast thawing procedures. *Reproductive Sciences*, **16**(4), 391–396. doi: 10.1177/1933719108327590

Gupta, M. K., Uhm, S. J. and Lee, H. T. (2007). Cryopreservation of immature and *in vitro* matured porcine oocytes by solid surface vitrification. *Theriogenology*, **67**(2), 238–248. doi: 10.1016/j.theriogenology.2006.07.015

Huang, J. Y., Buckett, W. M., Gilbert, L., Tan, S. L. and Chian, R. C. (2007). Retrieval of immature oocytes followed by *in vitro* maturation and vitrification: A case report on a new strategy of fertility preservation in women with borderline ovarian malignancy. *Gynecologic Oncology*, **105**(2), 542–544. doi: 10.1016/j.ygyno.2007.01.036

Jiang, M. X., Zhu, Y., Zhu, Z. Y., Sun, Q. Y. and Chen, D. Y. (2005). Effects of cooling, cryopreservation and heating on sperm proteins, nuclear DNA, and fertilization capability in mouse. *Molecular Reproduction and Development*, **72**(1), 129–134. doi: 10.1002/mrd.20328

Jiménez-Trigos, E., Vicente, J. S. and Marco-Jiménez, F. (2014). First pregnancy and live birth from vitrified rabbit oocytes after intra-oviductal transfer and *in vivo* fertilization. *Theriogenology*, **82**(4), 599–604. doi: 10.1016/j.theriogenology.2014.05.029

Johnson, M. H., Pickering, S. J. and George, M. A. (1988). The influence of cooling on the properties of the zona pellucida of the mouse oocyte. *Human Reproduction*, **3**(3), 383–387. doi: 10.1093/oxfordjournals.humrep.a136712

Kong, P. C., Zhu, Y., Wang, M. S., Li, H. P., Chen, X. J. and Jiang, M. X. (2013). Reprogramming of round spermatids by the germinal vesicle cytoplasm in mice. *PLOS ONE*, **8**(10), e78437. doi: 10.1371/journal.pone.0078437

Kuleshova, L., Gianaroli, L., Magli, C., Ferraretti, A. and Trounson, A. (1999). Birth following vitrification of a small number of human oocytes: Case report. *Human Reproduction*, **14**(12), 3077–3079. doi: 10.1093/humrep/14.12.3077

Kuwayama, M., Vajta, G., Ieda, S. and Kato, O. (2005a). Comparison of open and closed methods for vitrification of human embryos and the elimination of potential contamination. *Reproductive Biomedicine Online*, **11**(5), 608–614. doi: 10.1016/s1472-6483(10)61169-8

Kuwayama, M., Vajta, G., Kato, O. and Leibo, S. P. (2005b). Highly efficient vitrification method for cryopreservation of human oocytes. *Reproductive Biomedicine Online*, **11**(3), 300–308. doi: 10.1016/s1472-6483(10)60837-1

Martino, A., Songsasen, N. and Leibo, S. P. (1996). Development into blastocysts of bovine oocytes cryopreserved by ultra-rapid cooling. *Biology of Reproduction*, **54**(5), 1059–1069. doi: 10.1095/biolreprod54.5.1059

Rall, W. F. and Fahy, G. M. (1985). Ice-free cryopreservation of mouse embryos at −196 degrees C by vitrification. *Nature*, **313**(6003), 573–575. doi: 10.1038/313573a0

Rienzi, L., Martinez, F., Ubaldi, F., Minasi, M. G., Iacobelli, M., Tesarik, J. and Greco, E. (2004). Polscope analysis of meiotic spindle changes in living metaphase II human oocytes during the freezing and thawing procedures. *Human Reproduction*, **19**(3), 655–659. doi: 10.1093/humrep/deh101

Shanshan, G., Mei, L., Keliang, W., Yan, S., Rong, T. and Zi-Jiang, C. (2015). Effect of different rehydration temperatures on the survival of human vitrified-warmed oocytes. *Journal of Assisted Reproduction and Genetics*, **32**(8), 1197–1203. doi: 10.1007/s10815-015-0480-8

Song, W. Y., Sun, Y. P., Jin, H. X., Xin, Z. M., Su, Y. C., Guo, Y. H. and Chen, Z. J. (2010). [Clinical application of oocyte vitrification in failed testicular sperm extraction cycles: Report of 8 cases]. *Zhonghua Nan Ke Xue*, **16**(4), 305–309.

Tamura, A. N., Huang, T. T. and Marikawa, Y. (2013). Impact of vitrification on the meiotic spindle and components of the microtubule-organizing center in mouse mature oocytes. *Biology of Reproduction*, **89**(5), 112. doi: 10.1095/biolreprod.113.108167

Vajta, G., Holm, P., Kuwayama, M., Booth, P. J., Jacobsen, H., Greve, T. and Callesen, H. (1998). Open Pulled Straw (OPS) vitrification: A new way to reduce cryoinjuries of bovine ova and embryos. *Molecular Reproduction and Development*, **51**(1), 53–58. doi: 10.1002/(SICI)1098-2795(199809)51:1<53::AID-MRD6>3.0.CO;2-V

Zhou, C. J., Wang, D. H., Niu, X. X., Kong, X. W., Li, Y. J., Ren, J., Zhou, H. X., Lu, A., Zhao, Y. F. and Liang, C. G. (2016). High survival of mouse oocytes using an optimized vitrification protocol. *Scientific Reports*, **6**, 19465. doi: 10.1038/srep19465

Zhu, Z. Y., Chen, D. Y., Li, J. S., Lian, L., Lei, L., Han, Z. M. and Sun, Q. Y. (2003). Rotation of meiotic spindle is controlled by microfilaments in mouse oocytes. *Biology of Reproduction*, **68**(3), 943–946. doi: 10.1095/biolreprod.102.009910

Zhu, Y., Jiang, Y. H., He, Y. P., Zhang, X., Sun, Z. G., Jiang, M. X. and Wang, J. (2015). Knockdown of regulator of G-protein signalling 2 (Rgs2) leads to abnormal early mouse embryo development *in vitro*. *Reproduction, Fertility, and Development*, **27**(3), 557–566. doi: 10.1071/RD13269

https://doi.org/10.1017/S0967199423000540 Published online by Cambridge University Press