Personality Neuroscience

cambridge.org/pen

Editorial

Cite this article: Liu W, Pluta A, Charpentier CJ, and Rosenblau G. (2025) A computational cognitive neuroscience approach for characterizing individual differences in autism: Introduction to Special Issue. *Personality Neuroscience*. Vol **8**: e2, 1–7. doi: 10.1017/pen.2025.2

Received: 7 March 2025
Accepted: 8 March 2025

Corresponding author:
Gabriela Rosenblau;
Email: grosenblau@gwu.edu

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives licence (https://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided that no alterations are made and the original article is properly cited. The written permission of Cambridge University Press must be obtained prior to any commercial use and/or adaptation of the article.

A computational cognitive neuroscience approach for characterizing individual differences in autism: Introduction to Special Issue

Wenda Liu[1,2], Agnieszka Pluta[3], Caroline J. Charpentier[4,5,6] and Gabriela Rosenblau[1,2]

[1]Department of Psychological and Brain Sciences, George Washington University, Washington, DC, USA; [2]Autism and Neurodevelopmental Disorders Institute, George Washington University and Children's National Medical Center, Washington, DC, USA; [3]Faculty of Psychology, University of Warsaw, Warszawa, Poland; [4]Department of Psychology, University of Maryland College Park, College Park, MD, USA; [5]Brain and Behavior Institute, University of Maryland College Park, College Park, MD, USA and [6]Program in Neuroscience and Cognitive Science, University of Maryland College Park, College Park, MD, USA

Abstract

Traditional psychological research has often treated inter-subject variability as statistical noise (even, *nuisance* variance), focusing instead on averages rather than individual differences. This approach has limited our understanding of the substantial heterogeneity observed in neuropsychiatric disorders, particularly autism spectrum disorder (ASD). In this introduction to a special issue on this theme, we discuss recent advances in cognitive computational neuroscience that can lead to a more systematic notion of core symptom dimensions that differentiate between ASD subtypes. These advances include large participant databases and data-sharing initiatives to increase sample sizes of autistic individuals across a wider range of cultural and socioeconomic backgrounds. Our perspective helps to build bridges between autism symptomatology and individual differences in autistic traits in the non-autistic population and introduces finer-grained dynamic methods to capture behavioral dynamics at the individual level. We specifically focus on how cognitive computational models have emerged as powerful tools to better characterize autistic traits in the general population and autistic population, particularly with respect to social decision-making. We finally outline how we can combine and harness these recent advances, on the one hand, big data initiatives, and on the other hand, cognitive computational models, to achieve a more systematic and nuanced understanding of autism that can lead to improved diagnostic accuracy and personalized interventions.

For most of the past century, psychological research has viewed inter-subject variability in behavior as mere statistical noise around a *true* value, focusing primarily on central tendencies like averages or medians of distributions (Molenaar, 2004; Nesselroade, 2004; Rozin, 2001) – highlighting its problematic status, sometimes this variation is termed a *nuisance*. Traditional statistical models aimed to predict differences in the average population based on specific experimental manipulations (Molenaar, 2004; Nesselroade, 2004; Rozin, 2001). While these experimental designs and methods have helped to develop important theories across psychological domains, they have significant drawbacks. As noted by Wundt much earlier, experimental manipulations typically explain only a small amount of variance in the data and, relatedly, often lack reproducibility, contributing to a replication crisis across the field (Hedge et al., 2018; Kravitz & Mitroff, 2023; Siritzky et al., 2023). This apparent crisis has led to calls for better characterization of behavior through larger sample sizes, robust measures guided by theoretical assumptions formulated as *a priori* hypotheses, and increased transparency in the design and conduct of studies. Open science initiatives have contributed to making these changes by supporting data sharing and study preregistration (Collaboration, 2012, 2015; Kravitz & Mitroff, 2023; Siritzky et al., 2023).

Individual differences research, on the other hand, relies on large sample sizes and data-driven analysis of variability. Its broad, unconstrained study design enhances our ability to capture behavioral diversity and can produce more replicable findings compared to experimental studies with constrained homogeneous samples. Along with advancements in data science, particularly the big data approach (Adjerid & Kelley, 2018; Gomez-Marin et al., 2014; Harlow & Oswald, 2016), there have been calls for better characterization of interindividual differences across cognitive domains (Adjerid & Kelley, 2018; Eisenberg et al., 2019; Gomez-Marin et al., 2014; Harlow & Oswald, 2016). Researchers have long

suspected that deviations from the mean are not random noise but systematic variation. However, small sample sizes in most in-person psychological research have hindered the systematic investigation of these interindividual differences. Over the past couple of decades, individual differences have become a growing interest in psychological research. Personality psychology has spearheaded this movement, developing models of personality types (Ilmini & Fernando, 2017; Patzelt et al., 2018; Phan & Rauthmann, 2021), creativity (Lloyd-Cox et al., 2023; Mejia et al., 2021; Minai et al., 2021; Saunders & Bown, 2015), and cognitive styles (Riding & Rayner, 2013; Zhang, 2002) that could explain differences in perception (Haehner et al., 2023; Zhu et al., 2018), attention (Subramanian et al., 2013), and decision-making (Frolichs et al., 2022; King-Casas et al., 2008; Subramanian et al., 2013). While this field has provided evidence for interindividual variation in nonclinical groups, much less is known about the role of interindividual variability in neuropsychiatric disorders.

Across neuropsychiatric disorders, there is larger variability or greater interindividual differences in self-reported or observed behaviors such as personality traits and social skills (Jauk & Kanske, 2019) compared to control groups (Lochner & Stein, 2003; Steinhausen, 2009; Wolfers et al., 2019). These differences, combined (almost inevitably) with smaller sample sizes in clinical studies, present significant hurdles for improving diagnosis and treatment decisions (Beauchaine & Cicchetti, 2016; Jacob et al., 2019; Wolff et al., 2018). This issue is particularly pronounced in autism spectrum disorder (ASD), where formalizing heterogeneity is seen as critical for linking genetic profiles to neural and behavioral phenotypes (Pelphrey et al., 2011; Wolfers et al., 2019). Autism is marked by significant challenges in social interaction and communication, linked to differences in social perceptual and cognitive abilities across the diagnostic spectrum. The complex and diverse neurodevelopmental characteristics of autism are further compounded by interindividual variability observed in many other neuropsychiatric disorders (Jacob et al., 2019). This has led to a shift from viewing autism as unidimensional to one seeing it as an overall term including multiple syndromes, each resulting from different etiological pathways (Amaral et al., 2008; Geschwind & Levitt, 2007; Insel et al., 2010).

Addressing the heterogeneity in autism is imperative for two reasons. First, it can enhance diagnostic accuracy, which currently relies heavily on clinical observations only (Geschwind & Levitt, 2007; Insel et al., 2010; Wolff et al., 2018). Second, it can facilitate the development of more targeted cognitive-behavioral interventions for core autism symptoms, which remain elusive and rarely based on individual symptoms. However, in order to characterize and formalize the heterogeneity within ASD, several significant obstacles must be overcome. These include (1) increasing sample sizes of autistic individuals from a wider range of cultural and socioeconomic backgrounds, (2) building bridges between autism symptomatology and individual differences in autistic traits in the non-autistic population, and (3) introducing finer-grained dynamic methods to capture behavioral dynamics at the individual level. Here, we briefly describe emerging literature on the first two obstacles, before providing our perspective on how to overcome the third and more challenging one of developing finer-grained assessments and cognitive models of autism symptoms.

With respect to increasing sample sizes of autism research, there have been several initiatives spearheaded by the National Institute of Mental Health to support interdisciplinary research aimed at advancing the understanding of ASD and developing new interventions. The Autism Cluster of Excellence (ACE) initiative brings multiple research sites together jointly to investigate the underlying causes of ASD, including genetic, environmental, and developmental factors. ACE centers foster collaboration among researchers from diverse disciplines, including neuroscience, genetics, psychology, and education. The commitment to data sharing in the ACE program has led to large databases of behavioral and neural data such as the National Database for Autism Research and the Autism Brain Imaging Data Exchange. These data-sharing efforts encompass various types of data, including genetics, genomics, brain imaging, and behavioral assessments. This rich and diverse dataset is invaluable for testing the replicability of findings from smaller, single-site studies, enhancing the robustness and generalizability of research findings in the field of autism (Heinsfeld et al., 2018; Nielsen et al., 2013; Payakachat et al., 2016).

To address the second point, building bridges between autism symptomatology in individuals diagnosed with autism and autistic traits in the non-autistic population, there is a need to investigate autistic traits in larger and more diverse samples of individuals with an autism diagnosis alongside individuals without a diagnosis. Measuring self-reported autistic traits alongside objective measures of autistic symptom domains across diagnosed and undiagnosed individuals would shed light on the convergence of autism symptomatology and autistic traits in the general population. Initiatives like the Simons Foundation Autism Research Initiative provide an important resource for recruiting larger and heterogeneous groups of individuals with an autism diagnosis for behavioral studies. This initiative provides a valuable resource for researchers aiming to investigate ASD across different stages of life, from infancy through adulthood. The Simons Foundation Powering Autism Research for Knowledge (SPARK) initiative has aggregated a cohort of over 50,000 individuals with ASD and their families, who are interested in contributing to research. Researchers can access this large and diverse pool of participants nationwide for online behavioral studies. They can additionally apply to utilize existing phenotypic and genetic data on their recruited participants. SPARK has become increasingly popular for recruiting participants across developmental stages and levels of functioning, allowing studies to investigate variability in autism symptoms, including the co-occurrence of certain autism risk genes in phenotypes (Gaugler et al., 2014; Grove et al., 2019; Matoba et al., 2020; Myers et al., 2020; Wilfert et al., 2021), as well as a more thorough investigation of the influence of sex and gender differences in autism (Dillon et al., 2023; Fombonne et al., 2020; Saré & Smith, 2020). Large online studies leveraging such databases are beginning to shed light on the similarities and differences in autistic traits between people with autism or their family members and those without a close relative diagnosed with autism (Bora et al., 2017; Ruzich et al., 2016).

The third and most challenging obstacle is developing finer-grained assessments and cognitive models of autism symptoms that can help bridge the gap to animal models and genetics, on the one hand, and to real-world behavioral outcomes, on the other hand. This need is not unique to studying autism but is much needed across neuropsychiatric domains. Indeed, researchers have acknowledged the need for a computational psychiatry approach to precision phenotyping, increasing the precision with which we characterize certain sub-phenotypes of the autism spectrum (Tiego et al., 2023). A more nuanced approach to characterizing and quantifying the observed behavioral differences and their biological correlates can deepen our understanding of

https://doi.org/10.1017/pen.2025.2 Published online by Cambridge University Press

neuropsychiatric disorders and their symptoms (Friston et al., 2014; Hitchcock et al., 2022; Huys et al., 2016; Montague et al., 2012; Wang & Krystal, 2014). In this respect, computational psychiatry describes mathematical approaches to quantitatively analyze the complex interactions across biobehavioral system levels within and between neuropsychiatric disorders (Frässle et al., 2018; Karvelis et al., 2023; Petzschner et al., 2017; Stephan & Mathys, 2014; Wiecki et al., 2015). The hope of computational psychiatry is to identify nuanced patterns of behavior as well as their underlying cognitive mechanisms and neural implementation. This latter goal can be achieved through cognitive computational modeling.

Computational modeling can reveal clinically meaningful individual differences

Cognitive computational models have been widely used in the field of cognitive neuroscience (Castelfranchi & Falcone, 2010; Farrell & Lewandowsky, 2018; Kriegeskorte & Douglas, 2018; Lewandowsky & Farrell, 2010; Pitt et al., 2002; Sun, 2008). These models can be conceptualized as formal mathematical translations of theoretical assumptions. These mathematical models can reveal internal, unobservable states that govern behavioral output (Baker et al., 2009; Baker & Tenenbaum, 2014; Gluck et al., 2010; Just et al., 1999; Wolpert et al., 2003). By translating cognitive processes into mathematical terms and testing them on an individual level, cognitive computational models have revealed individual differences in a wide array of decision-making contexts. Previous studies have differentiated decision-making of more risk averse and risk seeking individuals (Daw et al., 2011; Jacob et al., 2019; Levy, 2017; Pushkarskaya et al., 2017, 2018), impulsive versus more deliberate individuals who plan ahead (Blankenstein et al., 2017; Kable & Glimcher, 2007, 2010; Kurzban et al., 2013), or those that learn from environmental feedback itself or through imitation of others, versus individuals that are more likely to engage in metacognition and represent the task structure (Charpentier et al., 2016, 2017, 2020; Feher da Silva et al., 2023; Ramsey et al., 2021; Vélez & Gweon, 2021). Cognitive computational models can enhance our understanding of behavioral dynamics over time, enabling a more nuanced characterization of individual differences and providing a fundamentally dynamic perspective on cognitive variability within individuals (Schurr et al., 2024).

Computational modeling can inform the links between autistic traits and social functioning

With respect to autism specifically, recent research has highlighted the utility of computational modeling in studying autistic traits within the general population. This is especially important because autistic traits are known to exist on a continuum within the general population (Robinson et al., 2011). Some studies suggest that clinically relevant autistic traits are an extension (or end point) of that continuum (Constantino & Todd, 2003; Ronald & Hoekstra, 2011; Skuse et al., 2005), while other studies suggest a discontinuity between autistic traits in the general population and those of individuals with an autism diagnosis (Abu-Akel et al., 2019; Frazier et al., 2009; Peralta & Cuesta, 2007). Irrespective of these two opposing positions, exploring autistic traits in the general population provides several advantages and can critically inform research on ASD. This claim is warranted by the observation that nonclinical groups with high autistic traits exhibit a higher degree of social functioning (De Groot & Van Strien, 2017) than individuals with ASD. It is, therefore, possible to examine the cognitive profiles or behavioral strategies that contribute to greater social functioning despite high autistic traits. Exploring autistic traits in neurotypical individuals also makes it possible to avoid the confounding effects of comorbid conditions that co-occur with ASD. Furthermore, it enables researchers to study larger cohorts (De Groot & Van Strien, 2017), making it possible to control confounding effects or specifically examine their interaction with autistic traits as a question of interest. The Broad Autism Phenotype questionnaire (Hurley et al., 2007), for instance, describes behavioral and cognitive tendencies that are less severe but rather stable characteristics, similar in nature to those found in individuals with an ASD. Importantly, autistic traits seem to scale with the genetic risk for ASD. Relatives of individuals with ASD without a diagnosis have been shown to exhibit more autistic traits than those without a close relative with an ASD diagnosis (Piven et al., 1997).

Recent studies that have leveraged computational modeling better to define cognitive mechanisms that co-occur with high autistic traits have produced promising avenues for autism research. For example, one study used a hierarchical Bayesian modeling framework to examine the integration of nonsocial and social cues through a reward-based learning task. They found that more pronounced autistic traits in a group of healthy control subjects were related to less integration of social cues in decision-making. Computational modeling further demonstrated that performance differences between individuals with low versus high autistic traits were not due to an inability to process the social stimuli (gaze direction) and their causes, but rather to the extent to which participants relied on social information to infer the nonsocial cue (Sevgi et al., 2020). Another recent large-scale study found that high autistic traits were associated with reduced goal emulation during observational learning. This means that participants with higher autistic traits were more prone to imitate the observer but showed a reduced tendency to represent the overall goal of the observed person, which corresponded to the reward structure of the task (Wu et al., 2024). This emerging literature on social decision-making and autistic traits suggests that high autistic traits do not amount to a general inability to process social cues (Sevgi et al., 2020). They do, however, have a nuanced influence on the extent to which different types of social information are used, which may result in less adaptive outcomes.

Computational modeling can help to specify differences in the cognitive mechanisms underlying behavioral phenotypes

Cognitive computational models have also revealed differences in the cognitive mechanisms of individuals with an ASD diagnosis. In line with previous theories (Baron-Cohen et al., 1985), some studies have shown that individuals with ASD have difficulties in representing their partners' intentions. Yoshida and colleagues (Yoshida et al., 2010), for instance, employed a stag-hunt game to characterize unobservable computational processes implicit in social interactions and to measure whether individuals prefer smaller individual versus larger joint rewards. They found that the decisions of autistic individuals were less guided by inferring their partners' beliefs and instead were guided by a fixed strategy compared to non-autistic control participants. Autistic individuals who showed less mental state inference had greater symptom load. Similarly, a study on social learning showed that autistic adolescents relied less on social knowledge and feedback to learn

https://doi.org/10.1017/pen.2025.2 Published online by Cambridge University Press

about peers' preferences (Rosenblau et al., 2021). However, this study specifically tested whether autistic teens could learn about the preferences of non-autistic peers. Given that autistic and non-autistic adolescents may have differing preferences, these disparities could contribute to the "double empathy problem" – a phenomenon where misunderstandings between autistic and non-autistic individuals arise from mutual challenges in understanding each other's perspectives (Milton, 2012).

Other studies using economic exchange tasks have shown more nuanced differences in autistic individuals, which result in more adaptive behavior. For instance, autistic individuals have been shown to assess their partner's cooperation history more accurately and reciprocate less when partners are untrustworthy (Maurer et al., 2018). In accordance with this finding, a study on information sampling for cooperation showed that autistic adolescents had lower overall expectations (i.e., priors) about their partner's reciprocation tendencies (Liu et al., 2024). Given the range of trustworthy and untrustworthy agents, the autistic priors more accurately reflected the overall trustworthiness distribution of their potential partners. Moreover, autistic adolescents shared less often with untrustworthy agents than the non-autistic sample, which suggests that they are less prosocial than non-autistic adolescents. Moreover, this type of strategic interaction was less related to social skills in the autism group. Participants' task behaviors were less strongly associated with social skills as measured by the social responsiveness scale (SRS) compared to the non-autistic group, in which the SRS was the strongest predictor of cooperation tendencies. These findings suggest differences between strategic decision-making in economic exchange and non-economic social interactions in autistic individuals. In non-autistic groups, individuals' strategic choices may be more reflective of their mentalizing abilities and prosocial tendencies.

In conclusion, computational approaches offer a promising way to identify nuanced behavioral patterns and their underlying cognitive mechanisms. Studies leveraging both advances in building big data platforms and computational modeling hold promise for better characterizing autism phenotypes. This could pave the way for defining autism subtypes more clearly and examining their etiology, developmental trajectories, and comorbidities. Moreover, these advances can predict responses to therapeutic interventions, leading to more personalized and effective treatment plans (Collin et al., 2022; Johnson et al., 2021). Finally, we can combine insights from larger studies on autistic traits in the general population and those that investigate variability in symptoms in individuals diagnosed with ASD to inform commonalities and differences between populations with high autistic traits and no ASD diagnosis and those with a diagnosis. It could also help to systematically investigate the roles of sex, gender, and gender diversity in populations with high autistic traits and ASD diagnoses, given emerging evidence of the importance of examining sex differences in autism (Bölte et al., 2023; Loomes et al., 2017). This special issue focuses on how an individual differences approach, rather than the focus on central tendencies, can deepen our understanding of autistic traits and symptoms. It emphasizes the importance of investigating variability in autistic traits and phenotypes to refine autism classification. The issue highlights cutting-edge methods suited to capturing heterogeneity in behavioral and brain function and showcases big data approaches for analyzing large samples as well as computational modeling approaches that can expose how differences in cognitive mechanisms underlie meaningful behavioral variability.

References

Abu-Akel, A., Allison, C., Baron-Cohen, S., & Heinke, D. (2019). The distribution of autistic traits across the autism spectrum: Evidence for discontinuous dimensional subpopulations underlying the autism continuum. *Molecular Autism*, *10*, 24. https://doi.org/10.1186/s13229-019-0275-3

Adjerid, I., & Kelley, K. (2018). Big data in psychology: A framework for research advancement. *The American Psychologist*, *73*, 899–917. https://doi.org/10.1037/amp0000190

Amaral, D. G., Schumann, C. M., & Nordahl, C. W. (2008). Neuroanatomy of autism. *Trends in Neurosciences*, *31*, 137–145. https://doi.org/10.1016/j.tins.2007.12.005

Baker, C. L., Saxe, R., & Tenenbaum, J. B. (2009). Action understanding as inverse planning. *Cognition*, *113*, 329–349. https://doi.org/10.1016/j.cognition.2009.07.005

Baker, C. L., & Tenenbaum, J. B. (2014). Modeling human plan recognition using Bayesian theory of mind. *Plan, Activity, and Intent Recognition: Theory and Practice*, *7*, 177–204. https://doi.org/10.1016/B978-0-12-398532-3.00007-5

Baron-Cohen, S., Leslie, A. M., & Frith, U. (1985). Does the autistic child have a "theory of mind"? *Cognition*, *21*, 37–46. https://doi.org/10.1016/0010-0277(85)90022-8

Beauchaine, T. P., & Cicchetti, D. (2016). A new generation of comorbidity research in the era of neuroscience and Research Domain Criteria. *Development and psychopathology*, *28*, 891–894. https://doi.org/10.1017/S0954579416000602

Blankenstein, N. E., Peper, J. S., Crone, E. A., & van Duijvenvoorde, A. C. K. (2017). Neural mechanisms underlying risk and ambiguity attitudes. *Journal of Cognitive Neuroscience*, *29*, 1845–1859. https://doi.org/10.1162/jocn_a_01162

Bölte, S., Neufeld, J., Marschik, P. B., Williams, Z. J., Gallagher, L., & Lai, M. C. (2023). Sex and gender in neurodevelopmental conditions. *Nature Reviews. Neurology*, *19*, 136–159. https://doi.org/10.1038/S41582-023-00774-6

Bora, E., Aydın, A., Saraç, T., Kadak, M. T., & Köse, S. (2017). Heterogeneity of subclinical autistic traits among parents of children with autism spectrum disorder: Identifying the broader autism phenotype with a data-driven method. *Autism Research: Official Journal of the International Society for Autism Research*, *10*, 321–326. https://doi.org/10.1002/aur.1661

Castelfranchi, C., & Falcone, R. (2010). *Trust theory: A socio-cognitive and computational model*. John Wiley & Sons.

Charpentier, C. J., Aylward, J., Roiser, J. P., & Robinson, O. J. (2017). Enhanced risk aversion, but not loss aversion, in unmedicated pathological anxiety. *Biological Psychiatry*, *81*, 1014–1022. https://doi.org/10.1016/J.BIOPSYCH.2016.12.010

Charpentier, C. J., De Neve, J. E., Li, X., Roiser, J. P., & Sharot, T. (2016). Models of affective decision making: How do feelings predict choice? *Psychological Science*, *27*, 763–775. https://doi.org/10.1037/npe0000096

Charpentier, C. J., Iigaya, K., & O'Doherty, J. P. (2020). A Neuro-computational account of arbitration between choice Imitation and Goal Emulation during Human Observational Learning. *Neuron*, *106*, 687–699.e7. https://doi.org/10.1016/J.NEURON.2020.02.028

Collaboration, O. S. (2012). An open, large-scale, collaborative effort to estimate the reproducibility of psychological science. *Perspectives on Psychological Science*, *7*, 657–660. https://doi.org/10.1177/1745691612462588

Collaboration, O. S. (2015). Estimating the reproducibility of psychological science. *Science*, *349*, aac4716. https://doi.org/10.1126/science.aac4716

Collin, C. B., Gebhardt, T., Golebiewski, M., Karaderi, T., Hillemanns, M., Khan, F. M., Salehzadeh-Yazdi, A., Kirschner, M., Krobitsch, S., Consortium, E.-S. P., & Kuepfer, L. (2022). Computational models for clinical applications in personalized medicine – Guidelines and recommendations for data integration and model validation. *Journal of Personalized Medicine*, *12*, 166. https://doi.org/10.3390/jpm12020166

Constantino, J. N., & Todd, R. D. (2003). Autistic traits in the general population: A twin study. *Archives of General Psychiatry*, *60*, 524–530. https://doi.org/10.1001/archpsyc.60.5.524

Daw, N. D., Gershman, S. J., Seymour, B., Dayan, P., & Dolan, R. J. (2011). Model-based influences on humans' choices and striatal prediction errors. *Neuron*, *69*, 1204–1215. https://doi.org/10.1016/j.neuron.2011.02.027

https://doi.org/10.1017/pen.2025.2 Published online by Cambridge University Press

De Groot, K., & Van Strien, J. W. (2017). Evidence for a broad autism phenotype. *Advances in Neurodevelopmental Disorders*, *1*, 129–140. https://doi.org/10.1007/s41252-017-0021-9

Dillon, E. F., Kanne, S., Landa, R. J., Annett, R., Bernier, R., Bradley, C., Carpenter, L., Kim, S. H., Parish-Morris, J., Schultz, R., Wodka, E. L., & SPARK consortium (2023). Sex differences in autism: Examining intrinsic and extrinsic factors in children and adolescents enrolled in a national ASD cohort. *Journal of Autism and Developmental Disorders*, *53*, 1305–1318. https://doi.org/10.1007/s10803-021-05385-y

Eisenberg, I. W., Bissett, P. G., Zeynep Enkavi, A., Li, J., MacKinnon, D. P., Marsch, L. A., & Poldrack, R. A. (2019). Uncovering the structure of self-regulation through data-driven ontology discovery. *Nature Communications*, *10*, 2319. https://doi.org/10.1038/s41467-019-10301-1

Farrell, S., & Lewandowsky, S. (2018). *Computational modeling of cognition and behavior*. Cambridge University Press.

Feher da Silva, C., Lombardi, G., Edelson, M., & Hare, T. A. (2023). Rethinking model-based and model-free influences on mental effort and striatal prediction errors. *Nature Human Behaviour*, *7*, 956–969. https://doi.org/10.1038/s41562-023-01573-1

Fombonne, E., Green Snyder, L., Daniels, A., Feliciano, P., Chung, W., & SPARK Consortium (2020). Psychiatric and medical profiles of autistic adults in the SPARK cohort. *Journal of Autism and Developmental Disorders*, *50*, 3679–3698. https://doi.org/10.1007/s10803-020-04414-6

Frässle, S., Yao, Y., Schöbi, D., Aponte, E. A., Heinzle, J., & Stephan, K. E. (2018). Generative models for clinical applications in computational psychiatry. *WIREs Cognitive science*, *9*, e1460. https://doi.org/10.1002/wcs.1460

Frazier, T. W., Youngstrom, E. A., Sinclair, L., Kubu, C. S., Law, P., Rezai, A., Constantino, J. N., & Eng, C. (2009). Autism spectrum disorders as a qualitatively distinct category from typical behavior in a large, clinically ascertained sample. *Assessment*, *17*, 308–320. https://doi.org/10.1177/1073191109356534

Friston, K. J., Stephan, K. E., Montague, R., & Dolan, R. J. (2014). Computational psychiatry: The brain as a phantastic organ. *Lancet Psychiatry*, *1*, 148–158. https://doi.org/10.1016/S2215-0366(14)70275-5

Frolichs, K. M. M., Rosenblau, G., & Korn, C. W. (2022). Incorporating social knowledge structures into computational models. *Nature Communications*, *13*, 6205. https://doi.org/10.1038/s41467-022-33418-2

Gaugler, T., Klei, L., Sanders, S. J., Bodea, C. A., Goldberg, A. P., Lee, A. B., Mahajan, M., Manaa, D., Pawitan, Y., Reichert, J., Ripke, S., Sandin, S., Sklar, P., Svantesson, O., Reichenberg, A., Hultman, C. M., Devlin, B., Roeder, K., & Buxbaum, J. D. (2014). Most genetic risk for autism resides with common variation. *Nature Genetics*, *46*, 881–885. https://doi.org/10.1038/ng.3039

Geschwind, D. H., & Levitt, P. (2007). Autism spectrum disorders: Developmental disconnection syndromes. *Current Opinion in Neurobiology*, *17*, 103–111. https://doi.org/10.1016/j.conb.2007.01.009

Gluck, K., Stanley, C., Moore, L., Reitter, D. & Halbrügge, M. (2010), Exploration for understanding in cognitive modeling. *Journal of Artificial General Intelligence*, *2*, 88–107. https://doi.org/10.2478/v10229-011-0011-7

Gomez-Marin, A., Paton, J. J., Kampff, A. R., Costa, R. M., & Mainen, Z. F. (2014). Big behavioral data: Psychology, ethology and the foundations of neuroscience. *Nature Neuroscience*, *17*, 1455–1462. https://doi.org/10.1038/nn.3812

Grove, J., Ripke, S., Als, T. D., Mattheisen, M., Walters, R. K., Won, H., Pallesen, J., Agerbo, E., Andreassen, O. A., Anney, R., Awashti, S., Belliveau, R., Bettella, F., Buxbaum, J. D., Bybjerg-Grauholm, J., Bækvad-Hansen, M., Cerrato, F., Chambert, K., Christensen, J. H., Churchhouse, C., … Børglum, A. D. (2019). Identification of common genetic risk variants for autism spectrum disorder. *Nature Genetics*, *51*, 431–444. https://doi.org/10.1038/s41588-019-0344-8

Haehner, P., Rakhshani, A., Fassbender, I., Lucas, R. E., Donnellan, M. B., & Luhmann, M. (2023). Perception of major life events and personality trait change. *European Journal of Personality*, *37*, 524–542. https://doi.org/10.1177/08902070221107973

Harlow, L. L., & Oswald, F. L. (2016). Big data in psychology: Introduction to the special issue. *Psychological Methods*, *21*, 447–457. https://doi.org/10.1037/met0000120

Hedge, C., Powell, G., & Sumner, P. (2018). The reliability paradox: Why robust cognitive tasks do not produce reliable individual differences. *Behavior Research Methods*, *50*, 1166–1186. https://doi.org/10.3758/s13428-017-0935-1

Heinsfeld, A. S., Franco, A. R., Craddock, R. C., Buchweitz, A., & Meneguzzi, F. (2017). Identification of autism spectrum disorder using deep learning and the ABIDE dataset. *NeuroImage. Clinical*, *17*, 16–23. https://doi.org/10.1016/j.nicl.2017.08.017

Hitchcock, P. F., Fried, E. I., & Frank, M. J. (2022). Computational psychiatry needs time and context. *Annual Review of Psychology*, *73*, 243–270. https://doi.org/10.1146/annurev-psych-021621-124910

Hurley, R. S., Losh, M., Parlier, M., Reznick, J. S., & Piven, J. (2007). The broad autism phenotype questionnaire. *Journal of Autism and Developmental Disorders*, *37*, 1679–1690. https://doi.org/10.1007/s10803-006-0299-3

Huys, Q. J., Maia, T. V., & Frank, M. J. (2016). Computational psychiatry as a bridge from neuroscience to clinical applications. *Nature Neuroscience*, *19*, 404–413. https://doi.org/10.1038/nn.4238

Ilmini, W. M. K. S., & Fernando, T. G. I. (2017). Computational personality traits assessment: A review. *IEEE International Conference on Industrial and Information Systems, ICIIS 2017 - Proceedings*, *2018-January*, 1–6. https://doi.org/10.1109/ICIINFS.2017.8300416

Insel, T., Cuthbert, B., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., Sanislow, C., & Wang, P. (2010). Research domain criteria (RDoC): Toward a new classification framework for research on mental disorders. *The American Journal of Psychiatry*, *167*, 748–751. https://doi.org/10.1176/appi.ajp.2010.09091379

Jacob, S., Wolff, J. J., Steinbach, M. S., Doyle, C. B., Kumar, V., & Elison, J. T. (2019). Neurodevelopmental heterogeneity and computational approaches for understanding autism. *Translational Psychiatry*, *9*, 63. https://doi.org/10.1038/s41398-019-0390-0

Jauk, E., & Kanske, P. (2019). Perspective change and personality state variability: An argument for the role of self-awareness and an outlook on bidirectionality (Commentary on Wundrack et al., 2018). *Journal of Intelligence*, *7*, 10. https://doi.org/10.3390/jintelligence7020010

Johnson, K. B., Wei, W.-Q., Weeraratne, D., Frisse, M. E., Misulis, K., Rhee, K., Zhao, J., & Snowdon, J. L. (2021). Precision medicine, AI, and the future of personalized health care. *Clinical and Translational Science*, *14*, 86–93. https://doi.org/10.1111/cts.12884

Just, M. A., Carpenter, P. A., & Varma, S. (1999). Computational modeling of high-level cognition and brain function. *Human Brain Mapping*, *8*, 128–136. https://doi.org/10.1002/(SICI)1097-0193(1999)8:2/3<128::AID-HBM10>3.0.CO;2-G

Kable, J. W., & Glimcher, P. W. (2007). The neural correlates of subjective value during intertemporal choice. *Nature Neuroscience*, *10*, 1625–1633. https://doi.org/10.1038/nn2007

Kable, J. W., & Glimcher, P. W. (2010). An "as soon as possible" effect in human intertemporal decision making: Behavioral evidence and neural mechanisms. *Journal of Neurophysiology*, *103*, 2513–2531. https://doi.org/10.1152/JN.00177.2009/ASSET/IMAGES/LARGE/Z9K0051000750009.JPEG

Karvelis, P., Paulus, M. P., & Diaconescu, A. O. (2023). Individual differences in computational psychiatry: A review of current challenges. *Neuroscience and Biobehavioral Reviews*, *148*, 105137. https://doi.org/10.1016/j.neubiorev.2023.105137

King-Casas, B., Sharp, C., Lomax-Bream, L., Lohrenz, T., Fonagy, P., & Montague, P. R. (2008). The rupture and repair of cooperation in borderline personality disorder. *Science*, *321*, 806–810. https://doi.org/10.1126/science.1156902

Kravitz, D. J., & Mitroff, S. R. (2023). Quantifying, and correcting for, the impact of questionable research practices on false discovery rates in psychological science. *Journal for Reproducibility in Neuroscience*. https://doi.org/0.36850/jrn.2023.e44

Kriegeskorte, N., & Douglas, P. K. (2018). Cognitive computational neuroscience. *Nature Neuroscience*, *21*, 1148–1160. https://doi.org/10.1038/s41593-018-0210-5

Kurzban, R., Duckworth, A., Kable, J. W., & Myers, J. (2013). An opportunity cost model of subjective effort and task performance. *Behavioral and Brain Sciences*, *36*, 661–679. https://doi.org/10.1017/S0140525X12003196

Levy, I. (2017). Neuroanatomical substrates for risk behavior. *The Neuroscientist*, *23*, 275–286. https://doi.org/10.1177/1073858416672414

Lewandowsky, S., & Farrell, S. (2010). *Computational modeling in cognition: Principles and practice*. SAGE Publications.

Liu, W., Shah, N., Ma, I., & Rosenblau, G. (2024). Strategic social decision making undergoes significant changes in typically developing and autistic early adolescents. *Developmental Science*, *27*, e13463. https://doi.org/10.1111/desc.13463

Lloyd-Cox, J., Pickering, A. D., Beaty, R. E., & Bhattacharya, J. (2023). Toward greater computational modeling in neurocognitive creativity research. *Psychology of Aesthetics, Creativity, and the Arts*, advance online publication. https://doi.org/10.1037/ACA0000627

Lochner, C., & Stein, D. J. (2003). Heterogeneity of obsessive-compulsive disorder: A literature review. *Harvard Review of Psychiatry*, *11*(3), 113–132. https://doi.org/10.1080/10673220303949

Loomes, R., Hull, L., & Mandy, W. P. L. (2017). What is the male-to-female ratio in autism spectrum disorder? A systematic review and meta-analysis. *Journal of the American Academy of Child & Adolescent Psychiatry*, *56*, 466–474. https://doi.org/10.1016/J.JAAC.2017.03.013

Matoba, N., Liang, D., Sun, H., Aygün, N., McAfee, J. C., Davis, J. E., Raffield, L. M., Qian, H., Piven, J., Li, Y., Kosuri, S., Won, H., & Stein, J. L. (2020). Common genetic risk variants identified in the SPARK cohort support DDHD2 as a candidate risk gene for autism. *Translational Psychiatry*, *10*, 265. https://doi.org/10.1038/s41398-020-00953-9

Maurer, C., Chambon, V., Bourgeois-Gironde, S., Leboyer, M., & Zalla, T. (2018). The influence of prior reputation and reciprocity on dynamic trust-building in adults with and without autism spectrum disorder. *Cognition*, *172*, 1–10. https://doi.org/10.1016/J.COGNITION.2017.11.007

Mejia, C., D'Ippolito, B., & Kajikawa, Y. (2021). Major and recent trends in creativity research: An overview of the field with the aid of computational methods. *Creativity and Innovation Management*, *30*, 475–497. https://doi.org/10.1111/CAIM.12453

Milton, D. E. M. (2012). On the ontological status of autism: The 'double empathy problem.' *Disability & Society*, *27*, 883–887. https://doi.org/10.1080/09687599.2012.710008

Minai, A. A., Doboli, S., & Iyer, L. R. (2021). Models of creativity and ideation: An overview. *Understanding Complex Systems*, 21–45. https://doi.org/10.1007/978-3-030-77198-0_2

Molenaar, P. C. M. (2004). A manifesto on psychology as idiographic science: Bringing the person back into scientific psychology, this time forever. *Measurement: Interdisciplinary Research and Perspectives*, *2*, 201–218. https://doi.org/10.1207/s15366359mea0204_1

Montague, P. R., Dolan, R. J., Friston, K. J., & Dayan, P. (2012). Computational psychiatry. *Trends in Cognitive Sciences*, *16*, 72–80. https://doi.org/10.1016/j.tics.2011.11.018

Myers, S. M., Challman, T. D., Bernier, R., Bourgeron, T., Chung, W. K., Constantino, J. N., Eichler, E. E., Jacquemont, S., Miller, D. T., Mitchell, K. J., Zoghbi, H. Y., Martin, C. L., & Ledbetter, D. H. (2020). Insufficient Evidence for "Autism-Specific" Genes. *American Journal of Human Genetics*, *106*, 587–595. https://doi.org/10.1016/j.ajhg.2020.04.004

Nesselroade, J. R. (2004). Interindividual differences in intraindividual change. *Best Methods for the Analysis of Change: Recent Advances, Unanswered Questions, Future Directions*, 92–105. https://doi.org/10.1037/10099-006

Nielsen, J. A., Zielinski, B. A., Fletcher, P. T., Alexander, A. L., Lange, N., Bigler, E. D., Lainhart, J. E., & Anderson, J. S. (2013). Multisite functional connectivity MRI classification of autism: ABIDE results. *Frontiers in Human Neuroscience*, *7*, 599. https://doi.org/10.3389/fnhum.2013.00599

Patzelt, E. H., Hartley, C. A., & Gershman, S. J. (2018). Computational phenotyping: Using models to understand individual differences in personality, development, and mental illness. *Personality Neuroscience*, *1*, e18. https://doi.org/10.1017/PEN.2018.14

Payakachat, N., Tilford, J. M., & Ungar, W. J. (2016). National database for autism research (NDAR): Big data opportunities for health services research and health technology assessment. *PharmacoEconomics*, *34*, 127–138. https://doi.org/10.1007/s40273-015-0331-6

Pelphrey, K. A., Shultz, S., Hudac, C. M., & Vander Wyk, B. C. (2011). Research review: Constraining heterogeneity: The social brain and its development in autism spectrum disorder. *Journal of Child Psychology and Psychiatry*, *52*, 631–644. https://doi.org/10.1111/J.1469-7610.2010.02349.X

Peralta, V., & Cuesta, M. J. (2007). A dimensional and categorical architecture for the classification of psychotic disorders. *World Psychiatry: Official Journal of the World Psychiatric Association (WPA)*, *6*, 100–101.

Petzschner, F. H., Weber, L. A. E., Gard, T., & Stephan, K. E. (2017). Computational psychosomatics and computational psychiatry: Toward a joint framework for differential diagnosis. *Biological Psychiatry*, *82*, 421–430. https://doi.org/10.1016/j.biopsych.2017.05.012

Phan, L. V., & Rauthmann, J. F. (2021). Personality computing: New frontiers in personality assessment. *Social and Personality Psychology Compass*, *15*, e12624. https://doi.org/10.1111/spc3.12624

Pitt, M. A., Myung, I. J., & Zhang, S. (2002). Toward a method of selecting among computational models of cognition. *Psychological Review*, *109*, 472–491. https://doi.org/10.1037/0033-295x.109.3.472

Piven, J., Palmer, P., Jacobi, D., Childress, D., & Arndt, S. (1997). Broader autism phenotype: Evidence from a family history study of multiple-incidence autism families. *The American Journal of Psychiatry*, *154*, 185–190. https://doi.org/10.1176/ajp.154.2.185

Pushkarskaya, H., Tolin, D. F., Henick, D., Levy, I., & Pittenger, C. (2018). Unbending mind: Individuals with hoarding disorder do not modify decision strategy in response to feedback under risk. *Psychiatry Research*, *259*, 506–513. https://doi.org/10.1016/j.psychres.2017.11.001

Pushkarskaya, H., Tolin, D., Ruderman, L., Henick, D., Kelly, J. M. L., Pittenger, C., & Levy, I. (2017). Value-based decision making under uncertainty in hoarding and obsessive- compulsive disorders. *Psychiatry Research*, *258*, 305–315. https://doi.org/10.1016/j.psychres.2017.08.058

Ramsey, R., Kaplan, D. M., & Cross, E. S. (2021). Watch and learn: The cognitive neuroscience of learning from others' actions. *Trends in Neurosciences*, *44*, 478–491. https://doi.org/10.1016/J.TINS.2021.01.007

Riding, R., & Rayner, S. (2013). *Cognitive styles and learning strategies: Understanding style differences in learning and behavior*. David Fulton Publishers.

Robinson, E. B., Koenen, K. C., McCormick, M. C., Munir, K., Hallett, V., Happé, F., Plomin, R., & Ronald, A. (2011). Evidence that autistic traits show the same etiology in the general population and at the quantitative extremes (5%, 2.5%, and 1%). *Archives of General Psychiatry*, *68*, 1113–1121. https://doi.org/10.1001/archgenpsychiatry.2011.119

Ronald, A., & Hoekstra, R. A. (2011). Autism spectrum disorders and autistic traits: A decade of new twin studies. *American Journal of Medical Genetics. Part B, Neuropsychiatric Genetics*, *156B*, 255–274. https://doi.org/10.1002/ajmg.b.31159

Rosenblau, G., Korn, C. W., Dutton, A., Lee, D., & Pelphrey, K. A. (2021). Neurocognitive mechanisms of social inferences in typical and autistic adolescents. *Biological Psychiatry. Cognitive Neuroscience and Neuroimaging*, *6*, 782–791. https://doi.org/10.1016/j.bpsc.2020.07.002

Rozin, P. (2001). Social psychology and science: Some lessons from Solomon Asch. *Personality and Social Psychology Review*, *5*, 2–14. https://doi.org/10.1207/S15327957PSPR0501_1

Ruzich, E., Allison, C., Smith, P., Watson, P., Auyeung, B., Ring, H., & Baron-Cohen, S. (2016). Subgrouping siblings of people with autism: Identifying the broader autism phenotype. *Autism research*, *9*, 658–665. https://doi.org/10.1002/aur.1544

Saré, R. M., & Smith, C. B. (2020). Association between sleep deficiencies with behavioral problems in autism spectrum disorder: Subtle sex differences. *Autism Research*, *13*, 1629–1822.. https://doi.org/10.1002/aur.2396

Saunders, R., & Bown, O. (2015). Computational social creativity. *Artificial Life*, *21*, 366–378. https://doi.org/10.1162/ARTL_a_00177

Schurr, R., Reznik, D., Hillman, H., Bhui, R., & Gershman, S. J. (2024). Dynamic computational phenotyping of human cognition. *Nature Human Behaviour*, *8*, 917–931. https://doi.org/10.1038/s41562-024-01814-x

Sevgi, M., Diaconescu, A. O., Henco, L., Tittgemeyer, M., & Schilbach, L. (2020). Social Bayes: Using Bayesian modeling to study autistic trait-related differences in social cognition. *Biological Psychiatry*, *87*, 185–193. https://doi.org/10.1016/j.biopsych.2019.09.032

Siritzky, E. M., Cox, P. H., Nadler, S. M., Grady, J. N., Kravitz, D. J., & Mitroff, S. R. (2023). Standard experimental paradigm designs and data exclusion

practices in cognitive psychology can inadvertently introduce systematic "shadow" biases in participant samples. *Cognitive Research: Principles and Implications*, *8*, 66. https://doi.org/10.1186/s41235-023-00520-y

Skuse, D. H., Mandy, W. P., & Scourfield, J. (2005). Measuring autistic traits: Heritability, reliability and validity of the Social and Communication Disorders Checklist. *The British Journal of Psychiatry*, *187*, 568–572. https://doi.org/10.1192/bjp.187.6.568

Steinhausen H. C. (2009). The heterogeneity of causes and courses of attention-deficit/hyperactivity disorder. *Acta Psychiatrica Scandinavica*, *120*, 392–399. https://doi.org/10.1111/j.1600-0447.2009.01446.x

Stephan, K. E., & Mathys, C. (2014). Computational approaches to psychiatry. *Current Opinion in Neurobiology*, *25*, 85–92. https://doi.org/10.1016/j.conb.2013.12.007

Subramanian, R., Yan, Y., Staiano, J., Lanz, O., & Sebe, N. (2013). On the relationship between head pose, social attention and personality prediction for unstructured and dynamic group interactions. *Proceedings of the 15th ACM on International Conference on Multimodal Interaction*, 3–10. https://doi.org/10.1145/2522848.2522862

Sun, R. (2008). Introduction to computational cognitive modeling. In R. Sun (Ed.), *The Cambridge Handbook of Computational Psychology* (pp. 3–19). Cambridge University Press. https://doi.org/10.1017/CBO9780511816772.003

Tiego, J., Martin, E. A., DeYoung, C. G., Hagan, K., Cooper, S. E., Pasion, R., Satchell, L., Shackman, A. J., Bellgrove, M. A., Fornito, A., & HiTOP Neurobiological Foundations Work Group (2023). Precision behavioral phenotyping as a strategy for uncovering the biological correlates of psychopathology. *Nature. Mental Health*, *1*, 304–315. https://doi.org/10.1038/s44220-023-00057-5

Vélez, N., & Gweon, H. (2021). Learning from other minds: An optimistic critique of reinforcement learning models of social learning. *Current Opinion in Behavioral Sciences*, *38*, 110–115. https://doi.org/10.1016/j.cobeha.2021.01.006

Wang, X. J., & Krystal, J. H. (2014). Computational psychiatry. *Neuron*, *84*, 638–654. https://doi.org/10.1016/j.neuron.2014.10.018

Wiecki, T. V., Poland, J., & Frank, M. J. (2015). Model-based cognitive neuroscience approaches to computational psychiatry: Clustering and classification. *Clinical Psychological Science*, *3*, 378–399. https://doi.org/10.1177/2167702614565359

Wilfert, A. B., Turner, T. N., Murali, S. C., Hsieh, P., Sulovari, A., Wang, T., Coe, B. P., Guo, H., Hoekzema, K., Bakken, T. E., Winterkorn, L. H., Evani, U. S., Byrska-Bishop, M., Earl, R. K., Bernier, R. A., SPARK Consortium, Zody, M. C., & Eichler, E. E. (2021). Recent ultra-rare inherited variants implicate new autism candidate risk genes. *Nature Genetics*, *53*, 1125–1134. https://doi.org/10.1038/s41588-021-00899-8

Wolfers, T., Floris, D. L., Dinga, R., van Rooij, D., Isakoglou, C., Kia, S. M., Zabihi, M., Llera, A., Chowdanayaka, R., Kumar, V. J., Peng, H., Laidi, C., Batalle, D., Dimitrova, R., Charman, T., Loth, E., Lai, M. C., Jones, E., Baumeister, S., Moessnang, C., ... **Beckmann, C. F.** (2019). From pattern classification to stratification: Towards conceptualizing the heterogeneity of Autism Spectrum Disorder. *Neuroscience and Biobehavioral Reviews*, *104*, 240–254. https://doi.org/10.1016/j.neubiorev.2019.07.010

Wolff, J. J., Jacob, S., & Elison, J. T. (2018). The journey to autism: Insights from neuroimaging studies of infants and toddlers. *Development and Psychopathology*, *30*, 479–495. https://doi.org/10.1017/S0954579417000980

Wolpert, D. M., Doya, K., & Kawato, M. (2003). A unifying computational framework for motor control and social interaction. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *358*, 593–602. https://doi.org/10.1098/rstb.2002.1238

Wu, Q., Oh, S., Tadayonnejad, R., Feusner, J. D., Cockburn, J., O'Doherty, J. P., & Charpentier, C. J. (2024). Individual differences in autism-like traits are associated with reduced goal emulation in a computational model of observational learning. *Nature. Mental health*, *2*, 1032–1044. https://doi.org/10.1038/s44220-024-00287-1

Yoshida, W., Dziobek, I., Kliemann, D., Heekeren, H. R., Friston, K. J., & Dolan, R. J. (2010). Cooperation and heterogeneity of the autistic mind. *Journal of Neuroscience*, *30*, 8815–8818. https://doi.org/10.1523/JNEUROSCI.0400-10.2010

Zhang, L. (2002). Measuring thinking styles in addition to measuring personality traits? *Personality and Individual Differences*, *33*, 445–458. https://doi.org/10.1016/S0191-8869(01)00166-0

Zhu, H., Li, L., Zhao, S., & Jiang, H. (2018). Evaluating attributed personality traits from scene perception probability. *Pattern Recognition Letters*, *116*, 121–126. https://doi.org/10.1016/j.patrec.2018.09.027

https://doi.org/10.1017/pen.2025.2 Published online by Cambridge University Press