J. London Math. Soc. (2) 71 (2005) 304–320 © 2005 London Mathematical Society
doi:10.1112/S0024610705006290

PURE-PROJECTIVE MODULES

GENA PUNINSKI AND PHILIPP ROTHMALER

Abstract

The question is addressed of when all pure-projective modules are direct sums of finitely presented modules. It is proved that this is the case over hereditary noetherian rings. Partial results are obtained for uniserial rings. Some of the methods are model-theoretic, and the techniques developed using these may be of interest in their own right.

1. *Introduction*

It is well known that a module is pure-projective if and only if it is a direct summand of a direct sum of finitely presented modules. Our principal concern is the question of when all pure-projective modules *are* direct sums of finitely presented modules (or, equivalently, of finitely generated modules, for finitely generated pure-projectives are finitely presented). This is a 'pure' version of the classical question of when projectives are free. By a theorem of Maranda, the answer is yes for abelian groups, that is, over the ring of integers; see [**7**, Theorem 30.2].

If the ring is Krull–Schmidt in the sense of [**8**], that is, when every finitely presented module is a direct sum of modules with a local endomorphism ring, then the Krull–Remak–Schmidt–Azumaya theorem implies that, again, the answer is yes. All right noetherian, (two-sided) serial rings are examples of such rings (by Fact 5.1, which holds for serial rings as well, and [**5**, Proposition 9.24] or [**12**, Lemma 2.11]).

Our development of the decomposition theory of pure-projective modules can be outlined as follows.

After recollection of some model-theoretic preliminaries in Section 2, we introduce pure-projectives and develop some of their model theory in Section 3, on the basis of on the second author's description of countably generated pure-projective modules, Fact 3.4. The main result in this section is what we call the telescoping map theorem, Theorem 3.7.

As a first application of this machinery, we give an easy proof of the Bass–Björk result, Theorem 4.2, that if every flat right R-module is projective, then every left R-module has the descending chain condition (DCC) on finitely generated submodules.

In Section 5 we deal with uniserial rings. The decomposition theory of pure-projectives is little understood there. Even over a uniserial domain R it is possible to have an indecomposable pure-projective module which is not finitely generated. In [**12**, proof of Lemma 14.21], such a module was constructed in the category $\mathrm{Add}(M)$ of a *cyclically presented* module M, that is, of a module of the form R/rR, where $r \in R$. Here, $\mathrm{Add}(M)$ denotes the smallest additive category containing

Received 2 March 2004.

2000 *Mathematics Subject Classification* 16D70, 16G30, 16D40, 16P40, 13L05, 16B70, 16U10, 16U20.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

M (whose objects are the direct summands of direct sums of copies of M). We are able, however, to answer our original question within this restricted category: we completely determine when every (pure-projective) module in $\mathrm{Add}(R/rR)$ is a direct sum of finitely generated modules. This is the same as determining when every projective module over the ring $S = \mathrm{End}(R/rR)$ is free; see Theorem 5.8.

In Section 6, we answer in the positive our original question for (two-sided) hereditary noetherian rings by showing that every pure-projective module over such a ring is a direct sum of finitely generated modules (Corollary 6.5). The heart of the proof is the special case of hereditary noetherian prime rings (Theorem 6.4). This section does not use the machinery developed earlier, but is rather based on a general result, Lemma 6.3, which states that, over a semihereditary semiprime Goldie ring, pure-projectives are torsion-splitting.

Unless stated otherwise, all our modules are right modules over an associative ring R with 1, that is, we work in the category Mod-R. All our unexplained terminology is standard and can be found in one of the texts cited. (In particular, as for ring-theoretic terms, when no side is specified, they mean either side; for example a hereditary noetherian ring is a two-sided hereditary, two-sided noetherian ring.)

2. *Free realizations*

Basic definitions from the model theory of modules can be found in [**11**]. Below we recall a few of them.

As is common, by a *tuple* we mean a finite sequence of elements, usually denoted by a lower-case letter with a bar over. Slightly abusing notation, we write $\overline{m} \in M$ to mean that each entry of the tuple $\overline{m}$ is in M.

By a *pointed module* we mean a pair $(M, \overline{m})$ where M is a module and $\overline{m} \in M$ is a tuple of elements from M. A *morphism* of pointed modules (or a *pointed morphism*), $f : (M, \overline{m}) \longrightarrow (N, \overline{n})$, is a homomorphism $f : M \longrightarrow N$ such that $f(\overline{m}) = \overline{n}$ coordinatewise (in particular, $\overline{m}$ and $\overline{n}$ have to have the same length).

Recall that a *pp-formula* is a formula of the form $\exists\, \overline{y}\ (\overline{y}A = \overline{x}B)$, where $\overline{x}$, $\overline{y}$ are tuples of variables, and A,B are rectangular matrices over the ring of appropriate size. We often use the shorthand $A \,|\, \overline{x}B$ (read 'A divides $\overline{x}B$') for this formula. For instance $a \,|\, x$ stands for $\exists\, y\ (ya = x)$.

Given a pointed module $(M, \overline{m})$ and a pp-formula $\varphi(\overline{x})$ of the form $\exists\, \overline{y}\ (\overline{y}A = \overline{x}B)$, one writes $M \models \varphi(\overline{m})$ (and states that $\overline{m}$ satisfies φ in M) if there is a tuple $\overline{n} \in M$ such that $\overline{n}A = \overline{m}B$. For instance, $M \models (a \,|\, x)(m)$ if and only if $m \in Ma$.

We write $\varphi \longrightarrow \psi$ if a tuple in a module satisfies ψ whenever it satisfies φ. (Therefore this is a statement about the entire category of all right R-modules.)

The *pp-type* of $\overline{m}$ in M, denoted $pp_M(\overline{m})$, is the set of all pp-formulae satisfied by the tuple $\overline{m}$ in M.

It is easily seen that homomorphisms preserve pp-formulae. If $f : M \longrightarrow N$ and $M \models \varphi(\overline{m})$, then $N \models \varphi(f(\overline{m}))$; hence also $pp_M(\overline{m}) \subseteq pp_N(f(\overline{m}))$. In particular, if $f : (M, \overline{m}) \longrightarrow (N, \overline{n})$ is a pointed morphism and $M \models \varphi(\overline{m})$, then $N \models \varphi(\overline{n})$; hence also $pp_M(\overline{m}) \subseteq pp_N(\overline{n})$.

There are two well known cases where the converse is true. (Another one is dealt with in Lemma 3.3.)

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

FACT 2.1 [**11**, Proposition 8.5, Theorem 2.8]. *Let M be finitely presented or N be pure-injective. Suppose that $\overline{m} \in M$ and $\overline{n} \in N$ are such that $pp_M(\overline{m}) \subseteq pp_N(\overline{n})$. Then there exists a pointed morphism $f : (M, \overline{m}) \longrightarrow (N, \overline{n})$.*

A pointed module $(M, \overline{m})$ is said to be a *free realization* of a pp-formula $\varphi(\overline{x})$ if (a) $M \models \varphi(\overline{m})$, and (b) if $N \models \varphi(\overline{n})$ for some tuple $\overline{n}$ in a module N, then there is a pointed morphism $(M, \overline{m}) \longrightarrow (N, \overline{n})$. For instance, if a is an element of the ring R, then (R, a) is a free realization of the formula $a \,|\, x$, and $(R/aR, 1 + aR)$ is a free realization of $xa = 0$. Both of these are *finitely presented* in the sense that the underlying module is.

Free realizations are never unique: one can always add arbitrary modules or take pure injective envelopes without disturbing conditions (a) and (b) above. In general, there is no 'minimal' free realization either.

However, as the following shows, all pp-formulae have a free realization, even a 'small' one.

FACT 2.2 [**11**, Proposition 8.4]. *Every pp-formula has a finitely presented free realization.*

The pp-type $p = pp_M(\overline{m})$ is called *finitely generated* if there exists a formula $\varphi \in p$ such that $\varphi \longrightarrow \psi$ for every $\psi \in p$. In this case φ is said to *generate* p. It is easy to verify that for every free realization $(M, \overline{m})$ (of some pp-formula φ), the pp-type $pp_M(\overline{m})$ is finitely generated (by φ).

It is also not hard to prove that any pointed module $(N, \overline{n})$ with underlying finitely presented module N is a free realization of some pp-formula. This yields the following.

FACT 2.3 [**11**, Proposition 8.4]. *Every pp-type in a finitely presented module is finitely generated.*

In fact one can pinpoint the generating formula precisely. Let M be given by generators $\overline{n}$ and relations $\overline{n}A = 0$, and let $\overline{m} = \overline{n}B$. Then $pp_M(\overline{m})$ is generated by the formula $\exists\, \overline{y}\ (\overline{y}B = \overline{x} \wedge \overline{y}A = 0)$. This can be generalized as follows.

FACT 2.4 [**16**, Lemma 1.5]. *Suppose that $(M, \overline{m})$ is a pointed module such that $pp_M(\overline{m})$ is finitely generated, and $\overline{n}$ is contained in the submodule generated by $\overline{m}$. Then $pp_M(\overline{n})$ is finitely generated.*

More precisely, if $\overline{n} = \overline{m}A$ and $pp_M(\overline{m})$ is generated by $\varphi(\overline{x})$, then $pp_M(\overline{n})$ is generated by $\exists\, \overline{y}\ (\overline{y}A = \overline{x} \wedge \varphi(\overline{y}))$.

Proof. Let $\psi \in pp_M(\overline{n})$, and denote the formula $\psi(\overline{x}A)$ by $\delta(\overline{x})$. The latter formula is in $pp_M(\overline{m})$; hence $\varphi \longrightarrow \delta$. To prove $\exists\, \overline{y}\ (\overline{y}A = \overline{x} \wedge \varphi(\overline{y})) \longrightarrow \psi(\overline{x})$, let $\overline{a}$ satisfy $\exists\, \overline{y}\ (\overline{y}A = \overline{x} \wedge \varphi(\overline{y}))$ in some module N. We have to show that it also satisfies ψ. Choose witnesses $\overline{b}$ satisfying φ such that $\overline{b}A = \overline{a}$. As $\varphi \longrightarrow \delta$, the tuple $\overline{b}$ also satisfies δ, that is, $\overline{a} = \overline{b}A$ satisfies ψ, as desired. □

We conclude with an algebraic description of finite generation of pp-types.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

LEMMA 2.5. *Let $(M, \overline{m})$ be a pointed module. Then the following are equivalent.*

(1) *$p = pp_M(\overline{m})$ is finitely generated.*

(2) *There is a finitely presented pointed module $(N, \overline{n})$ and a 'universal' pointed morphism $f : (N, \overline{n}) \longrightarrow (M, \overline{m})$, that is, given a finitely presented pointed module $(K, \overline{k})$ and a pointed morphism $g : (K, \overline{k}) \longrightarrow (M, \overline{m})$, the diagram in Figure 1 can be completed as shown.*




FIGURE 1.

Proof. (1) $\Rightarrow$ (2): Let φ generate p. Pick a finitely presented free realization $(N, \overline{n})$ of φ and let f be the corresponding map.

We know that $pp_K(\overline{k})$ is generated by a certain $\psi(\overline{x})$. Applying g, we obtain $\psi \in p$, and therefore $\varphi \longrightarrow \psi$; hence also $pp_K(\overline{k}) \subseteq pp_N(\overline{n})$. Thus the desired h exists by Fact 2.1.

(2) $\Rightarrow$ (1): Suppose that there is a universal pointed morphism $f : (N, \overline{n}) \longrightarrow (M, \overline{m})$. Let φ generate $pp_N(\overline{n})$. We claim that φ generates p. Applying f, we obtain $\varphi \in p$.

Suppose that $\psi \in p$. Let $(K, \overline{k})$ be a free realization of ψ and $g : (K, \overline{k}) \longrightarrow (M, \overline{m})$ be the corresponding map. By the universal property, g factors through f via a pointed morphism $h : (K, \overline{k}) \longrightarrow (N, \overline{n})$. Applying h to $K \models \psi(\overline{k})$, we obtain $N \models \psi(\overline{n})$. Thus $\varphi \longrightarrow \psi$. □

3. *Pure-projective modules*

A module is said to be *pure-projective* if it is projective with respect to pure-exact sequences. Warfield [**20**] showed that a module is pure-projective if and only if it is a direct summand of a direct sum of finitely presented modules, cf. [**21**, 33.6]. In other words, the class of pure-projective right R-modules is precisely the class of objects in Add(mod-R) (where, as usual, mod-R stands for the full subcategory of Mod-R of finitely presented right R-modules).

We start with a remark that is crucial when trying to find certain finitely presented direct summands in a pure-projective module.

FACT 3.1 [**21**, 34.1(3)]. *If N is a finitely generated pure submodule of a pure-projective module M, then N is a direct summand of M. (This is because, by* [**21**, 34.1(2)]*, the factor M/N is pure-projective; hence the pure-exact sequence $0 \longrightarrow N \longrightarrow M \longrightarrow M/N \longrightarrow 0$ splits.)*

Pure-projective modules share many properties with finitely presented modules, the most important model-theoretic one being the following, which was noticed by Ivo Herzog, cf. [**16**, remark before 2. 3 (a)].

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

FACT 3.2. *Every tuple in a pure-projective module has a finitely generated pp-type.*

Here is another similarity between pure-projective and finitely presented modules (cf. Fact 2.1).

LEMMA 3.3. *Let M be a pure-projective and N be an arbitrary module. Suppose that $\overline{m} \in M$ and $\overline{n} \in N$ are such that $pp_M(\overline{m}) \subseteq pp_N(\overline{n})$.*

Then there is a pointed morphism $f : (M, \overline{m}) \longrightarrow (N, \overline{n})$. Consequently, $(M, \overline{m})$ is a free realization of any formula that generates $pp_M(\overline{m})$.

Proof. Being pure-projective, M is a direct summand of a direct sum of finitely presented modules M_i, $i \in I$. Decompose $\overline{m}$ as $(\overline{m}_1, \ldots, \overline{m}_k)$ in $\bigoplus_{i\in I} M_i$. If φ_i generates $pp_{M_i}(\overline{m}_i)$, then $\varphi = \varphi_1 + \ldots + \varphi_k$ generates $p = pp_M(\overline{m})$; hence, by hypothesis, $\varphi \in pp_N(\overline{n})$. Therefore we can write $\overline{n} = \overline{n}_1 + \ldots + \overline{n}_k$ so that $N \models \varphi_i(\overline{n}_i)$ for each i.

Since each M_i is finitely presented, there is a pointed morphism $f_i : (M_i, \overline{m}_i) \longrightarrow (N, \overline{n}_i)$. Then, for $f = f_1 \oplus \ldots \oplus f_k : \bigoplus_{i\in I} M_i \longrightarrow N$, we have $f(\overline{m}) = \overline{n}$. Now restrict f to M. □

By a variant of Kaplansky's theorem (see [**5**, Theorem 2.47]), every pure-projective module is a direct sum of countably generated (and automatically pure-projective) modules. This enables us to make extensive use of the following key fact, a partial converse to Fact 3.2 (which is implicit also in [**17**, Theorem 2.2]).

FACT 3.4 [**16**, Corollary 2.9, Lemma 3.9]. *A countably generated module is pure-projective if and only if all tuples have a finitely generated pp-type.*

We can improve on this by keeping track of the pp-formulae involved. Besides, it suffices to consider pp-types of generators.

PROPOSITION 3.5. *Let $m_1, m_2, \ldots$ be a countable sequence of generators for a module M such that $\varphi_i(x_1, \ldots, x_i)$ generates $p_i = pp_M(m_1, \ldots, m_i)$. Then M is a direct summand of $N = \bigoplus_{i\geqslant 1} N_i$, where N_i is a finitely presented free realization of $\varphi_i(0, \ldots, 0, x)$. In particular, M is pure-projective.*

Proof. Let (N_i, n_i') be a finitely presented free realization of $\varphi_i(0, \ldots, 0, x)$, and set $M_i = \bigoplus_{k=1}^{i} N_k$. By induction on i, we will construct elements $n_i \in M_i \subseteq N$ and an ascending chain of morphisms $f_i : M_i \to M$ such that $M_i \models \varphi_i(n_1, \ldots, n_i)$ and $f_i(n_j) = m_j$ for every $j \leqslant i$. Then the map $f = \bigcup_{i\geqslant 1} f_i$ is an epimorphism $N \longrightarrow M$.

Further, the choice of φ_i yields the type inclusion

$$pp_M(m_1, \ldots, m_i) \subseteq pp_{M_i}(n_1, \ldots, n_i) = pp_N(n_1, \ldots, n_i).$$

Therefore $g(m_i) = n_i$ defines a morphism $M \longrightarrow N$, for if $m_1 r_1 + \ldots + m_i r_i = 0$, by this inclusion, also $n_1 r_1 + \ldots + n_i r_i = 0$. Since $fg(m_i) = f(n_i) = m_i$ for every i, the map f splits, and so M is a direct summand of N, as desired.

It remains to find n_i, as described. For $i = 1$, take $n_1 = n_1'$. Since N_1 is finitely presented, there is a pointed morphism $f_1 : (N_1, n_1) \longrightarrow (M, m_1)$.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

Suppose that we have already constructed n_k and f_k for every $k \leqslant i$. Clearly $\varphi_i(x_1, \ldots, x_i) \longrightarrow \exists\, x_{i+1}\, \varphi_{i+1}(x_1, \ldots, x_i, x_{i+1})$. Since $M_i \models \varphi_i(n_1, \ldots, n_i)$, there is $n''_{i+1} \in M_i$ such that $M_i \models \varphi_{i+1}(n_1, \ldots, n_i, n''_{i+1})$. Applying, f_i we obtain $M \models \varphi_{i+1}(m_1, \ldots, m_i, f_i(n''_{i+1}))$. However, $M \models \varphi_{i+1}(m_1, \ldots, m_i, m_{i+1})$ also, and so $M \models \varphi_{i+1}(0, \ldots, 0, m_{i+1} - f_i(n''_{i+1}))$. Since (N_{i+1}, n'_{i+1}) is a free realization of $\varphi_{i+1}(0, \ldots, 0, x)$, there is a pointed morphism $f'_{i+1} : (N_{i+1}, n'_{i+1}) \longrightarrow (M, m_{i+1} - f_i(n''_{i+1}))$. Now take $f_{i+1} = f_i \oplus f'_{i+1}$ and $n_{i+1} = n''_{i+1} + n'_{i+1}$. Then

$$f_{i+1}(n_{i+1}) = f_i(n''_{i+1}) + f'_{i+1}(n'_{i+1}) = f_i(n''_{i+1}) + m_{i+1} - f_i(n''_{i+1}) = m_{i+1}\,.$$

Finally, adding $M_i \models \varphi_{i+1}(n_1, \ldots, n_i, n''_{i+1})$ and $N_{i+1} \models \varphi_{i+1}(0, \ldots, 0, n'_{i+1})$, we obtain $M_{i+1} \models \varphi_{i+1}(n_1, \ldots, n_i, n_{i+1})$. □

In the following lemma, which will be used in the next section, we address the question about when a pp-type in a direct limit is finitely generated.

LEMMA 3.6. *Let $\langle (M_i, \overline{m}_i), f_{ij} \rangle$ be a directed system of pointed modules over the directed set I, where $(M_i, \overline{m}_i)$ is a free realization of φ_i for every $i \in I$. Further, let $\langle (M, \overline{m}), f_i \rangle$ be the corresponding direct limit. Then the following are equivalent.*

(1) *$pp_M(\overline{m})$ is finitely generated.*
(2) *There exists i such that $\varphi_i \longrightarrow \varphi_j$ for every $j \geqslant i$.*
(3) *There exists i such that all φ_j with $j \geqslant i$ are equivalent to φ_i.*

Whenever these conditions hold, $pp_M(\overline{m})$ is generated by φ_i.

Proof. Applying f_{ij} to $M_i \models \varphi_i(\overline{m}_i)$, we obtain $M_j \models \varphi_i(\overline{m}_j)$; hence $\varphi_j \longrightarrow \varphi_i$ for all $j \geqslant i$. Similarly, $f_i : M_i \longrightarrow M$ yields $\varphi_i \in pp_M(\overline{m})$ for every i. Denote this latter type by p.

(1) ⇒ (3): Assume that p is generated by φ. As $\varphi_j \in p$, it follows that $\varphi \longrightarrow \varphi_j$ for every j. However, $M \models \varphi(\overline{m})$ implies that $M_i \models \varphi(\overline{m}_i)$ for sufficiently large i. Since $(M_i, \overline{m}_i)$ is a free realization of φ_i, we obtain $\varphi_i \longrightarrow \varphi$ for these i; hence also $\varphi_i \longrightarrow \varphi \longrightarrow \varphi_j \longrightarrow \varphi_i$ for every $j \geqslant i$.

(2) ⇒ (1): Suppose that i is as in (2). We claim that φ_i generates p. We know already that $\varphi_i \in p$. If now $\psi \in p$, as above it follows that $\varphi_j \longrightarrow \psi$ for sufficiently large j. Choosing $j \geqslant i$, by (2), we obtain $\varphi_i \longrightarrow \varphi_j \longrightarrow \psi$.

As (3) ⇒ (2) is trivial, this completes the proof. □

The next result, which was inspired by Dung and Facchini [**3**, Theorem 4.9] (see also [**5**, Proposition 9.30]), will be used in the main theorem on uniserial rings below. Some more explanations of motivation may be in order. Although this is not the point of view taken in the next proof, a direct limit (M, f_i) of a direct system of modules (N_i, f_{ij}), $i, j \in I$, can be regarded as an epimorphic image of the direct sum $\bigoplus_{i \in I} N_i$. Further, the corresponding epimorphism is easily seen to be pure, cf. [**21**, 33.9(2)]. Thus, if M itself is pure-projective, then this map splits, that is, M is a direct summand of $\bigoplus_{i \in I} N_i$. However, this is a mere existence statement, which does not indicate the complement, let alone the map constituting the splitting. This exactly is what the (proof of the) next theorem does, in the case when M is countably generated. (Note that it also constitutes a generalization of Eilenberg's trick.)

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

THEOREM 3.7 (telescoping map theorem). *Suppose that M is a countably generated pure-projective module. If M is the direct limit of a direct system (N_i, f_{ij}), $i, j \in I$, then $M \oplus \bigoplus_{i\in I} N_i \cong \bigoplus_{i\in I} N_i$.*

Proof. Represent the elements of M as (equivalence classes of) pairs (m, i) with $i \in I$ and $m \in N_i$, and where $(m, i) = (n, j)$ if there exists $k \geqslant i, j$ such that $f_{ik}(m) = f_{jk}(n)$.

Let $m_1, m_2, \ldots$ be a countable list of generators of M. By Fact 3.4, the pp-type of any finite tuple of elements from this list is generated by a single formula. Therefore there is a chain of indices $i_1 < i_2 < \ldots \in I$ such that $m_k = (n_k, i_k)$ and the tuple $(m_1, \ldots, m_{k-1}, m_k)$ has the same pp-type in M as does the tuple $(f_{i_1,i_k}(n_1), \ldots, f_{i_{k-1},i_k}(n_{k-1}), n_k)$ in N_{i_k}.

Denote $\{i_1, i_2, \ldots\}$ by I'. First we show that

$$M \cong \varinjlim_{i,j\in I'}(N_i, f_{ij}).$$

Since any element $m \in M$ is a linear combination of some of the (n_k, i_k), we may write m as (n, j), where j is the highest occurring index from I' (in that linear combination) and n is the corresponding linear combination of the $f_{i_k,j}(n_k)$. Hence the natural map

$$\varinjlim_{i,j\in I'}(N_i, f_{ij}) \longrightarrow \varinjlim_{i,j\in I}(N_i, f_{ij})$$

(induced by the inclusion $I' \subseteq I$) is onto.

To prove that this map is also mono, let $m = (n, j)$ go to zero, that is, $m = 0$ in M. Notice that the aforementioned equality of pp-types is preserved by passing to the corresponding linear combinations; hence $pp_M(m) = pp_{N_j}(n)$, and so $n = 0$ as well. However, $m = (n, j) = 0$ also in $\varinjlim_{i,j\in I'}(N_i, f_{ij})$.

Thus we may assume that

$$M = \varinjlim_{i,j,\in I'}(N_i, f_{ij}),$$

and if we prove that $M \oplus \bigoplus_{i\in I'} N_i \cong \bigoplus_{i\in I'} N_i$, then adding $\bigoplus_{i\in I\setminus I'} N_i$ to both sides finishes off the proof. We may therefore even assume that $I = \omega = \{1, 2, \ldots\}$, that $m_i = (n_i, i)$ with $n_i \in N_i$, and that $pp_M(m_1, \ldots, m_{k-1}, m_k) = pp_{N_k}(f_{1k}(n_1), \ldots, f_{k-1,k}(n_{k-1}), n_k)$. Then every element $m = m_1r_1 + m_2r_2 + \ldots + m_kr_k$ in M has a representation (n, k) with $n = f_{1,k}(n_1)r_1 + f_{2,k}(n_2)r_2 + \ldots + n_kr_k$.

Since M is pure-projective, by Lemma 3.3, there exists a morphism $g_k : M \longrightarrow N_k$ such that $g_k(m_i) = f_{ik}(n_i)$ for every $i \leqslant k$. It is easily checked that therefore $g_k(m) = n$ when $m = (n, k)$ is a representation, as in the previous paragraph.

Let $N = \bigoplus_{i\in I} N_i$, and define $g : M \longrightarrow N$ by sending $m \in M$ to

$$(g_1(m), g_2(m) - f_{12}g_1(m), \ldots, g_k(m) - f_{k-1,k}g_{k-1}(m), \ldots).$$

Since $g_{j+1}(m_i) = f_{j,j+1}g_j(m_i)$ for every $j \geqslant i$, this map is well defined.

We claim that $\ker(g) = 0$ (and so $\mathrm{im}(g) \cong M$). Indeed, let $m \in M$ be in the kernel and represent it by $m = (n, k)$ as above so that $g_k(m) = n$. From $g(m) = 0$, we successively obtain $0 = g_1(m) = \ldots = g_k(m) = n$; hence $m = (0, k)$ is zero in M.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

Now define $h : N \longrightarrow M$ by $h(l_1, \ldots, l_k, 0, \ldots) = (l_1, 1) + \ldots + (l_k, k)$. Then

$$\begin{aligned} hg(m_i) &= h(g_1(m_i), g_2(m_i) - f_{12}g_1(m_i), \ldots, g_i(m_i) - f_{i-1,i}g_{i-1}(m_i), 0, \ldots) \\ &= (g_1(m_i), 1) + (g_2(m_i), 2) - (f_{12}g_1(m_i), 2) + \ldots + (g_i(m_i), i) \\ &\quad - (f_{i-1,i}g_{i-1}(m_i), i) \\ &= (g_i(m_i), i) = (n_i, i) = m_i. \end{aligned}$$

Thus g splits, that is, $N = \mathrm{im}(g) \oplus \ker(h)$. As $\mathrm{im}(g) \cong M$, it remains to prove that $\ker(h) \cong N$. To this end, let $s : N \longrightarrow N$ be the map that sends $n = (l_1, l_2, \ldots, l_k, 0, \ldots)$ to $(l_1, l_2 - f_{12}(l_1), \ldots, l_k - f_{k-1,k}(l_{k-1}), -f_{k,k+1}(l_k), 0, \ldots)$.

We claim that $\ker(s) = 0$; indeed, if $s(n) = 0$, we successively obtain $0 = l_1 = \ldots = l_k$, whence $n = 0$. Hence $\mathrm{im}(s) \cong N$, and so it suffices to prove that $\mathrm{im}(s) = \ker(h)$.

For $\mathrm{im}(s) \subseteq \ker(h)$, consider $n = (l_1, l_2, \ldots, l_k, 0, \ldots) \in N$. Then $hs(n) = (l_1, 1) + (l_2, 2) - (f_{12}(l_1), 2) + \ldots + (l_k, k) - (f_{k-1,k}(l_{k-1}), k) - (f_{k,k+1}(l_k), k + 1) = 0$, since $(l_i, i) = (f_{i,i+1}(l_i), i + 1)$ for all i.

We are left with $\ker(h) \subseteq \mathrm{im}(s)$. Let n be as before and $h(n) = 0$, that is, $(l_1, 1) + \ldots + (l_k, k) = 0$. By the definition of direct limit, there is $j \geqslant k$ such that $f_{1j}(l_1) + \ldots + f_{kj}(l_k) = 0$. We may take j to be strictly bigger than k.

Consider the element $n' \in N$ whose ith component n'_i is defined as follows. If $i \leqslant k$, set $n'_i = f_{1i}(l_1) + \ldots + f_{i-1,i}(l_{i-1}) + l_i$. If $k < i \leqslant j$, set $n'_i = f_{k,i}(n'_k)$. For $i > j$, set $n'_i = 0$. Then easy calculations show that $n'_{i+1} - f_{i,i+1}(n'_i) = l_{i+1}$ for all $i < k$ and $n'_{i+1} - f_{i,i+1}(n'_i) = 0$ for $k \leqslant i < j$, while $n'_{j+1} - f_{j,j+1}(n'_j) = -f_{j,j+1}(n'_j) = 0$, because $n'_j = f_{1j}(l_1) + \ldots + f_{kj}(l_k) = 0$ by the choice of j.

Consequently, $s(n') = (l_1, l_2, \ldots, l_k, 0, \ldots) = n$, as desired, which completes the proof. □

In the proof of Theorem 5.8 we need one more technical fact about pure-projectives. For this, recall that a module is called *Bezout* if each of its finitely generated submodules is cyclic. A module M is said to be *endo-Bezout* if ${}_SM$ is Bezout, where $S = \mathrm{End}(M)$.

LEMMA 3.8. *Given a pure-projective module M, the following are equivalent.*

(1) *M is endo-Bezout.*

(2) *For any one-place pp-formulae φ and ψ, if the pp-type of some element of M is generated by φ and the pp-type of some element of M is generated by ψ, then there is an element of M whose pp-type is generated by $\varphi + \psi$.*

Proof. Let m and n be arbitrary elements of M such that $pp_M(m)$ is generated by φ and $pp_M(n)$ is generated by ψ.

(1) $\Rightarrow$ (2): Condition (1) yields an element $k \in M$ such that $Sm + Sn = Sk$. We claim that $r = pp_M(k)$ is generated by $\varphi + \psi$.

Write $k = fm + gn$, where f and g are in S. Then $M \models \varphi(m)$ implies that $M \models \varphi(fm)$, and $M \models \psi(n)$ implies that $M \models \psi(gn)$. Then $M \models (\varphi + \psi)(k)$, that is, $\varphi + \psi \in r$.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

It remains to prove that $\varphi + \psi \longrightarrow \xi$ for every $\xi \in r$. Indeed, from $m \in Sk$ and $M \models \xi(k)$, we obtain $M \models \xi(m)$; hence $\varphi \longrightarrow \xi$, as $pp_M(m)$ is generated by φ. Similarly, $\psi \longrightarrow \xi$, and so $\varphi + \psi \longrightarrow \xi$.

(2) $\Rightarrow$ (1): Condition (2) yields an element $k \in M$ whose pp-type is generated by $\varphi + \psi$. As $\varphi \longrightarrow \varphi + \psi$, Fact 2.1 implies that $m \in Sk$. Similarly, $n \in Sk$, and hence $Sm + Sn \subseteq Sk$.

If, conversely, $l \in Sk$, then we have $M \models (\varphi + \psi)(l)$. Write $l = m' + n'$ accordingly so that $M \models \varphi(m') \wedge \psi(n')$. As above, there are $f, g \in S$ such that $f(m) = m'$ and $g(n) = n'$. Then $l = m' + n' \in Sm + Sn$. □

4. *A result of Bass and Björk revisited*

As a first application of our techniques, using an idea of Sakhayev [**18**], we give a new proof of a result of Bass and Björk, Theorem 4.2, a proof that does not use Björk's theorem. We start with a special case of Prest's lemma.

REMARK 4.1. Let φ be the pp-formula $A \,|\, \overline{x}$ and ψ be the pp-formula $B \,|\, \overline{x}$. Then $\varphi \longrightarrow \psi$ if and only if there is C such that $A = CB$.

Proof. If $A = CB$, then the implication that $\varphi \longrightarrow \psi$ is obvious. For the converse, assume that $\varphi \longrightarrow \psi$, and let M be a free module generated by a tuple $\overline{m}$ of the same length as $\overline{x}$. Since $\overline{m}A$ trivially satisfies φ, it also satisfies ψ, that is, $M \models B \,|\, \overline{m}A$. Therefore there is some matrix C such that $(\overline{m}C)B = \overline{m}A$; hence $A = CB$, as desired. □

We emphasize the fact that the following new proof makes no reference to Björk's theorem.

THEOREM 4.2 (Bass–Björk result). *If every flat right R-module is projective, then every left R-module has the DCC on finitely generated submodules.*

Proof. Let $M_1 \supseteq M_2 \supseteq \ldots$ be a chain of left R-modules with each M_i generated by some $\overline{m}_i$. Then $\overline{m}_{i+1} = B_i \overline{m}_i$ for appropriate matrices B_i.

Correspondingly, let N be the right module with generators $\overline{n}_1, \overline{n}_2, \ldots$ and relations $\overline{n}_i = \overline{n}_{i+1} B_i$, $i = 1, 2, \ldots$ (see Figure 2).




FIGURE 2.

Consider the free module N_i generated by a tuple $\overline{x}_i$ (matching $\overline{n}_i$) and the morphism $f_{i,i+1} : N_i \longrightarrow N_{i+1}$ given by $\overline{x}_i \longmapsto \overline{x}_{i+1} B_i$. It is easy to verify that N, together with the maps $f_i : N_i \longrightarrow N$ given by $f_i(\overline{x}_i) = \overline{n}_i$, is the direct limit of the thus given directed system.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

By Fact 2.4, the pp-type of $f_{ij}(\overline{x}_i)$ in N_j is generated by the formula $B_{j-1} \cdot \ldots \cdot B_i \,|\, \overline{x}$. In particular, the pp-type of $f_{1j}(\overline{x}_1)$ in N_j is generated by the formula $B_{j-1} \cdot \ldots \cdot B_1 \,|\, \overline{x}$, which we denote by φ_j. It is easy to see that, in fact, $(N_j, f_{1j}(\overline{x}_1))$ is a free realization of φ_j.

Being a direct limit of free modules, N is flat, and hence, by hypothesis, projective. In particular, N is pure-projective, and so, by Fact 3.2, $pp_N(\overline{n}_1)$ is finitely generated. Therefore, Lemma 3.6 yields i such that $\varphi_i \longrightarrow \varphi_j$ for every $j \geqslant i$, that is, $B_{i-1} \cdot \ldots \cdot B_1 \,|\, \overline{x} \longrightarrow B_{j-1} \cdot \ldots \cdot B_1 \,|\, \overline{x}$. By Remark 4.1, there is a matrix C_{j-1} such that $C_{j-1}B_{j-1} \cdot \ldots \cdot B_1 = B_{i-1} \cdot \ldots \cdot B_1$. However, $\overline{m}_i = B_{i-1} \cdot \ldots \cdot B_1 \overline{m}_1 = C_{j-1}B_{j-1} \cdot \ldots \cdot B_1 \overline{m}_1 = C_{j-1}\overline{m}_j$ for every $j \geqslant i$. Consequently, the chain $M_1 \supseteq M_2 \supseteq \ldots$ stabilizes after i steps. □

Notice that, applying Theorem 3.7 to the directed system leading to N above, we obtain $N \oplus R^{(\omega)} \cong R^{(\omega)}$, a fact that is known as Eilenberg's trick for *any* (countably generated) projective module.

5. *Uniserial rings*

The main objective of this section is to study the category $\mathrm{Add}(M)$ for a cyclically presented module $M = R/rR$ over a uniserial ring R. (Recall that a module M is *uniserial* if the lattice of submodules of M is a chain, while a ring R is *uniserial* if R_R and ${}_RR$ are uniserial modules.) We will determine when exactly this category is *trivial* in the sense that each module in $\mathrm{Add}(M)$ is a direct sum of copies of M. (Recall that the objects of $\mathrm{Add}(M)$ are the direct summands of all the direct sums $M^{(I)}$ of copies of M, so all of them are pure-projective.) In other words, we will answer our principal question stated in the introduction for this very special kind of pure-projective. For the remainder of this section, we will work in this setting of $\mathrm{Add}(M)$, where M is a fixed cyclically presented module R/rR over a uniserial ring R, where without loss, $r \in \mathrm{Jac}(R)$. This is because every uniserial ring is local; hence, if r is not in the Jacobson radical, then r is a unit and so $R/rR = 0$. Note that (even for an arbitrary finitely generated right module M) the study of $\mathrm{Add}(M)$ is equivalent to the study of projective right modules over $S = \mathrm{End}(M)$, since there exists an equivalence between $\mathrm{Add}(M)$ and the category of projective right S-modules (see [**1**, Lemma 29.4] or [**5**, Theorem 4.7]).

Any cyclically presented R-module is uniserial and hence indecomposable. The next fundamental result states that these are the building blocks for the finitely presented modules.

Fact 5.1 (Drozd and Warfield, see [**12**, Theorem 2.3]). *Every finitely presented module over a uniserial ring can be decomposed into a finite direct sum of uniserial cyclically presented modules. (This also holds for serial rings, that is, rings R such that both ${}_RR$ and R_R are direct sums of uniserial modules.)*

It was noticed by the first author ([**12**, Proposition 2.20], see also [**5**, Theorem 9.19]) that such a decomposition is unique.

The following result sheds some light on the structure of the category $\mathrm{Add}(M)$.

Fact 5.2 [**3**, Corollary 2.8]. *If S is the endomorphism ring of a uniserial module M, then every finitely generated projective right S-module is free. Consequently,*

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

every finitely generated module in $\mathrm{Add}(M)$ *is isomorphic to a direct sum of finitely many copies of* M.

As noted in [**5**, Proposition 3.12 and remarks thereafter], homing into the ring constitutes a duality between the finitely generated projective left modules and the finitely generated projective right modules (even for an arbitrary ring), and therefore all the left ones are free if and only if all the right ones are also. Thus, for S, the endomorphism ring of a uniserial module, both the left *and* the right finitely generated projectives are free. Note that such rings are even *projective free* in the terminology of Cohn [**2**, §0.2] (another left–right symmetric notion, see [**2**, remark after Proposition 0.2.7]), which means that finitely generated projectives are free *of unique rank*. Indeed, being semilocal, S has finite dual Goldie dimension. Therefore the rank of every finitely generated free right module is uniquely determined.

It is known that there are uniserial domains over which $\mathrm{Add}(M)$ can have a nontrivial decomposition theory, that is, there exists a projective right S-module that is not free (see [**13**, Corollary 8.2]). We will give a precise criterion for this to happen.

First we establish some important properties of cyclically presented modules over uniserial rings.

LEMMA 5.3. *Every cyclically presented module over a uniserial ring is endo-Bezout.*

Proof. According to [**12**, Corollary 11.15], every finitely presented module over a uniserial ring is endo-distributive (that is, distributive as a module over its own endomorphism ring). Since M is uniserial, by [**5**, Theorem 9.1], $S = \mathrm{End}(M)$ has at most two maximal right ideals, and $S/\mathrm{Jac}(S)$ is a direct sum of at most two skew fields. Now, by [**19**, 3.33], distributivity is equivalent to the Bezout property over any ring which is abelian (von Neumann) regular after factorization by the Jacobson radical. □

Note that every pure-projective module over a uniserial ring is endo-distributive, but it is not known if it has to be endo-Bezout.

The previous lemma allows us to describe the generation of pp-types in (pure-projective) modules in $\mathrm{Add}(M)$.

LEMMA 5.4. *Let* R *be a uniserial ring and let* $M = R/rR$, *where* $0 \neq r \in \mathrm{Jac}(R)$.

The pp-types of non-zero elements in any $N \in \mathrm{Add}(M)$ *are generated by formulae of the form* $a \mid x \wedge xb = 0$, *where* $0 \neq a \in R$ *and* $0 \neq b \in \mathrm{Jac}(R)$ *such that* $r = ab$.

Further, for non-zero elements $m \in M$, *the ring element* a *can be chosen in the* rR*-coset* m *itself.*

Proof. We first prove the assertion for M. Therefore, let $m = a + rR$ be a non-zero element of M. As $(R/rR, 1 + rR)$ is a free realization of $xr = 0$, Fact 2.4 tells us that (M, m) is a free realization of the formula $\exists y\ (ya = x \wedge yr = 0)$. As R is uniserial, either $a \in rR$, which contradicts $m \neq 0$, or else $r = ab$ for some $b \in R$. Then, as is easily seen, b must be in $\mathrm{Jac}(R)$ and $\exists y\ (ya = x \wedge yr = 0)$ is equivalent to $a \mid x \wedge xb = 0$.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

In order to prove this for any $N \in \mathrm{Add}(M)$, let $0 \neq n \in N$. Then N is pure-projective, and $pp_N(n)$ is generated by a formula $\varphi_1 + \ldots + \varphi_k$, where φ_i generates the pp-type of some $m_i \in M$. As M is endo-Bezout, Lemma 3.8 yields some element $0 \neq m \in M$ with $pp_N(n) = pp_M(m)$, and the assertion follows from the first part of the proof. □

Next we cite two more technical facts, where, for $r, s \in R$, we write $r \leqslant s$ if $RrR \supseteq RsR$, and $r \sim s$ if $RrR = RsR$.

FACT 5.5 [**14**, Corollary 3.2]. *Let R be a uniserial ring and let $0 \neq a, b, c \in \mathrm{Jac}(R)$. Then the following are equivalent.*

(1) $a \mid x \wedge xb = 0 \longrightarrow ca \mid x$.
(2) $a \sim cab$.

Notice that $a \sim cab$ is equivalent to $a \in RcabR$, where $a \leqslant cab$ always holds.

FACT 5.6 [**12**, Proposition 2.21]. *Let r, s be elements of a uniserial ring R. Then the following are equivalent.*

(1) $R/rR \cong R/sR$.
(2) $r \sim s$ (*that is*, $RrR = RsR$).
(3) $r = usv$ *for some units* $u, v \in R$.

The following construction is used in the theorem below and may be of interest in its own right.

LEMMA 5.7. *Suppose that $r_1, r_2, \ldots \in \mathrm{Jac}(R)$ are such that $r_i \cdot \ldots \cdot r_2 \neq 0$ for every $i > 1$.*

Then the module N with generators $n_1, n_2, \ldots$ and relations $n_1 r_1 = 0$ and $n_{i+1} r_{i+1} = n_i$ for all $i \geqslant 1$ is not finitely generated.

Proof. Depict N as shown in Figure 3.




FIGURE 3.

If N were finitely generated, it would be generated by a single n_k. We show that this is impossible. In fact, we will reach a contradiction from the assumption that $n_{k+1} = n_k s$ for some $s \in R$.

Indeed, as $n_k = n_{k+1} r_{k+1}$, this assumption yields $n_{k+1}(1 - r_{k+1}s) = 0$; hence $n_{k+1} = 0$, for, by hypothesis, $1 - r_{k+1}s$ is a unit. Then $n_{k+1} = n_k = \ldots = n_1 = 0$. By the definition of N, n_1 must be a linear combination of the 'relations', whence

$$n_1 = n_1 r_1 t_1 + (n_2 r_2 - n_1) t_2 + \ldots + (n_m r_m - n_{m-1}) t_m$$

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

for some ring elements t_i. Comparing coefficients, we obtain

$$1 = r_1t_1 - t_2, \quad 0 = r_2t_2 - t_3, \quad \ldots, \quad 0 = r_{m-1}t_{m-1} - t_m, \quad 0 = r_mt_m.$$

These imply that

$$0 = r_mt_m = r_mr_{m-1}t_{m-1} = \ldots = r_m \cdot \ldots \cdot r_2t_2 = r_m \cdot \ldots \cdot r_2(r_1t_1 - 1).$$

The right-hand factor is a unit again, so $r_m \cdot \ldots \cdot r_2$ must be 0, contradicting the hypothesis. □

We are ready now to give the promised criterion.

THEOREM 5.8. *Let R be a uniserial ring, and let $0 \neq r \in \mathrm{Jac}(R)$, $M = R/rR$ and $S = \mathrm{End}(M)$. Then the following are equivalent.*

(1) $\mathrm{Add}(M)$ *is trivial, that is, every module in* $\mathrm{Add}(M)$ *is isomorphic to some* $M^{(I)}$.

(2) *Every projective right S-module is free.*

(3) *If $r \sim a \sim b$ in R, then $r \neq ab$.*

(4) *Every non-zero element of any module $N \in \mathrm{Add}(M)$ is contained in a direct summand of N isomorphic to M.*

(5) *Every module in* $\mathrm{Add}(M)$ *contains a direct summand that is isomorphic to M.*

(6) *Every module in* $\mathrm{Add}(M)$ *contains a non-zero finitely generated direct summand.*

(7) *Every module in* $\mathrm{Add}(M)$ *is a direct sum of finitely generated modules.*

Proof. The equivalence of (1) and (2) follows from the aforementioned equivalence of categories. The equivalence of (5) and (6) as well as that of (1) and (7) follows from Fact 5.2. The implication that (5) ⇒ (1) is [**4**, Corollary 2.10]. Since (4) ⇒ (5) is trivial, we are left with (1) ⇒ (3) and (3) ⇒ (4).

(1) ⇒ (3): Assume, on the contrary, that there are $a, b \in R$ such that $ab = r$ and $a, b \sim r$, In particular, $0 \neq a, b \in \mathrm{Jac}(R)$. We will produce a module N in $\mathrm{Add}(M)$ which is not of the form $M^{(I)}$.

To this end, we choose as follows non-zero elements $r_1, r_2, \ldots$ in $\mathrm{Jac}(R)$ such that $r_i \sim r$ and $r_i \cdot \ldots \cdot r_1 \sim r$ for every i.

Since $a \sim b \sim ab = r$, we may start with $r_1 = b$ and $r_2 = a$. Now suppose that $r_1, \ldots, r_i$ have already been chosen so that $r_i \cdot \ldots \cdot r_1 \sim r$. Then $r_i \cdot \ldots \cdot r_1 \sim b$. By Fact 5.6, there are units $u, v \in R$ such that $r_i \cdot \ldots \cdot r_1 = ubv$. Then $u^{-1}r_i \cdot \ldots \cdot r_1 = bv$; hence $au^{-1}r_i \cdot \ldots \cdot r_1 = abv = rv \sim r$, whence we may take $r_{i+1} = au^{-1}$.

We claim that the module N with generators $n_1, n_2, \ldots$ and relations $n_1r_1 = 0$ and $n_{i+1}r_{i+1} = n_i$ for $i \geqslant 1$ is a uniserial pure-projective that is (countably but) not finitely generated.

As cyclic modules over uniserial rings are uniserial and N is the union of a chain of cyclic modules, we see that N is uniserial. Further, Lemma 5.7 states that N is not finitely generated. However, it is countably generated, and so to show that it is pure-projective, by Proposition 3.5, it suffices to check that every $p_i = pp_N(n_i)$ is finitely generated, which we do next.

Fix i, and consider, for all $j > i$, the submodule $N_j = n_jR$ of N. As noted before Fact 5.5, $pp_{N_j}(n_i)$ is generated by the formula $r_j \cdot \ldots \cdot r_{i+1} \,|\, x \wedge xr_i \cdot \ldots \cdot r_1 = 0$, which we denote by φ_j. Given $k > j > i$, we clearly have $\varphi_k \longrightarrow \varphi_j$. We claim

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

that these formulae are in fact equivalent, for which we just have to show that $\varphi_j \longrightarrow r_k \cdot \ldots \cdot r_{i+1} \,|\, x$. However, this latter follows from Fact 5.5 with $c = r_k \cdot \ldots \cdot r_{j+1}$, $a = r_j \cdot \ldots \cdot r_{i+1}$, and $b = r_i \cdot \ldots \cdot r_1$, for $r \sim r_j \leqslant r_j \cdot \ldots \cdot r_{i+1} = a \leqslant cab = r_k \cdot \ldots \cdot r_1 \sim r$.

Thus the formulae φ_j are equivalent for all $j > i$. Therefore, the types $pp_{N_j}(n_i)$ are the same for all $j > i$ as well. Hence they are the same as $pp_N(n_i)$, which means that the latter is finitely generated, as desired.

Now that we have proved the claim about N, we may apply to it the (telescoping) Theorem 3.7 to obtain the decomposition

$$N \oplus \bigoplus_i R/(r_i \cdot \ldots \cdot r_1 R) \cong \bigoplus_i R/(r_i \cdot \ldots \cdot r_1 R).$$

However, $R/(r_i \cdot \ldots \cdot r_1 R) \cong R/rR$ by Fact 5.6. Consequently, $N \oplus M^{(\omega)} \cong M^{(\omega)}$, whence $N \in \mathrm{Add}(M)$.

Finally, if N were of the form $M^{(I)}$, then being uniserial, it would have to be isomorphic to M, which is impossible, as the latter is finitely generated and N is not.

(3) $\Rightarrow$ (4): Suppose that N is in $\mathrm{Add}(M)$ and $0 \neq n \in N$. We must prove that n is contained in a direct summand isomorphic to M. By Lemma 5.4, $pp_N(n)$ is generated by a formula of the form $a \,|\, x \wedge xb = 0$, where $ab = r$, $a \neq 0$, and $0 \neq b \in \mathrm{Jac}(R)$.

If a is invertible, then $pp_N(n)$ is obviously generated by the quantifier-free formula $xa^{-1}r = 0$, and hence nR is a (finitely generated) pure submodule of the pure-projective module N, and consequently, by Fact 3.1, a direct summand. It remains for us to notice that $nR \cong R/a^{-1}rR \cong R/rR \cong M$ to see that in this case we are done.

Therefore assume that a is not invertible. Then $0 \neq a, b \in \mathrm{Jac}(R)$, which will allow us to apply Fact 5.5 as follows.

Choose $n_1 \in N$ such that $n_1 a = n$. The type $pp_N(n_1)$ is also generated by a formula $a_1 \,|\, x \wedge xb_1 = 0$ for some $0 \neq a_1 \in R$ and $0 \neq b_1 \in \mathrm{Jac}(R)$ such that $a_1 b_1 = r$.

If a_1 is invertible, then $n_1 R$ is a direct summand isomorphic to M as before, and we are done again. If not, $a_1 \in \mathrm{Jac}(R)$ as well.

Since $pp_N(n)$ is generated by $a \,|\, x \wedge xb = 0$, it follows that $a \,|\, x \wedge xb = 0 \longrightarrow a_1 a \,|\, x$. Hence from Fact 5.5, we obtain $a \sim a_1 ab = a_1 r$. Then $r \leqslant a_1 r \sim a \leqslant ab = r$; hence $a_1 r \sim r$. By Fact 5.6, this yields units u and v such that $r = u a_1 r v$. As, clearly, $rv \sim r$, (3) therefore implies that $ua_1 \not\sim r$; hence $a_1 \not\sim r$.

Next choose $n_2 \in N$ such that $n_2 a_1 = n_1$ and a formula $a_2 \,|\, x \wedge xb_2 = 0$ generating its pp-type in N, where $a_2 b_2 = r$. We get the diagram shown in Figure 4.




FIGURE 4.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

Again, if a_2 is invertible, we finish as before. We conclude by showing that there is no other possibility.

Indeed, if a_2 were not invertible, it would be in $\mathrm{Jac}(R)$. Since $pp_N(n_1)$ is generated by $a_1 \mid x \wedge xb_1 = 0$, we would therefore obtain, as above, $a_1 \mid x \wedge xb_1 = 0 \longrightarrow a_2a_1 \mid x$; hence $a_1 \sim a_2a_1b_1 = a_2r$, and thus $r \leqslant a_2r \sim a_1 \leqslant a_1b_1 = r$. However this would imply that $a_1 \sim r$, contradicting the above. □

For the convenience of the reader, we conclude with a direct proof of (4) ⇒ (1), because it is based on an argument that is very useful in applications of Kaplansky's theorem. First of all, by the latter, we may assume that the module $N \in \mathrm{Add}(M)$ in question is countably generated, say by $n_1, n_2, \ldots$. Now choose a direct summand $M_1 \cong M$ of N containing n_1, and write $n_2 = m_{21} + n_2' \in N = M_1 \oplus N_1$ accordingly. Then choose a direct summand $M_2 \cong M$ of N_1 containing n_2', and write $n_3 = m_{31} + m_{32} + n_3' \in M_1 \oplus M_2 \oplus N_2$. Continuing like this, we obtain direct summands $M_i \cong M$ and a decomposition $N = M_1 \oplus \ldots \oplus M_k \oplus N_k$ such that $n_k \in M_1 \oplus \ldots \oplus M_k$. Then the submodule $\bigoplus_{i=1}^{\infty} M_i$ of N contains all generators n_i, and is therefore all of N, as desired.

6. *Noetherian rings*

Here we answer our main question in the affirmative for hereditary noetherian rings by showing that every pure-projective module over such rings is a direct sum of finitely generated modules.

The main idea is to use the torsion theory generated by singular modules. Recall that a ring R is *right Goldie* if R has finite right uniform dimension and the ascending chain condition (ACC) on right annihilators.

Let R be a semiprime right Goldie ring. Then, by Goldie's theorem, the set of regular elements of R is a right Ore set, and R has a (right) classical quotient ring $Q = Q(R)$. Further, Q is a semisimple artinian ring and, as a left R-module, it is flat (being a union of left modules $Ra^{-1} \cong {}_RR$, where a is a regular element of R).

We say that an element of a module is *torsion* if it is annihilated by a regular ring element. A module is said to be *torsion* if every element of it is. By [**10**, Proposition 2.3.5], an R-module M is torsion if and only if it is *singular*, that is, if and only if the annihilator of any $m \in M$ is an essential right ideal of R. The torsion elements of M form the *torsion submodule* $T(M)$ of M. Note that $P = M/T(M)$ is a *torsion-free* module, that is, $T(P) = 0$.

The first goal is to prove that, in pure-projective modules, the torsion part splits off. This is true even over semihereditary semiprime Goldie rings, as we show in the following two lemmas.

LEMMA 6.1. *Let M be a finitely generated module over a semihereditary semiprime Goldie ring. Then $M = T(M) \oplus P$, where P is projective.*

Proof. This can be proved as [**10**, Lemma 5.7.4]. It is worthwhile, however, to emphasize where the left and the right Goldie conditions come in now, which is why we reproduce the proof.

Since $P = M/T(M)$ is finitely generated torsion-free, and ${}_RQ$ is flat by the right Goldie condition, tensoring P with Q embeds P (as a right R-module) in Q^n. As R is also left Goldie, any $q \in Q$ can be written as $c^{-1}d$, where $c \in R$ is regular.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

Multiplying by the common left denominator on the left, we obtain an embedding of P in R^n.

Since R is (right) semihereditary, P is projective; hence the short exact sequence $0 \longrightarrow T(M) \longrightarrow M \longrightarrow P \longrightarrow 0$ splits. □

Note that semihereditarity is symmetric for rings of finite uniform dimension [**9**, Theorem 7.64].

REMARK 6.2. Every torsion-free module N over a semihereditary semiprime Goldie ring is flat. (Indeed, in the case when N is finitely generated, it is even projective by the previous proof; for the general case, write N as a direct limit of its finitely generated (projective) submodules.)

LEMMA 6.3. *Let M be a pure-projective module over a semihereditary semiprime Goldie ring. Then $M = T(M) \oplus P$, where P is projective.*

Proof. As M is pure-projective, M is a direct summand of a direct sum $N = \bigoplus_{i\in I} M_i$ of finitely presented R-modules M_i. Let $\pi : N \longrightarrow M$ be the map that splits the inclusion $M \subseteq N$.

By the previous lemma, $M_i = T(M_i) \oplus P_i$, where P_i is projective. Then $T(N) = \bigoplus_{i\in I} T(N_i)$ and hence $N = T(N) \oplus P$, where $P = \bigoplus_{i\in I} P_i$ is projective.

Clearly $T(M) \subseteq T(N)$ and $\pi(T(N)) \subseteq T(M)$; hence π also splits the inclusion $T(M) \subseteq T(N)$. Since $T(N)$ is a direct summand of N, so is $T(M)$, whence the latter is a direct summand also of M.

Thus $M = T(M) \oplus P$, where $P = M/T(M)$ is a torsion-free module. By the above remark, P is flat. However, every flat pure-projective module is projective. □

In order to decompose a pure-projective module into a direct sum of finitely generated submodules, we need to put some additional restrictions on the ring.

THEOREM 6.4. *Let M be a pure-projective module over a hereditary noetherian prime ring R. Then M is a direct sum of finitely generated modules.*

Proof. By Lemma 6.3, $M = T(M) \oplus P$, where $T(M)$ is torsion, and P is projective. Since R is right hereditary, by a result of Albrecht, P is a direct sum of finitely generated right ideals of R (see [**2**, comments on Theorem 0.2.9]).

As above, $T(M)$ is a direct summand of a direct sum of finitely generated torsion modules. However, by [**10**, Lemma 5.7.4], every finitely generated torsion module over a hereditary noetherian prime ring has finite length. Thus we may invoke the Krull–Remak–Schmidt–Azumaya theorem in the form of [**1**, Theorem 26.5] applied to the case where M is countably generated (which suffices, as pointed out before Fact 3.4). □

COROLLARY 6.5. *Every pure-projective module over a hereditary noetherian ring is a direct sum of finitely generated modules.*

Proof. By [**10**, Theorem 5.4.6], any hereditary noetherian ring is a direct sum of hereditary noetherian prime rings and an artinian (hereditary) ring. Thus the proof splits into two respective parts, the former being dealt with in the theorem.

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

In the latter, artinian, case, finitely generated modules have finite length, so the Krull–Remak–Schmidt–Azumaya argument works again. □

It is not clear to us to what other noetherian rings this result might be extended. The following example shows at least a limitation for our proof to work.

EXAMPLE 6.6. Let R be a local commutative noetherian domain which is not uniserial (the power series ring $F[[X,Y]]$ over a field F will do). Consider a non-unit $0 \neq c \in R$ and $a, b \in R$ such that the right ideals aR and bR are incomparable. Let M be the finitely presented module $\langle x, y \mid xac + ybc = 0 \rangle$. It is not hard to check that M is indecomposable (see [**6**, proof of Theorem 20.42]).

Since $(xa + yb)c = 0$ and $xa + yb \neq 0$ in M, we conclude that $T(M) \neq 0$. Indeed, $xa + yb = 0$ in M means that $xa + yb = (xac + ybc)d$ in the free module generated by x and y. Equating coefficients at x, we obtain $a = acd$; hence $cd = 1$, a contradiction.

By similar calculations, $x \notin T(M)$. Thus $T(M)$ is not a direct summand of M, and $M/T(M)$ is not flat.

References

1. F. W. ANDERSON and K. R. FULLER, *Rings and categories of modules*, 2nd edn, Graduate Texts in Mathematics 13 (Springer, New York, 1992).
2. P. M. COHN, *Free rings and their relations* (Academic Press, 1971).
3. N. V. DUNG and A. FACCHINI, 'Weak Krull–Schmidt for infinite direct sum of uniserial modules', *J. Algebra* 193 (1997) 102–121.
4. N. V. DUNG and A. FACCHINI, 'Direct summands of serial modules', *J. Pure Appl. Algebra* 133 (1998) 93–106.
5. A. FACCHINI, *Module theory: endomorphism rings and direct sum decompositions in some classes of modules*, Progress in Mathematics 167 (Birkhäuser, Basel, 1998).
6. C. FAITH, *Algebra.* II: *Ring theory*, Grundlehren der Mathematischen Wissenschaften 191 (Springer, Berlin, 1976).
7. L. FUCHS, *Infinite Abelian groups – Vol.* 1 (Academic Press, 1970).
8. I. HERZOG, 'A test for finite representation type', *J. Pure Appl. Algebra* 95 (1994) 151–182.
9. T. Y. LAM, *Lectures on modules and rings*, Graduate Texts in Mathematics 131 (Springer, New York, 1999).
10. J. C. MCCONNEL and J. C. ROBSON, *Noncommutative Noetherian rings*, Graduate Studies in Mathematics 30 (American Mathematical Society, Providence, RI, 2000) (revised edition).
11. M. PREST, *Model theory and modules*, London Mathematical Society Lecture Note Series 130 (Cambridge University Press, Cambridge, 1987).
12. G. PUNINSKI, *Serial rings* (Kluwer, Dordrecht, 2001).
13. G. PUNINSKI, 'Pure projective modules over a nearly simple uniserial domain and decompositions of uniserial modules', *J. Pure Appl. Algebra* 163 (2001) 319–337.
14. G. PUNINSKI, 'When a superdecomposable pure injective module over a serial ring exists', *Rings, modules and algebras. Proceedings of the Algebra Conference, Venice*, Lecture Notes in Pure and Applied Mathematics 236 (ed. A. Facchini, E. Houston and L. Salce, Marcel Dekker, 2004) 449–464.
15. G. PUNINSKI and P. ROTHMALER, 'When every finitely generated flat module is projective', *J. Algebra* 277 (2004) 542–558.
16. P. ROTHMALER, 'Mittag–Leffler modules and positive atomicity', Habilitationsschrift, Kiel, 1994.
17. P. ROTHMALER, 'Mittag–Leffler modules', *J. Pure Appl. Algebra* 88 (1997) 227–239.
18. I. SAKHAYEV, 'Bass's perfect rings and their generalizations', *Abelian Groups and Modules* (*Tomsk*) 10 (1991) 106–124.
19. A. A. TUGANBAEV, *Semidistributive rings and modules*, Mathematics and its Applications 449 (Kluwer, Dordrecht, 1998).
20. R. B. WARFIELD, 'Purity and algebraic compactness for modules', *Pacific J. Math.* 28 (1969) 699–719.
21. R. WISBAUER, *Foundations of module and ring theory*, Algebra, Logic and Application Series 3 (Gordon & Breach, 1991).

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press

Gena Puninski and Philipp Rothmaler
Department of Mathematics
Ohio State University
435 Galvin Hall
4240 Campus Drive
Lima
OH 45804
USA

puninskiy.1@osu.edu
rothmaler.1@osu.edu

https://doi.org/10.1112/S0024610705006290 Published online by Cambridge University Press