Forum of Mathematics, Pi (2026), **Vol. 14**:e16 1–4
doi:10.1017/fmp.2026.10033

CAMBRIDGE UNIVERSITY PRESS

RESEARCH ARTICLE

Optimal bounds in Bend-and-Break

Eric Jovinelly[1], Brian Lehmann[2] and Eric Riedl[3]

[1]Department of Mathematics, Brown University, Box 1917, 151 Thayer Street, Providence, RI, 02912, USA;
E-mail: eric_jovinelly@brown.edu.
[2]Department of Mathematics, Boston College, 140 Commonwealth Ave, Chestnut Hill, MA, 02467, USA;
E-mail: lehmannb@bc.edu (Corresponding author).
[3]Department of Mathematics, University of Notre Dame, 255 Hurley Hall, Notre Dame, IN, 46556, USA;
E-mail: eriedl@nd.edu.

Received: 24 December 2025; **Revised:** 6 March 2026; **Accepted:** 11 April 2026

2020 Mathematics Subject Classification: *Primary* – 14E30; *Secondary* – 14H10

Abstract
We improve the Bend-and-Break result of Miyaoka and Mori by establishing the optimal degree bound. Our result also yields optimal bounds on lengths of extremal rays of log canonical pairs.

Contents

1 Introduction **1**
1.1 Extremal rays 2
2 Breaking curves: low degree rational curves **2**
References **4**

1. Introduction

Mori's Bend-and-Break lemma is a fundamental tool for working with curves on projective varieties. Different versions of this important result have been established by (Mor79; Mor82; MM86; Kol96). Our main goal is to strengthen (MM86, Theorem 5) and to apply it to lengths of extremal rays, answering questions posed by (Kol96; Nik96; Mat02; Fuj11) (and others).

Theorem 1.1. *Let X be a projective variety over an algebraically closed field of arbitrary characteristic. Let H be a nef $\mathbb{R}$-Cartier divisor on X. Suppose there exists an irreducible curve $C \subset X$ contained in the smooth locus of X such that*

$$K_X \cdot C < 0.$$

Then for every closed point $x \in C$, there exists a rational curve R containing x such that

$$H \cdot R \leq (\dim X + 1)\frac{H \cdot C}{-K_X \cdot C}.$$

The constant $(\dim X + 1)$ in Theorem 1.1 improves the constant $(2 \dim X)$ given in (MM86, Theorem 5). Our improvement is optimal as we may let X be $\mathbb{P}^n$ and H be a hyperplane.

The proof of (MM86, Theorem 5) uses the fact that a one-dimensional family of maps $C \to X$ with a fixed point must break off a rational curve. Our key technical statement ("Bend-and-Shatter,"

© The Author(s), 2026. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

https://doi.org/10.1017/fmp.2026.10033 Published online by Cambridge University Press

Lemma 2.1) shows that a k-dimensional family of curves $C \to X$ that fixes k points must break off k rational curves. When combined with the reduction steps of (MM86, Theorem 5) and (Kol96, II.5.8 Theorem), we obtain a quick proof of Theorem 1.1.

1.1. Extremal rays

One of Mori's first applications for Bend-and-Break was the study of extremal rays of the pseudo-effective cone of curves. (Mor82, Theorem 1.4) proved that for a smooth projective variety X every K_X-negative extremal ray of the pseudo-effective cone contains a rational curve C satisfying $-K_X \cdot C \leq \dim X + 1$.

As a consequence of new advances in subadjunction theory, Kawamata proved a bound on the length of extremal rays for klt pairs (X, Δ) with $\mathbb{Q}$-coefficients. More precisely, (Kaw91) showed that the length of such an extremal ray is bounded by $2n$ where n is the dimension of a fiber in the exceptional locus of the corresponding contraction. This was extended to dlt pairs with $\mathbb{R}$-coefficients by Shokurov in the appendix to (Nik96) and by (BCHM10, Theorem 3.8.1). Using Theorem 1.1 in place of (MM86, Theorem 5) in these arguments, we obtain the optimal degree bound:

Theorem 1.2. *Let (X, Δ) be a dlt pair over an algebraically closed field of characteristic 0. Suppose that $\pi : X \to Z$ is the contraction of a $(K_X + \Delta)$-negative extremal face $\mathcal{R}$ of $\overline{\mathrm{NE}}(X)$. For any positive-dimensional irreducible component F of a fiber of π, there is a rational curve C in F satisfying:*

1. *The class of C is contained in the face $\mathcal{R}$.*
2. *The deformations of C sweep out F.*
3. $-(K_X + \Delta) \cdot C \leq \dim F + 1$.

If furthermore (X, Δ) is klt and π is a birational contraction, then we can ensure a strict inequality in (3).

The arguments of (Fuj11, Theorem 18.2) extend this result to lc pairs with $\mathbb{R}$-coefficients.

Theorem 1.3. *Let (X, Δ) be an lc pair over an algebraically closed field of characteristic 0. Then every $(K_X + \Delta)$-negative extremal ray of the pseudo-effective cone of curves is generated by a rational curve C with $-(K_X + \Delta) \cdot C \leq \dim X + 1$.*

Note that this length bound was known previously for toric varieties by (Fuj03) and in the setting of LCIQ singularities by (CT09).

2. Breaking curves: low degree rational curves

In this section, we establish Bend-and-Shatter and use it to prove Theorem 1.1. For a numerical curve class β on a projective variety X, we let $\overline{\mathcal{M}}_{g,n}(X, \beta)$ denote the Kontsevich moduli stack of stable maps and let $\mathcal{M}_{g,n}(X, \beta)$ denote the open substack of maps with smooth irreducible domain. We refer to (Kon95; BM96; FP97; AO01) for more details about the definition of these spaces and the structures that they carry (such as forgetful morphisms).

Lemma 2.1 (Bend-and-Shatter). *Let X be a projective variety over an algebraically closed field of arbitrary characteristic. Consider a non-negative integer g and a positive integer r such that $2g-2+r > 0$. Fix a stable irreducible marked curve $(C, q_1, \ldots, q_r) \in \mathcal{M}_{g,r}$.*

For some $k \leq r$, let $p_1, \ldots, p_k$ be points of X and let β be a numerical curve class. Suppose there exists a k-dimensional locally closed substack $S \subset \mathcal{M}_{g,r}(X, \beta)$ parametrizing pointed maps $s : (C, q_1, \ldots, q_r) \to X$ with $s(q_i) = p_i$ for all $i \leq k$. Then there exists a stable map $s' : (C', q'_1, \ldots, q'_r) \to X$ in the closure of S in $\overline{\mathcal{M}}_{g,r}(X, \beta)$ such that

- *$s'(q'_i) = p_i$ for all $i \leq k$;*
- *for each $i \leq k$ there is a tree of rational curves $C'_i \subset C'$ such that $q'_i \in C'_i$ and s' does not contract C'_i to a point; and*

https://doi.org/10.1017/fmp.2026.10033 Published online by Cambridge University Press

- *the stabilization of* $(C', q_1', \ldots, q_r')$ *is* $(C, q_1, \ldots, q_r)$. *In particular, the stabilization map* $(C', q_1', \ldots, q_r') \to (C, q_1, \ldots, q_r)$ *contracts* C_i' *to* $q_i \in C$.

Proof. Let U be the preimage of S in $\mathcal{M}_{g,r+k}(X, \beta)$ under the map $\pi : \overline{\mathcal{M}}_{g,r+k}(X, \beta) \to \overline{\mathcal{M}}_{g,r}(X, \beta)$ which forgets the last k points. Thus, U parametrizes maps in S together with the choice of k additional points $\{q_{r+i}\}_{i=1}^{k}$ in $C \setminus \{q_i\}_{i=1}^{r}$. We also let $\psi : \overline{\mathcal{M}}_{g,r+k}(X, \beta) \to \overline{\mathcal{M}}_{g,r+k}$ be the forgetful map and let $\phi : \overline{\mathcal{M}}_{g,r+k} \to \overline{\mathcal{M}}_{g,r}$ be the map forgetting the last k markings. Note that the fiber F of ϕ over $(C, q_1, \ldots, q_r)$ is irreducible since it is birational to C^k. By construction the closure of the image of U under ψ is F. Note that the nonempty fibers of $\psi : U \to F$ are isomorphic to S and thus have dimension k.

Fix a very ample line bundle $\mathcal{L}$ on X. Set $U_0 = U$ and for $1 \leq i \leq k$ define $U_i \subset \mathcal{M}_{g,r+k}(X, \beta)$ inductively by choosing a general section D_i of $\mathcal{L}$ and letting U_i be the intersection of U_{i-1} with $ev_{r+i}^{-1}(D_i)$ where ev_{r+i} denotes the evaluation map at the corresponding marked point. In particular we choose D_i so that $p_i \notin D_i$. We prove by induction that the dimension of the fiber of $\psi|_{U_i}$ over a general point of F is at least $k - i$.

By induction we know the fiber V_{i-1} of $\psi|_{U_{i-1}}$ over a general point $(C, q_1, \ldots, q_{r+k})$ of F has positive dimension. Note that the image of V_{i-1} in $\mathcal{M}_{g,0}(X, \beta)$ does not depend on the choice of marked points $q_{r+i}, \ldots, q_{r+k}$. Furthermore, this image must have positive dimension since V_{i-1} has positive dimension and parametrizes maps from a fixed marked curve. Thus for a general point of F the image $s(q_{r+i})$ sweeps out a locus of dimension ≥ 1 in X as we vary $s \in V_{i-1}$. We conclude that the preimage of the general divisor D_i under the evaluation map $ev_{r+i}|_{U_{i-1}}$ meets V_{i-1}. In particular the dimension of the general fiber of $\psi|_{U_i}$ is at most one less than the dimension of the general fiber of $\psi|_{U_{i-1}}$, proving the claim.

Let $\overline{U_k}$ be the closure of U_k in $\overline{\mathcal{M}}_{g,r+k}(X, \beta)$. There is an element of $\overline{U_k}$ lying over the locus in F where q_{r+i} specializes to q_i for each i. Let $s' : (C', q_1', \ldots, q_{r+k}') \to X$ be the corresponding stable map. Because $\pi(s')$ lies in the closure $\overline{S}$ of S in $\overline{\mathcal{M}}_{g,r}(X, \beta)$, we know that the stabilization of $(C', q_1', \ldots, q_r')$ must be $(C, q_1, \ldots, q_r)$. Thus each q_i' is contained in a tree of rational curves (which also contains q_{r+i}') that is contracted by the stabilization map. Likewise, because $\pi(s')$ lies in $\overline{S}$ we see that $s'(q_i') = p_i$ for all $i \leq k$. For all $i, j \leq k$, generality of D_j ensures it is disjoint from p_i. Because $s'(q_{r+i}')$ must lie in D_i, we see that the tree of rational curves containing q_i' has to map to a curve in X connecting p_i to D_i; in particular, some component is not contracted by s'. □

The next proposition relates the dimension of a family of curves to the number of rational curves that can be broken off using Lemma 2.1.

Proposition 2.2. *Let X be a projective variety and let C be a smooth projective curve of genus g over an algebraically closed field of arbitrary characteristic. Suppose $M \subset \mathrm{Mor}(C, X)$ is an irreducible locally closed subvariety. Set $k = \lfloor \frac{\dim M}{\dim X + 1} \rfloor$ and let $s : C \to X$ be any map parametrized by M. If $2g - 2 + k > 0$, then the closure of the image of M in $\overline{\mathcal{M}}_{g,0}(X, \beta)$ contains a point parametrizing a map $s' : C' \to X$ satisfying:*

- *C' consists of the union of C with at least k trees of rational curves, and*
- *at least k of these trees contain an irreducible component T such that s' realizes T as a noncontracted rational curve on X that passes through a general point of $s(C)$.*

Proof. Let $q_1, \ldots, q_k \in C$ be k general points in C. Let $T_i = \{\tilde{s} \in M \mid \tilde{s}(q_i) = s(q_i)\}$. Since $s \in T_i$ for all i, the intersection $S := \cap_i T_i$ is nonempty. Moreover, as $\mathrm{codim}(T_i, M) \leq \dim X$, we get $\mathrm{codim}(S, M) \leq k(\dim X)$. Thus $\dim S \geq k$.

Because $2g - 2 + k > 0$, the pointed curve $(C, q_1, \ldots, q_k)$ has finite automorphism group, and hence the natural map $\pi : S \to \overline{\mathcal{M}}_{g,k}(X, \beta)$ is generically finite. Apply Lemma 2.1 to a dense locally closed subset of $\pi(S)$ and let $s' : (C', q_1', \ldots, q_k') \to X$ be the stable map it identifies. The desired stable map is obtained from s' by forgetting the k marked points and stabilizing. □

We are now equipped to prove Theorem 1.1 via a dimension counting argument.

https://doi.org/10.1017/fmp.2026.10033 Published online by Cambridge University Press

Proof of Theorem 1.1. First suppose that our ground field is algebraically closed of characteristic $p > 0$ and that H is $\mathbb{Q}$-Cartier. After rescaling H we may suppose it is Cartier. We write $i : C' \to X$ for the normalization of C. For $m > 0$, let $s_m : C' \to X$ be the precomposition of i with the m^{th} iterate of the Frobenius. The dimension d_m of $\mathrm{Mor}(C', X)$ at s_m satisfies

$$d_m \geq p^m(-K_X \cdot i_* C') - g \dim X,$$

where g is the genus of C'. Let $k_m = \lfloor \frac{d_m}{\dim X + 1} \rfloor$. For large m, Proposition 2.2 allows us to find a deformation of s_m that breaks off k_m rational curves through k_m general points of $s(C')$. Because H is nef, at least one of these rational curves has H-degree at most

$$\frac{H \cdot s_{m*} C'}{k_m} = \frac{p^m (H \cdot i_* C')}{k_m} \leq \frac{p^m (H \cdot i_* C')}{p^m(-K_X \cdot i_* C') - (g+1)\dim X} (\dim X + 1).$$

For large enough m the floor of this upper bound is at most $(\dim X + 1)\frac{H \cdot i_* C'}{-K_X \cdot i_* C'}$. This proves that a general point of $s(C')$ is contained in a rational curve whose H-degree satisfies the desired inequality. Since the existence of such a rational curve through a point is a closed condition, this statement holds for every closed point in $s(C')$, and in particular for x.

The extension to ample $\mathbb{Q}$-Cartier divisors in characteristic 0 uses the spreading out argument of (MM86, Step 3 of proof of Theorem 5). The extension to nef $\mathbb{R}$-Cartier divisors follows as in (Kol96, Steps 4 and 5 of proof of II.5.8 Theorem). □

Acknowledgements. We thank Izzet Coskun, Brendan Hassett, James McKernan, Joaquín Moraga, and Sho Tanimoto for helpful conversations. We are grateful to Yuchen Liu for suggesting we think about lengths of extremal rays.

Competing interests. The author have no competing interests to declare.

Funding statement. Eric Jovinelly was supported by an NSF postdoctoral research fellowship, DMS-2303335. Brian Lehmann was supported by Simons Foundation grant award number 851129. Eric Riedl was supported by NSF CAREER grant DMS-1945944 and Simons Foundation grants 00011850 and 00013673.

References

[AO01] D. Abramovich and F. Oort, 'Stable maps and Hurwitz schemes in mixed characteristics', in *Advances in Algebraic Geometry Motivated by Physics (Lowell, MA, 2000)*, volume 276 of *Contemp. Math.* (Amer. Math. Soc., Providence, RI, 2001), 89–100.

[BCHM10] C. Birkar, P. Cascini, C. D. Hacon and J. McKernan, 'Existence of minimal models for varieties of log general type', *J. Amer. Math. Soc.* **23**(2) (2010), 405–468.

[BM96] K. Behrend and Yu. Manin, 'Stacks of stable maps and Gromov-Witten invariants', *Duke Math. J.* **85**(1) (1996), 1–60.

[CT09] J.-C. Chen and H.-H. Tseng, 'Cone theorem via Deligne-Mumford stacks', *Math. Ann.* **345**(3) (2009), 525–545.

[FP97] W. Fulton and R. Pandharipande, 'Notes on stable maps and quantum cohomology', in *Algebraic Geometry—Santa Cruz 1995*, volume 62, Part 2 of *Proc. Sympos. Pure Math.* (Amer. Math. Soc., Providence, RI, 1997), 45–96.

[Fuj03] O. Fujino, 'Notes on toric varieties from Mori theoretic viewpoint', *Tohoku Math. J. (2)*, **55**(4) (2003), 551–564.

[Fuj11] O. Fujino, 'Fundamental theorems for the log minimal model program', *Publ. Res. Inst. Math. Sci.* **47**(3) (2011), 727–789.

[Kaw91] Y. Kawamata, 'On the length of an extremal rational curve', *Invent. Math.* **105**(3) (1991), 609–611.

[Kol96] J. Kollár, *Rational Curves on Algebraic Varieties*, volume 32 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* (Springer-Verlag, Berlin, 1996).

[Kon95] M. Kontsevich, 'Enumeration of rational curves via torus actions', in *The Moduli Space of Curves (Texel Island, 1994)*, volume 129 of *Progr. Math.* (Birkhäuser Boston, Boston, MA, 1995), 335–368.

[Mat02] K. Matsuki, *Introduction to the Mori Program*. Universitext (Springer-Verlag, New York, 2002).

[MM86] Y. Miyaoka and S. Mori, 'A numerical criterion for uniruledness', *Ann. of Math. (2)* **124**(1) (1986), 65–69.

[Mor79] S. Mori, 'Projective manifolds with ample tangent bundles', *Ann. of Math. (2)* 110(3) (1979), 593–606.

[Mor82] S. Mori, 'Threefolds whose canonical bundles are not numerically effective', *Ann. of Math. (2)* **116**(1) (1982), 133–176.

[Nik96] V. V. Nikulin, 'The diagram method for 3-folds and its application to the Kähler cone and Picard number of Calabi-Yau 3-folds. I', in *Higher-dimensional Complex Varieties (Trento, 1994)* (de Gruyter, Berlin, 1996), 261–328. With an appendix by Vyacheslav V. Shokurov.

https://doi.org/10.1017/fmp.2026.10033 Published online by Cambridge University Press