Forum of Mathematics, Sigma (2025), **Vol. 13**:e77 1–15
doi:10.1017/fms.2024.143

CAMBRIDGE UNIVERSITY PRESS

RESEARCH ARTICLE

Global F-regularity for weak del Pezzo surfaces

Tatsuro Kawakami [1] and Hiromu Tanaka [2]

[1]Department of Mathematics, Graduate School of Science, Kyoto University, 606-8502, Japan
E-mail: tkawakami@math.kyoto-u.ac.jp (corresponding author).
[2]Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914, Japan;
E-mail: tanaka@ms.u-tokyo.ac.jp.

Received: 29 April 2024; **Revised:** 5 August 2024; **Accepted:** 17 November 2024

2020 Mathematics Subject Classification: *Primary* – 14J45; *Secondary* – 13A35

Abstract

Let k be an algebraically closed field of characteristic $p > 0$. Let X be a normal projective surface over k with canonical singularities whose anticanonical divisor is nef and big. We prove that X is globally F-regular except for the following cases: (1) $K_X^2 = 4$ and $p = 2$, (2) $K_X^2 = 3$ and $p \in \{2, 3\}$, (3) $K_X^2 = 2$ and $p \in \{2, 3\}$, (4) $K_X^2 = 1$ and $p \in \{2, 3, 5\}$. For each degree K_X^2, the assumption of p is optimal.

Contents

1 Introduction **1**
2 Preliminaries **3**
2.1 Notation and terminology . . . 3
2.1.1 Singularities in minimal model program . . . 3
2.1.2 Weak del Pezzo surfaces . . . 4
2.2 F-splitting and global F–regularity . . . 4
2.3 Reflexive Cartier operator . . . 6
3 Proofs of main theorems **7**
3.1 Criterion of the F-splitting of klt surfaces . . . 7
3.2 Global F-splitting: Proof of Theorem B . . . 8
3.3 Global F-regularity: Proof of Theorem A . . . 13
References **14**

1. Introduction

We work over an algebraically closed field of characteristic $p > 0$. Fano varieties play a significant role in the classification of algebraic varieties. In positive characteristic, properties defined by the Frobenius morphism such as (global) F-splitting or global F-regularity are useful. Therefore, it is natural to ask when Fano varieties are F-split or globally F-regular. For smooth del Pezzo surfaces, Hara [Har98a] proved the following result:

Theorem 1.1 [Har98a, Example 5.5]. *Let X be a smooth del Pezzo surface over an algebraically closed field of characteristic $p > 0$. Then X is globally F-regular except for the following cases:*

© The Author(s), 2025. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

(1) $K_X^2 = 3$ *and* $p = 2$.
(2) $K_X^2 = 2$ *and* $p \in \{2, 3\}$.
(3) $K_X^2 = 1$ *and* $p \in \{2, 3, 5\}$.

Our aim is to generalize Hara's result to the case when $-K_X$ is nef and big, or equivalently, X is canonical. Here, we say that a variety is *canonical* if it has only canonical singularities. In fact, the following theorem holds.

Theorem A. *Let k be an algebraically closed field of characteristic $p > 0$. Let X be a canonical projective surface over k whose anticanonical divisor is nef and big. Then X is globally F-regular except for the following cases:*

(1) $K_X^2 = 4$ *and* $p = 2$.
(2) $K_X^2 = 3$ *and* $p \in \{2, 3\}$.
(3) $K_X^2 = 2$ *and* $p \in \{2, 3\}$.
(4) $K_X^2 = 1$ *and* $p \in \{2, 3, 5\}$.

Theorem A has an important role for investigating of global F-regularity of smooth Fano threefolds and smooth del Pezzo varieties (see [KT24b, KT24c] for details).

Remark 1.2. For each degree K_X^2, the assumption on p is optimal. In fact, for each case listed above, there exists a canonical del Pezzo surface that is not strongly F-regular (see [KN23, Table 1] and [KT24a, Table 1]).

The assumption on p is still optimal even if we assume that X is smooth since taking the minimal resolution does not change the degree and globally F-regularity (Corollary 2.5). Similarly, even if we replace 'nef and big' by 'ample', the conclusion of Theorem A is the same. This is because X is globally F-regular if and only if its anticanonical model is globally F-regular (Proposition 2.4). In particular, Theorem A does not imply Theorem 1.1.

Remark 1.3. For each prime number p, there exists a Kawamata log terminal (klt) del Pezzo surface X (i.e., a normal projective surface such that $(X, 0)$ is klt and $-K_X$ is ample) that is not F-split [CTW18, Theorem 1.1].

We now focus on the proof of Theorem A. We first investigate when X as in Theorem A is F-split. The proof is divided into two cases: the case where $K_X^2 \geq 5$ and the case where $K_X^2 \leq 4$.

First, we consider the case where $K_X^2 \geq 5$. We may assume that X is obtained by taking a blowup $f\colon X \to \mathbb{P}^2$ along some points. Recall that if there exists an effective divisor $\Delta_{\mathbb{P}^2}$ on $\mathbb{P}^2$ such that the divisor Δ on X defined by $K_X + \Delta = f^*(K_{\mathbb{P}^2} + \Delta_{\mathbb{P}^2})$ is effective, then the following holds (Proposition 2.4):

$$(\mathbb{P}^2, \Delta_{\mathbb{P}^2}) \text{ is } F\text{-split} \Leftrightarrow (X, \Delta) \text{ is } F\text{-split} \Rightarrow X \text{ is } F\text{-split}.$$

Such a divisor $\Delta_{\mathbb{P}^2}$ can be found by utilizing an inversion of adjunction for F-splitting (Proposition 2.6). However, since we can only assume that the blowup points are in *almost general position*, the situation is more complicated than the smooth del Pezzo cases, which are obtained by blowing up points in *general position*. For more details, see Proposition 3.8.

Next, we consider the case where $K_X^2 \leq 4$. In the case of smooth del Pezzo surfaces, Hara [Har98a] investigated F-splitting using the following two steps:

(i) The reduction of F-splitting to the vanishing $H^1(X, \Omega_X^1(pK_X)) = 0$ via the Cartier operator.
(ii) Embedding X as a hypersurface or a complete intersection of a weighted projective space and proving the vanishing $H^1(X, \Omega_X^1(pK_X)) = 0$.

For a smooth weak del Pezzo surface X, we can also reduce the F-splitting of X to the vanishing $H^1(X, \Omega_X^1(pK_X)) = 0$. However, Step (ii), proving $H^1(X, \Omega_X^1(pK_X)) = 0$, is not easy because X is not embedded in a weighted projective space via $|-mK_X|$ for $m \in \mathbb{Z}_{>0}$. Therefore, by replacing X with its anticanonical model, we embed X into a weighted projective space via $|-mK_X|$. However, in this case, Step (i), the reduction of the F-splitting of X to the vanishing $H^1(X, \Omega_X^1(pK_X)) = 0$, is not

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

straightforward since Cartier operator is defined on smooth schemes. To address this issue, we use the reflexive Cartier operator introduced in [Kaw22b]. Indeed, we utilize the fact that the reflexive Cartier operator behaves well on F-pure klt surfaces (Lemma 3.1).

Combining the above results, we obtain the following theorem.

Theorem B. *Let k be an algebraically closed field of characteristic $p > 0$. Let X be a canonical F-pure projective surface over k whose anticanonical divisor is ample. Then X is F-split except for the following cases:*

(1) $K_X^2 = 3$ *and* $p = 2$.
(2) $K_X^2 = 2$ *and* $p \in \{2, 3\}$.
(3) $K_X^2 = 1$ *and* $p \in \{2, 3, 5\}$.

Remark 1.4.

(1) When $K_X^2 = 4$ and $p = 2$, there exists a a canonical del Pezzo surface with D_5^0-singularity, which is not F-pure [KN23, Proposition 3.16].
(2) When $K_X^2 = 3$ and $p = 3$, there exists a canonical del Pezzo surface with E_6^0-singularity, which is not F-pure [KN23, Proposition 3.22].

Finally, we now overview how to deduce Theorem A from Theorem B. Let X be a canonical weak del Pezzo surface. Replacing X with its anticanonical model, we can assume that $-K_X$ is ample. For each degree K_X^2, we will find an optimal bound on p that ensures all the singularities of X are strongly F-regular (see Lemma 3.9). We then conclude by the following well-known fact that asserts the equivalence between F-splitting and global F-regularity for strongly F-regular $\mathbb{Q}$-Gorenstein Fano varieties:

Theorem 1.5 (cf. [KT23, Proof of Theorem 6.2]). *Let X be a normal projective variety such that $-K_X$ is an ample $\mathbb{Q}$-Cartier $\mathbb{Z}$-divisor. Suppose that X is strongly F-regular. If X is F-split, then X is globally F-regular.*

2. Preliminaries

2.1. Notation and terminology

In this subsection, we summarize notation and basic definitions used in this article.

(1) Throughout the paper, p denotes a prime number and we work over an algebraically closed field k of characteristic $p > 0$. We set $\mathbb{F}_p := \mathbb{Z}/p\mathbb{Z}$. We denote by $F\colon X \to X$ the absolute Frobenius morphism on an $\mathbb{F}_p$-scheme X.
(2) We say that X is a *variety* (over k) if X is an integral scheme that is separated and of finite type over k. We say that X is a *curve* (resp. *surface*) if X is a variety of dimension one (resp. two).
(3) For a variety X, we define the *function field* $K(X)$ of X as the stalk $\mathcal{O}_{X,\xi}$ at the generic point ξ of X.
(4) We say that a $\mathbb{Q}$-divisor D on a normal variety X is *simple normal crossing* if for every point $x \in \operatorname{Supp} D$, the local ring $\mathcal{O}_{X,x}$ is regular and there exists a regular system of parameters $x_1, \dots, x_d$ of the maximal ideal $\mathfrak{m}$ of $\mathcal{O}_{X,x}$ and $1 \le r \le d$ such that $\operatorname{Supp}(D|_{\operatorname{Spec} \mathcal{O}_{X,x}}) = \operatorname{Spec}(\mathcal{O}_{X,x}/(x_1 \cdots x_r))$.
(5) Given a variety X, a projective birational morphism $\pi\colon Y \to X$ is called a *log resolution of* X if Y is a smooth variety and $\operatorname{Exc}(f)$ is a simple normal crossing divisor.
(6) Given a variety X and a closed subscheme Z, we denote by $\operatorname{Bl}_Z X$ the blowup of X along Z.
(7) Given a normal variety X and a $\mathbb{Z}$-divisor D on X, we define a reflexive sheaf $\Omega_X^{[i]}(D)$ by $j_*(\Omega_U^i \otimes \mathcal{O}_U(D))$, where $j\colon U \hookrightarrow X$ is the open immersion from the smooth locus U of X.

2.1.1. Singularities in minimal model program

For the definitions of singularities in minimal model program (e.g., canonical and klt), we refer to [KM98, Section 2.3]. Take a normal surface X. Let $f\colon Y \to X$ be the minimal resolution. We only need the following characterizations in this paper.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

(1) X is canonical if and only if K_X is $\mathbb{Q}$-Cartier and $K_Y \sim_{\mathbb{Q}} f^*K_X$.
(2) X is klt if and only if K_X is $\mathbb{Q}$-Cartier, f is a log resolution of X and all the coefficients of the $\mathbb{Q}$-divisor Γ defined by $K_Y + \Gamma \sim_{\mathbb{Q}} f^*K_X$ are < 1.

By definition, we have

$$\text{canonical} \Rightarrow \text{klt}.$$

Moreover, the following implications hold for the surface case:

- canonical $\Rightarrow$ Gorenstein.
- klt $\Rightarrow$ $\mathbb{Q}$-factorial.

2.1.2. Weak del Pezzo surfaces

Given a normal projective Gorenstein surface X, we say that X is *del Pezzo* (resp. *weak del Pezzo*) if $-K_X$ is ample (resp. nef and big). In what follows, we summarize some properties on (weak) del Pezzo surfaces for later usage.

Let Z be a canonical weak del Pezzo surface. The *anticanonical model* Y of Z is defined as the Stein factorisation of the morphism $\varphi_{|-mK_Z|}\colon Z \to \mathbb{P}^{h^0(Z,-mK_Z)-1}$ induced by the complete linear system $|-mK_Z|$, where m is a positive integer such that $|-mK_X|$ is base point free (whose existence is guaranteed by [Tan15, Theorem 0.4]). Then Y is canonical, because have $K_Z \sim h^*K_Y$ for the induced morphism $h\colon Z \to Y$. Moreover, h is obtained by contracting all the (-2)-curves on Y. In particular, the minimal resolution X of Z coincides with the minimal resolution of Y:

$$f\colon X \xrightarrow{g} Z \xrightarrow{h} Y.$$

Moreover, X is a smooth weak del Pezzo surface.

There is a natural one-to-one correspondence between

- smooth weak del Pezzo surfaces and
- canonical del Pezzo surfaces.

Indeed, if Y is a canonical del Pezzo surface, then its minimal resolution X is a smooth weak del Pezzo surface. Conversely, given a smooth weak del Pezzo surface X, its anticanonical model Y is a canonical del Pezzo surface.

2.2. F-splitting and global F–regularity

In this subsection, we gather basic facts on F-splitting and global F-regularity.

Definition 2.1. Let X be a normal variety, and let Δ be an effective $\mathbb{Q}$-divisor on X.

(1) We say that (X, Δ) is *F-split* if

$$\mathcal{O}_X \xrightarrow{F^e} F^e_*\mathcal{O}_X \hookrightarrow F^e_*\mathcal{O}_X(\lfloor (p^e - 1)\Delta \rfloor)$$

splits as an $\mathcal{O}_X$-module homomorphism for every $e \in \mathbb{Z}_{>0}$.
(2) We say that (X, Δ) is *sharply F-split* if

$$\mathcal{O}_X \xrightarrow{F^e} F^e_*\mathcal{O}_X \hookrightarrow F^e_*\mathcal{O}_X(\lceil (p^e - 1)\Delta \rceil)$$

splits as an $\mathcal{O}_X$-module homomorphism for some $e \in \mathbb{Z}_{>0}$.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

(3) We say that (X, Δ) is *globally F-regular* if, given an effective $\mathbb{Z}$-divisor E, there exists $e \in \mathbb{Z}_{>0}$ such that

$$\mathcal{O}_X \xrightarrow{F^e} F^e_* \mathcal{O}_X \hookrightarrow F^e_* \mathcal{O}_X(\lceil (p^e - 1)\Delta \rceil + E)$$

splits as an $\mathcal{O}_X$-module homomorphism.

We say that X is *F-split* (resp. *globally F-regular*) if so is $(X, 0)$.

Remark 2.2. We have the following implications:

$$\text{globally } F\text{-regular} \Longrightarrow \text{sharply } F\text{-split} \Longrightarrow F\text{-split}$$

where the former implication is easy and the latter one holds by the same argument as in [Sch08, Proposition 3.3]. Moreover, if the condition $(\star)$ holds, then (X, Δ) is sharply F-split if and only if (X, Δ) is F-split.

$(\star)$ $(p^e - 1)\Delta$ is a $\mathbb{Z}$-divisor for some $e \in \mathbb{Z}_{>0}$.

In particular, X is F-split if and only if $F\colon \mathcal{O}_X \to F_*\mathcal{O}_X$ splits as an $\mathcal{O}_X$-module homomorphism. In this paper, we only treat the case when $(\star)$ holds, and hence being F-split is equivalent to being sharply F-split. For more foundational properties, we refer to [SS10].

We shall also use the local versions of F-splitting and global F-regularity.

Definition 2.3. Given a normal variety X, we say that X is *F-pure* (resp. *strongly F-regular*) if there exists an open cover $X = \bigcup_{i \in I} X_i$ such that X_i is F-split (resp. globally F-regular) for every $i \in I$.

In what follows, we summarize some F-splitting criteria, which are well known to experts.

Proposition 2.4. *Let $f : X \to Y$ be a birational morphism of normal projective varieties. Take an effective $\mathbb{Q}$-divisor Δ_Y on Y such that $(p^e - 1)(K_Y + \Delta_Y)$ is Cartier for some $e \in \mathbb{Z}_{>0}$. Assume that the $\mathbb{Q}$-divisor Δ defined by $K_X + \Delta = f^*(K_Y + \Delta_Y)$ is effective. Then (X, Δ) is F-split (resp. globally F-regular) if and only if (Y, Δ_Y) is F-split (resp. globally F-regular).*

Proof. If (X, Δ) is F-split, then so is (Y, Δ_Y), which can be checked by taking the push-forward. As for the opposite implication, the same argument as in the first paragraph of the proof of [HX15, Proposition 2.11] works. □

Corollary 2.5. *Let Y be a canonical projective surface, and let $f : X \to Y$ be its minimal resolution. Then X is F-split (resp. globally F-regular) if and only if Y is F-split (resp. globally F-regular).*

Proof. The assertion immediately follows from Proposition 2.4 by using $K_X = f^*K_Y$. □

Proposition 2.6. *Let X be a normal projective Gorenstein variety. Take a normal prime Cartier divisor S and an effective $\mathbb{Q}$-Cartier $\mathbb{Q}$-divisor B on X such that $S \not\subset \operatorname{Supp} B$. Assume that*

(1) *$(S, B|_S)$ is F-split, and*
(2) *there is a positive integer $e \in \mathbb{Z}_{>0}$ such that $(p^e - 1)(K_X + S + B)$ is Cartier and*

$$H^1(X, \mathcal{O}_X(-S - (p^e - 1)(K_X + S + B))) = 0.$$

Then $(X, S + B)$ is F-split.

Proof. The same argument as in [CTW17, Lemma 2.7] works. □

Example 2.7. We now summarize some easy cases for later usage, although all of them are well known to experts,

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

(1) If $P, Q \in \mathbb{P}^1$ are distinct points, then $(\mathbb{P}^1, P+Q)$ is F-split (Proposition 2.6).
(2) Let L, L', L'' be three lines on $\mathbb{P}^2$ such that $L + L' + L''$ is simple normal crossing. Then $(\mathbb{P}^2, L + L' + L'')$ is F-split by (1) and Proposition 2.6.
(3) For each $i \in \{1, 2\}$, let F_i and F_i' be distinct fibers of the i-th projection $\mathrm{pr}_i \colon \mathbb{P}^1 \times \mathbb{P}^1 \to \mathbb{P}^1$. Then $(\mathbb{P}^1 \times \mathbb{P}^1, F_1 + F_1' + F_2 + F_2')$ is F-split by (1) and Proposition 2.6.

2.3. *Reflexive Cartier operator*

Throughout this subsection, we use the following convention unless stated otherwise.

Convention 2.8. *Let X be a normal variety and D a $\mathbb{Z}$-divisor on X. Let U be the smooth locus of X and $j \colon U \hookrightarrow X$ the inclusion. By abuse of notation, $D|_U$ is denoted by D.*

The Frobenius pushforward of the de Rham complex

$$F_*\Omega_U^\bullet \colon F_*\mathcal{O}_U \xrightarrow{F_*d} F_*\Omega_U^1 \xrightarrow{F_*d} \cdots$$

is a complex of $\mathcal{O}_U$-modules. Tensoring with $\mathcal{O}_U(D)$, we obtain a complex

$$F_*\Omega_U^\bullet \colon F_*\mathcal{O}_U(pD) \xrightarrow{F_*d \otimes \mathcal{O}_U(D)} F_*\Omega_U^1(pD) \xrightarrow{F_*d \otimes \mathcal{O}_U(D)} \cdots.$$

We define coherent $\mathcal{O}_U$-modules $B_U^i(pD)$ and $Z_U^i(pD)$ by

$$B_U^i(pD) \coloneqq \mathrm{Im}(F_*d \otimes \mathcal{O}_U(D) \colon F_*\Omega_U^{i-1}(pD) \to F_*\Omega_U^i(pD)),$$
$$Z_U^i(pD) \coloneqq \mathrm{Ker}(F_*d \otimes \mathcal{O}_U(D) \colon F_*\Omega_U^i(pD) \to F_*\Omega_U^{i+1}(pD)),$$

for all $i \geq 0$. Then $B_U^i(pD)$ and $Z_U^i(pD)$ are locally free [Kaw22b, Lemma 3.2]. When $D = 0$, we simply denote $B_U^i(pD)$ and $Z_U^i(pD)$ by B_U^i and Z_U^i, respectively. Then $B_U^i(pD) = B_U^i \otimes \mathcal{O}_U(D)$ and $Z_U^i(pD) = Z_U^i \otimes \mathcal{O}_U(D)$ holds [Kaw22b, Remark 3.3]. In particular, we note that $B_U^i(pD)$ and $Z_U^i(pD)$ do not mean $B_U^i \otimes \mathcal{O}_U(pD)$ and $Z_U^i \otimes \mathcal{O}_U(pD)$.

By [Kaw22b, Lemma 3.2], there exists an exact sequence

$$0 \to B_U^i(pD) \to Z_U^i(pD) \xrightarrow{C_U^i(D)} \Omega_U^i(D) \to 0, \tag{2.8.1}$$

and the map $C_U^i(D)$ coincides with $C_U^i \otimes \mathcal{O}_U(D)$, where C_U^i is the usual Cartier operator.

Definition 2.9. We define reflexive $\mathcal{O}_X$-modules $B_X^{[i]}(pD)$ and $Z_X^{[i]}(pD)$ by

$$B_X^{[i]}(pD) \coloneqq j_*B_U^i(pD) \text{ and}$$
$$Z_X^{[i]}(pD) \coloneqq j_*Z_U^i(pD)$$

for all $i \geq 0$. The *i-th reflexive Cartier operator*

$$C_X^{[i]}(D) \colon Z_X^{[i]}(pD) \to \Omega_X^{[i]}(D)$$

associated to D is defined as $j_*C_U^i(D)$ for all $i \geq 0$.

Lemma 2.10. *There exist the following exact sequences:*

$$0 \to Z_X^{[i]}(pD) \to F_*\Omega_X^{[i]}(pD) \xrightarrow{d'} B_X^{[i+1]}(pD), \tag{2.10.1}$$

$$0 \to B_X^{[i]}(pD) \to Z_X^{[i]}(pD) \xrightarrow{C_X^{[i]}(D)} \Omega_X^{[i]}(D), \tag{2.10.2}$$

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

for all $i \geq 0$. *Moreover,* $d'|_U \colon F_*\Omega_X^{[i]}(pD)|_U \to B_X^{[i+1]}(pD)|_U$ *and* $C_X^{[i]}(D)|_U \colon Z_X^{[i]}(pD)|_U \to \Omega_X^{[i]}(D)|_U$ *are surjective, and the homomorphism* $C_X^{[i]}(D)|_U$ *coincides with* $C_U^i \otimes \mathcal{O}_U(D)$.

Proof. Taking $B = 0$ in [Kaw22b, Lemma 3.5], we obtain the assertion. □

Remark 2.11. Taking $i = 0$ in equation (2.10.1), we obtain an exact sequence

$$0 \to \mathcal{O}_X(D) \to F_*\mathcal{O}_X(pD) \to B_X^{[1]}(pD),$$

and the first map is induced by the Frobenius homomorphism. In particular,

$$B_X^{[1]}(pD) = j_*\mathrm{Coker}(F \colon \mathcal{O}_U(D) \to F_*\mathcal{O}_U(pD))$$

holds.

3. Proofs of main theorems

3.1. Criterion of the F-splitting of klt surfaces

In this subsection, we provide a criterion for the F-splitting of klt surfaces (Proposition 3.2).

Lemma 3.1. *Let X be an F-pure klt surface and D a $\mathbb{Z}$-divisor. Then the sequence*

$$0 \to B_X^{[1]}(pD) \to Z_X^{[1]}(pD) \xrightarrow{C_X^{[1]}(D)} \Omega_X^{[1]}(D) \to 0 \tag{3.1.1}$$

is exact.

Proof. It is enough to show that $C_X^{[1]}(D)$ is surjective, as the other parts has been settled in Lemma 2.10. Since the assertion is local on X, we may assume that X is affine and has a unique singular point Q. If $p \neq 5$ or the singularity Q is not rational double point (RDP) of type E_8^1, then X is F-liftable by [KT24a, Theorem A]. Then the surjectivity of $C^{[1]}(D)$ follows from [Kaw22b, Lemma 3.8].

Suppose that $p = 5$ and the singularity Q is of type E_8^1. Then we may assume that $D = 0$ by [Lip69, Section 24] (see also [LMM21, Table 2]). Then the desired surjectivity follows from [Kaw22b, Proposition 4.4] and [KT24a, Theorem B]. □

Proposition 3.2. *Let X be an F-pure klt projective surface. Suppose that the following conditions hold:*

(1) $H^0(X, \Omega_X^{[1]}(K_X)) = 0$.
(2) $H^1(X, \Omega_X^{[1]}(pK_X)) = 0$.
(3) $H^0(X, \mathcal{O}_X((p+1)K_X)) = 0$.

Then X is F-split.

Proof. Recall that X is F-split if and only if the evaluation map

$$\mathrm{Hom}_{\mathcal{O}_X}(F_*\mathcal{O}_X, \mathcal{O}_X) \xrightarrow{F^*} \mathrm{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{O}_X)(\cong H^0(X, \mathcal{O}_X))$$

is surjective. Then, as in [BK05, 1.3.9 Remarks (ii)], the surjectivity is equivalent to the injectivity of the map

$$F \colon H^2(X, \mathcal{O}_X(K_X)) \to H^2(X, \mathcal{O}_X(pK_X)) \tag{3.2.1}$$

induced by Frobenius by Serre duality. Let U be the smooth locus of X. Since U is F-pure, the exact sequence

$$0 \to \mathcal{O}_U \to F_*\mathcal{O}_U \to B_U^1 \to 0$$

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

splits locally by definition. Tensoring with $\mathcal{O}_U(K_X)$, we obtain a locally split exact sequence

$$0 \to \mathcal{O}_U(K_U) \to F_*\mathcal{O}_U(pK_U) \to B_U^{[1]}(pK_U) \to 0.$$

Since the above exact sequence splits locally, taking pushforward preserves exactness on the right. Thus, taking the pushforward by the inclusion $U \hookrightarrow X$, we obtain the following locally split exact sequence

$$0 \to \mathcal{O}_X(K_X) \to F_*\mathcal{O}_X(pK_X) \to B_X^{[1]}(pK_X) \to 0.$$

Therefore, for the injectivity of equation (3.2.1), it suffices to show that $H^1(X, B_X^{[1]}(pK_X)) = 0$. By Lemma 3.1 and the condition (1), it is enough to prove that $H^1(X, Z_X^{[1]}(pK_X)) = 0$. By equation (2.10.1), we have an exact sequence

$$0 \to Z_X^{[1]}(pK_X) \to F_*\Omega_X^{[1]}(pK_X) \to B_X^{[2]}(pK_X).$$

Let $\mathcal{B} \coloneqq \mathrm{Im}(F_*\Omega_X^{[1]}(pK_X) \to B_X^{[2]}(pK_X))$. Then we obtain an exact sequence

$$H^0(X, \mathcal{B}) \to H^1(X, Z_X^{[1]}(pK_X)) \to H^1(X, \Omega_X^{[1]}(pK_X)) \overset{(2)}{=} 0.$$

Since we have

$$\mathcal{B} \subset B_X^{[2]}(pK_X) \subset F_*\Omega_X^{[2]}(pK_X) = H^0(X, \mathcal{O}_X((p+1)K_X)) \overset{(3)}{=} 0,$$

we conclude that $H^1(X, Z_X^{[1]}(pK_X)) = 0$. □

The condition (3) of Proposition 3.2 is satisfied if $-K_X$ is big. In what follows, we see when the condition (1) of Proposition 3.2 is satisfied.

Definition 3.3 (Log liftability). Let X be a normal projective surface. We say that X is *log liftable* if there exists a log resolution $f\colon Y \to X$ of X such that $(Y, \mathrm{Exc}(f))$ lifts to the ring $W(k)$ of Witt vectors. For the definition of liftability of a log smooth pair, we refer to [Kaw22a, Definition 2.6].

Lemma 3.4. *Let X be a normal projective F-pure surface such that $-K_X$ is a nef and big $\mathbb{Q}$-Cartier $\mathbb{Z}$-divisor. Then X is log liftable if and only if $H^0(X, \Omega_X^{[1]}(K_X)) = 0$.*

Proof. Since $H^2(X, \mathcal{O}_X) \cong H^0(X, \mathcal{O}_X(K_X)) = 0$, the 'if' direction is [Kaw22c, Theorem 2.8]. We prove the 'only if' direction. Let $f\colon Y \to X$ be a log resolution such that $(Y, E \coloneqq \mathrm{Ex}(f))$ lifts to $W(k)$. Since $f_*(\Omega_Y^1(\log E) \otimes \mathcal{O}_Y(f^*K_X)) = \Omega_X^{[1]}(K_X)$ by [KT24a, Theorem B], we have $H^0(X, \Omega_X^{[1]}(K_X)) = H^0(Y, \Omega_Y^1(\log E) \otimes \mathcal{O}_Y(f^*K_X))$. Then the vanishing follows from [Kaw22a, Theorem 2.11]. □

3.2. Global F-splitting: Proof of Theorem B

In the following proposition, we investigate F-splitting of F-pure canonical del Pezzo surfaces. For the proof, we confirm when the condition (2) of Proposition 3.2 is satisfied.

Proposition 3.5. *Let X be an F-pure canonical del Pezzo surface. Suppose that one of the following holds.*

(1) $K_X^2 = 1$ *and* $p > 5$.
(2) $K_X^2 = 2$ *and* $p > 3$.
(3) $K_X^2 = 3$ *and* $p > 2$.
(4) $K_X^2 = 4$.

Then X is F-split.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

Proof. In each case, X is log liftable by [KN22, Theorem 1.7 (1)], and thus the condition (1) of Proposition 3.2 is satisfied. Thus, it suffices to confirm the condition (2) of Proposition 3.2, that is, $H^1(X, \Omega_X^{[1]}(pK_X)) = 0$. By Serre duality of Cohen–Macaulay sheaves [KM98, Theorem 5.71], we have $H^1(X, \Omega_X^{[1]}(-pK_X)) \cong H^1(X, \Omega_X^{[1]}(pK_X))$. Since X has only hypersurface singularities, Ω_X^1 is torsion-free by [Lip65, Section 8 (1)], and the natural map $\Omega_X^1 \to \Omega_X^{[1]}$ is injective. Since $\mathcal{O}_X(-pK_X)$ is Cartier, we have an exact sequence

$$0 \to \Omega_X^1 \otimes \mathcal{O}_X(-pK_X) \to \Omega_X^{[1]}(-pK_X) \to \mathcal{C} \to 0$$

for some coherent sheaf $\mathcal{C}$ satisfying $\dim \mathrm{Supp}(\mathcal{C}) = 0$. Since $H^1(X, \mathcal{C}) = 0$, it suffices to show that

$$H^1(X, \Omega_X^1 \otimes \mathcal{O}_X(-pK_X)) = 0.$$

In what follows, we divide the proof into the cases according to (1)–(4) in the proposition.

The case (1): In this case, X is a hypersurface of $P := \mathbb{P}(1, 1, 2, 3)$ of degree 6 [BT22, Theorem 2.15]. By [Mor75, Theorem 1.7], the non-Gorenstein locus of P is $\{[0 : 0 : 0 : 1], [0 : 0 : 1 : 0]\}$, and this locus coincides with the singular locus of P (Remark 3.6). Thus, X is contained in the smooth locus of P since it is Gorenstein. We define invertible sheaves $\mathcal{O}_X(n)$ by $\mathcal{O}_P(n)|_X$ for all $n \in \mathbb{Z}$.

By adjunction, we have $\omega_X = \mathcal{O}_X(-1)$, and thus we aim to show that

$$H^1(X, \Omega_X^1 \otimes \mathcal{O}_X(p)) = 0.$$

By the conormal exact sequence, we have an exact sequence

$$\mathcal{O}_X(-X + p) = \mathcal{O}_X(p - 6) \to \Omega_P^1|_X \otimes \mathcal{O}_X(p) \to \Omega_X^1 \otimes \mathcal{O}_X(p) \to 0.$$

Since $\mathcal{O}_X(p-6)$ is torsion-free and the first map is injective outside the singular points of X, we obtain an exact sequence

$$0 \to \mathcal{O}_X(p - 6) \to (\Omega_P^1 \otimes \mathcal{O}_P(p))|_X \to \Omega_X^1 \otimes \mathcal{O}_X(p) \to 0.$$

Since $p \geq 7$, we have $H^2(X, \mathcal{O}_X(p - 6)) = 0$ by Serre duality, and hence it suffices to show that $H^1(X, (\Omega_P^1 \otimes \mathcal{O}_P(p))|_X) = 0$. We have an exact sequence

$$\Omega_P^{[1]}(p - 6) = \Omega_P^{[1]}(p) \otimes \mathcal{O}_P(-6) \to \Omega_P^{[1]}(p) \to \Omega_P^{[1]}(p)|_X \to 0.$$

Here, we obtain the first equality as follows:

$$\Omega_P^{[1]}(p) \otimes \mathcal{O}_P(-6) = (\Omega_P^{[1]} \otimes \mathcal{O}_P(p))^{**} \otimes \mathcal{O}_P(-6) = (\Omega_P^{[1]} \otimes \mathcal{O}_P(p - 6))^{**} = \Omega_P^{[1]}(p - 6)$$

since $\mathcal{O}_P(-6)$ is Cartier. In particular, the first term of the above exact sequence is torsion-free, and thus the first map is injective since it is injective outside the singular points of P.

Moreover, since X is contained in the smooth locus of P, it follows that $\Omega_P^{[1]}(p)|_X = (\Omega_P^1 \otimes \mathcal{O}_P(p))|_X$. Thus, we obtain an exact sequence

$$0 \to \Omega_P^{[1]}(p - 6) \to \Omega_P^{[1]}(p) \to (\Omega_P^1 \otimes \mathcal{O}_P(p))|_X \to 0.$$

By Bott vanishing on P [Fuj07, Corollary 1.3], we have

$$H^1(P, \Omega_P^{[1]}(p)) = H^2(P, \Omega_P^{[1]}(p - 6)) = 0$$

since $p \geq 7$. Therefore, we obtain $H^1(X, (\Omega_P^1 \otimes \mathcal{O}_P(p))|_X) = 0$.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

The case (2): In this case, X is a hypersurface of $P \coloneqq \mathbb{P}(1,1,1,2)$ of degree 4 [BT22, Theorem 2.15]. By [Mor75, Theorem 1.7], the non-Gorenstein locus of P is $\{[0:0:0:1]\}$, and this locus coincides with the singular locus of P (Remark 3.6). Thus, X is contained in the smooth locus of P since it is Gorenstein.

By adjunction, we have $\omega_X = \mathcal{O}_X(-1)$, and thus we aim to show that

$$H^1(X, \Omega^1_X \otimes \mathcal{O}_X(p)) = 0.$$

As in the case (1), by the conormal exact sequence and the torsion-freeness of $\mathcal{O}_X(p-4)$, we have an exact sequence

$$0 \to \mathcal{O}_X(p-4) \to \Omega^1_P|_X \otimes \mathcal{O}_X(p) \to \Omega^1_X \otimes \mathcal{O}_X(p) \to 0.$$

Since $p \geq 5$, we have $H^2(X, \mathcal{O}_X(p-4)) = 0$, and it suffices to show that

$$H^1(X, \Omega^1_P|_X \otimes \mathcal{O}_X(p)) = 0.$$

As in the case (1), we have an exact sequence

$$0 \to \Omega^{[1]}_P(p-4) \to \Omega^{[1]}_P(p) \to \Omega^1_P|_X \otimes \mathcal{O}_X(p) \to 0.$$

By Bott vanishing [Fuj07, Corollary 1.3], we have

$$H^1(P, \Omega^{[1]}_P(p)) = H^2(P, \Omega^{[1]}_P(p-4)) = 0$$

since $p \geq 5$. Therefore, we obtain $H^1(X, \Omega^1_P(p)|_X) = 0$.

The case (3): In this case, X is a hypersurface of $P \coloneqq \mathbb{P}^3$ of degree 3 [BT22, Theorem 2.15]. By adjunction, we have $\omega_X = \mathcal{O}_X(-1)$, and thus we aim to show that $H^1(X, \Omega^1_X(p)) = 0$. By the conormal exact sequence and the torsion-freeness of $\mathcal{O}_X(p-3)$, we have an exact sequence

$$0 \to \mathcal{O}_X(p-3) \to \Omega^1_P(p)|_X \to \Omega^1_X(p) \to 0.$$

Since $p \geq 3$, we have $H^2(X, \mathcal{O}_X(p-3)) = 0$, and it suffices to show that $H^1(X, \Omega^1_P(p)|_X) = 0$. We have an exact sequence

$$0 \to \Omega^1_P(p-3) \to \Omega^1_P(p) \to \Omega^1_P(p)|_X \to 0.$$

By Bott vanishing [Fuj07, Corollary 1.3], we have $H^1(P, \Omega^1_P(p)) = 0$. By [Tot24, Proposition 1.3], we also have $H^2(P, \Omega^1_P(p-3)) = 0$ since $p \geq 3$. Therefore, we obtain $H^1(X, \Omega^1_P(p)|_X) = 0$.

The case (4): In this case, X is a complete intersection of two quadric hypersurfaces Q and Q' of $P \coloneqq \mathbb{P}^4$ [BT22, Theorem 2.15]. By adjunction, we have $\omega_X = \mathcal{O}_X(-1)$, and thus we aim to show that $H^1(X, \Omega^1_X(p)) = 0$. By the conormal exact sequence and the torsion-freeness of $\mathcal{O}_X(p-2)$, we have an exact sequence

$$0 \to \mathcal{O}_X(p-2) \to (\Omega^1_Q \otimes \mathcal{O}_Q(p))|_X \to \Omega^1_X \otimes \mathcal{O}_X(p) \to 0.$$

Since $p \geq 2$, we have $H^2(X, \mathcal{O}_X(p-2)) = 0$, and hence it suffices to show that $H^1(X, (\Omega^1_Q \otimes \mathcal{O}_Q(p))|_X) = 0$.

We define invertible sheaves $\mathcal{O}_Q(n)$ as $\mathcal{O}_P(n) \otimes \mathcal{O}_Q$ for all $n \in \mathbb{Z}$. We have an exact sequence

$$\Omega^1_Q \otimes \mathcal{O}_Q(p-2) \to \Omega^1_Q \otimes \mathcal{O}_Q(p) \to (\Omega^1_Q \otimes \mathcal{O}_Q(p))|_X \to 0.$$

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

Since X is regular in codimension one, Ω^1_Q is torsion-free by [Lip65, Section 8 (1)]. Thus, we have an exact sequence

$$0 \to \Omega^1_Q \otimes \mathcal{O}_Q(p-2) \to \Omega^1_Q \otimes \mathcal{O}_Q(p) \to (\Omega^1_Q \otimes \mathcal{O}_Q(p))|_X \to 0.$$

Therefore, it suffices to show that

$$H^1(Q, \Omega^1_Q \otimes \mathcal{O}_Q(p)) = 0 \text{ and } H^2(Q, \Omega^1_Q \otimes \mathcal{O}_Q(p-2)) = 0,$$

and in particular, the following claim finishes the proof of the case (4):

Claim. We have

(i) $H^1(Q, \Omega^1_Q \otimes \mathcal{O}_Q(n)) = 0$ for every $n \in \mathbb{Z} \setminus \{0\}$ and
(ii) $H^2(Q, \Omega^1_Q \otimes \mathcal{O}_Q(n)) = 0$ for every $n \in \mathbb{Z}$.

We have

(a) $H^i(P, \Omega^1_P(n)) = 0$ for every $n \in \mathbb{Z} \setminus \{0\}$ and $i \in \{1, 2, 3\}$, and
(b) $H^j(P, \Omega^1_P(n)) = 0$ for every $n \in \mathbb{Z}$ and $j \in \{2, 3\}$.

Indeed, (a) follows from Bott vanishing and Serre duality. Then (a), together with [Tot24, Proposition 1.3], implies (b). By the following exact sequence

$$0 \to \Omega^1_P(n-2) \to \Omega^1_P(n) \to \Omega^1_P(n)|_Q \to 0,$$

we get

(i)' $H^1(Q, \Omega^1_P(n)|_Q) = 0$ for every $n \in \mathbb{Z} \setminus \{0\}$, and
(ii)' $H^2(Q, \Omega^1_P(n)|_Q) = 0$ for every $n \in \mathbb{Z}$.

By the conormal exact sequence and the torsion-freeness of $\mathcal{O}_P(n-2)$, we have an exact sequence

$$0 \to \mathcal{O}_P(n-2) \to \Omega^1_P(n)|_Q \to \Omega^1_Q \otimes \mathcal{O}_Q(n) \to 0$$

for every $n \in \mathbb{Z}$. Since $H^2(P, \mathcal{O}_P(n)) = H^3(P, \mathcal{O}_P(n)) = 0$ for every $n \in \mathbb{Z}$, we have the claim. □

Remark 3.6. Take positive integers q_1, q_2, q_3 such that $\gcd(q_1, q_2, q_3) = 1$. Set $P := \mathbb{P}(1, q_1, q_2, q_3)$. Then it is well known (cf. [Ful93, Section 2.2]) that P coincides with the projective $\mathbb{Q}$-factorial toric threefold associated to the fan in $\mathbb{R}^3$ that is generated by four rays $\mathbb{R}u, \mathbb{R}e_1, \mathbb{R}e_2, \mathbb{R}e_3$, where e_1, e_2, e_3 is the standard basis of $\mathbb{Z}^3$ and

$$u := -(q_1e_1 + q_2e_2 + q_3e_3).$$

In the above proof, we have used the results (1) and (2).

(1) $\mathbb{P}(1, 1, 2, 3)$ has exactly two singular points, which corresponds to the cones $\mathbb{R}u + \mathbb{R}e_1 + \mathbb{R}e_2$ and $\mathbb{R}u + \mathbb{R}e_1 + \mathbb{R}e_3$ [CLS11, Theorem 1.3.12].
(2) $\mathbb{P}(1, 1, 1, 2)$ has a unique singular point, which corresponds to the cone $\mathbb{R}u + \mathbb{R}e_1 + \mathbb{R}e_2$ [CLS11, Theorem 1.3.12].

From now on, we focus on the case where $K_X^2 \geq 5$.

Proposition 3.7. *The following assertions hold.*

(1) *Fix an integer m satisfying $1 \leq m \leq 5$. Let $P_1, \ldots, P_m$ be distinct points on $\mathbb{P}^2$ such that the blowup X of $\mathbb{P}^2$ along $\{P_1, \ldots, P_m\}$ is a weak del Pezzo surface. Then X is F-split.*
(2) *Fix an integer n satisfying $1 \leq n \leq 4$. Let $Q_1, \ldots, Q_n$ be distinct points on $\mathbb{P}^1 \times \mathbb{P}^1$ such that the blowup X of $\mathbb{P}^1 \times \mathbb{P}^1$ along $\{Q_1, \ldots, Q_n\}$ is a weak del Pezzo surface. Then X is F-split.*

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

Proof. Let us show (1). In what follows, we only treat the case when $m = 5$, as otherwise the problem is easier. Let L (resp. L') be the line on $\mathbb{P}^2$ passing through P_1 and P_2 (resp. P_3 and P_4). Since X is weak del Pezzo, we obtain $L \neq L'$. Pick a general line L'' on $\mathbb{P}^2$ passing through P_5. Then $L + L' + L''$ is simple normal crossing. Therefore, $(\mathbb{P}^2, L + L' + L'')$ is F-split (Example 2.7(2)), which implies that so is X (Proposition 2.4). Thus, (1) holds. The proof of (2) is similar to that of (1). Indeed, for each projection $\mathrm{pr}_i \colon \mathbb{P}^1 \times \mathbb{P}^1 \to \mathbb{P}^1$, it is enough to take two fibers F_i and F_i' such that $F_1 \cup F_1' \cup F_2 \cup F_2'$ contains $\{Q_1, \ldots, Q_n\}$ (Example 2.7(3)). □

Proposition 3.8. *Let X be a smooth weak del Pezzo surface satisfying $K_X^2 \geq 5$. Then X is F-split.*

Proof. By [Dol12, Theorem 8.1.15], we may assume that there is a birational morphism $f \colon X \to \mathbb{P}^2$. In what follows, we only treat the case when $K_X^2 = 5$, as the other cases are simpler. There are the following five cases [Dol12, Section 8.5].

(i) P, Q, R, S.
(ii) $P' \succ P, Q, R$.
(iii) $P' \succ P, Q' \succ Q$.
(iv) $P'' \succ P' \succ P, Q$.
(v) $P'' \succ P' \succ P \succ Q$.

For the definition of $P' \succ P$, we refer to [Dol12, Section 7.3.2]. For example, in the case (iii), we have $X = Y'' \to Y' \to Y = \mathbb{P}^2$, where $Y' = \mathrm{Bl}_{P \amalg Q} Y$, $Y'' = \mathrm{Bl}_{P' \amalg Q'}$, and P' and Q' are points on Y' lying over P and Q, respectively.

The case (i) has been settled in Proposition 3.7, because we have $X = \mathrm{Bl}_{P \amalg Q \amalg R \amalg S} \mathbb{P}^2$ for distinct points $P, Q, R, S \in \mathbb{P}^2$ in this case. In the case (ii), P, Q, R are distinct points on $\mathbb{P}^2$, and we have $X = \mathrm{Bl}_{P'} Y'$ for $Y' := \mathrm{Bl}_{P \amalg Q \amalg R} \mathbb{P}^2$ and a closed point P' lying over P. Take the line $L_1 := \overline{PQ}$ passing through P and Q. Let L_2 and L_3 be general lines passing through P and R, respectively. Then $(\mathbb{P}^2, L_1 + L_2 + L_3)$ is F-split (Example 2.7(2)). Since Δ is effective for the divisor Δ defined by $K_X + \Delta = f^*(K_{\mathbb{P}^2} + L_1 + L_2 + L_3)$, it follows that X is F-split (Proposition 2.4). Similarly, (iii) is settled by taking the line $L_1 := \overline{PQ}$ and general lines L_2 and L_3 passing through P and Q, respectively.

Let us treat the case (iv). In this case, we have $X = Y''' \to Y'' \to Y' \to Y = \mathbb{P}^2$, where $Y' := \mathrm{Bl}_{P \amalg Q} Y, Y'' := \mathrm{Bl}_{P'} Y', Y''' := \mathrm{Bl}_{P''} Y''$, and P' (resp. P'') is lying over P (resp. P'). Let L_1 be the line on $Y = \mathbb{P}^2$ such that $P \in L_1$ and $P' \in L_1'$ for the proper transform L_1' of L_1 on Y'. Let L_2 and L_3 be general lines passing through P and Q, respectively. Then we can check that the divisor Δ defined by $K_X + \Delta = f^*(K_{\mathbb{P}^2} + L_1 + L_2 + L_3)$ is effective. Since $(\mathbb{P}^2, L_1 + L_2 + L_3)$ is F-split (Example 2.7(2)), so is X. This completes the proof for the case (iv).

Let us consider the case (v). In this case, we apply a similar method to that of (iv) after replacing $\mathbb{P}^2$ by $\mathbb{F}_1$. We have a sequence of one-point blowups:

$$f \colon X = Y''' \to Y'' \to Y' \to Y = \mathbb{F}_1 \to Z = \mathbb{P}^2,$$

where $Y := \mathrm{Bl}_Q \mathbb{P}^2 = \mathbb{F}_1$, $Y' := \mathrm{Bl}_P Y, Y'' := \mathrm{Bl}_{P'} Y'$, $Y''' := \mathrm{Bl}_{P''} Y''$, and P, P', P'' are lying over Q, P, P', respectively. For the (-1)-curve C on Y, we have $P \in C$. It is well known that there is another section $\widetilde{C}$ of the $\mathbb{P}^1$-bundle $\pi \colon Y = \mathbb{F}_1 \to B = \mathbb{P}^1$ such that $C \cap \widetilde{C} = \emptyset$ and $\widetilde{C}^2 = 1$. Let F is a fiber of π. Since $(K_Y + C + \widetilde{C}) \cdot F = 0$, there exists $n \in \mathbb{Z}$ such that $K_Y + C + \widetilde{C} \sim nF$. Then $n = C \cdot nF = C \cdot (K_Y + C + \widetilde{C}) = -2$. Since the proper transform C' of C on Y' satisfies $C'2 = -2$, we obtain

(1) $P' \notin C'$,

as otherwise, the proper transform C'' of C' on Y'' would satisfy $C'\prime 2 = -3$, which contradicts the fact that Y'' is weak del Pezzo.

We now treat the case when $P' \in F_P'$, where F_P' denotes the proper transform of the fiber F_P of $\pi \colon Y = \mathbb{F}_1 \to \mathbb{P}^1$ passing through P. Let $\widetilde{F}$ be a general fiber of π. As we have seen above,

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

$K_Y + \widetilde{F} + C + F_P + \widetilde{C} \sim 0$. Since $\widetilde{F}$ is nef and $\mathbb{F}_1$ is toric, we obtain $H^1(\mathbb{F}_1, \mathcal{O}_{F_1}(\widetilde{F})) = 0$ by [Tot24, Proposition 1.3]. Moreover, $(\widetilde{F}, (C + F_P + \widetilde{C})|_{\widetilde{F}}) = (\widetilde{F}, C|_{\widetilde{F}} + \widetilde{C}|_{\widetilde{F}})$ is F-split (Example 2.7 (1)). Thus, $(Y, C + F_P + \widetilde{C} + \widetilde{F})$ is F-split (Proposition 2.6), which implies that X is F-split (Proposition 2.4). In what follows, we assume that

(2) $P' \notin F'_P$ for the proper transform F'_P of the fiber F_P of $\pi : Y = \mathbb{F}_1 \to \mathbb{P}^1$ passing through P.

Claim. There is a section D of $\pi\colon Y = \mathbb{F}_1 \to \mathbb{P}^1$ such that

(a) $D \sim \widetilde{C} + F$,
(b) $P \in D$, and
(c) $P' \in D'$ for the proper transform D' of D on Y'.

Proof of Claim. Since $\widetilde{C} + F$ is an ample Cartier divisor on $Y = \mathbb{F}_1$, it follows that $|\widetilde{C} + F|$ is very ample [Har77, Ch. V, Corollary 2.18]. Then there is an effective Cartier divisor D on $Y = \mathbb{F}_1$ satisfying (a)–(d).

(d) D is smooth at P.

In fact, since $|\widetilde{C} + F|$ is very ample, the elements of $H^0(Y, \mathcal{O}_Y(\widetilde{C} + F))$ separate tangent vectors. Let $s_{P'} \in \mathfrak{m}_P/\mathfrak{m}_P^2$ is an element that corresponds to P'. We take D as a divisor of zeros of a global section $s \in H^0(Y, \mathcal{O}_Y(\widetilde{C} + F))$ that maps to $s_{P'} \in \mathfrak{m}_P/\mathfrak{m}_P^2$. Then (a)–(c) are satisfied. Since $s \in \mathfrak{m}_P/\mathfrak{m}_P^2$ is non-zero, the divisor D is smooth at P, i.e., (d) is satisfied. Since $D \cdot F = (\widetilde{C} + F) \cdot F = 1$, we can write $D = D_0 + F_1 + \cdots + F_r$, where $r \geq 0$, D_0 is a section of $\pi\colon Y = \mathbb{F}_1 \to \mathbb{P}^1$, and each F_i is a fiber of π. It suffices to prove $r = 0$. Suppose $r > 0$. The following holds:

$$D_0 \cdot C + r = (D_0 + F_1 + \cdots + F_r) \cdot C = D \cdot C = (\widetilde{C} + F) \cdot C = 1. \tag{3.8.1}$$

We now treat the case when $D_0 \neq C$. In this case, $D_0 \cdot C \geq 0$ and (3.8.1) imply $r = 1$ and $D_0 \cdot C = 0$. Hence, we get $D_0 \cap C = \emptyset$. Since $P \in C$, we have $P \notin D_0$. By (b), we obtain $P \in D = D_0 + F_1$, and thus $P \in F_1$. Hence, $D = D_0 + F_P$, where F_P denotes the fiber passing through P. Since $P' \notin C'$, we obtain $P' \in D' \setminus C' \subset F'_P$. This contradicts (2).

Hence, we may assume that $D_0 = C$. We then get $D = C + F_1 + \cdots + F_r$. Since $P \in C$, we obtain $P \notin F_1 \cup \cdots \cup F_r$ by (d). Then $P' \notin F'_1 \cup \cdots \cup F'_r$, where $F'_1, F'_2, \ldots, F'_r$ are proper transforms of $F_1, F_2, \ldots, F_r$ on Y', and thus we obtain $P' \in D' \setminus \{F'_1 \cup \cdots \cup F'_r\} \subset C'$ by (c). This contradicts (1). This completes the proof of the claim. □

We have $C \cdot D = 1$. Hence, $C \cap D = P$ and $C + D$ is a simple normal crossing divisor. Since both C and D are sections of $\pi\colon Y = \mathbb{F}_1 \to \mathbb{P}^1$, it follows that $C + D + \widetilde{F}$ is still simple normal crossing for a general fiber $\widetilde{F}$ of π. Then we see that $(Y, C + D + \widetilde{F})$ is F-split (Proposition 2.6, Example 2.7(1)). Since the divisor Δ defined by $K_X + \Delta = f^*(K_Y + C + D + \widetilde{F})$ is effective, X is F-split (Proposition 2.4). This completes the proof of Proposition 3.8. □

Proof of Theorem B. If $K_X^2 \leq 4$ (resp. $K_X^2 \geq 5$), then the assertion follows from Proposition 3.5 (resp. Proposition 3.8). □

3.3. Global F-regularity: Proof of Theorem A

In this subsection, we deduce Theorem A from Theorem B.

Lemma 3.9. *Let X be a canonical del Pezzo surface. Suppose that one of the following holds.*

(1) $p > 5$
(2) $K_X^2 \geq 2$ *and* $p > 3$.
(3) $K_X^2 \geq 4$ *and* $p > 2$.
(4) $K_X^2 \geq 5$.

Then X is strongly F-regular.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

Remark 3.10. Combining [KN23, Table 1] and [KT24a, Table 1], we can see that the assumption of p is optimal for each degree.

Proof. Strongly F-regular surface singularities are completely classified by Hara [Har98b, Theorem 1.1]. In what follows, we confirm the singularities on X satisfying one of (1)–(4) are all strongly F-regular.

(1) follows from [Har98b, Theorem 1.1]. In what follows, let $f\colon Y \to X$ be the minimal resolution. Then Y is a smooth weak del Pezzo surface (Subsection 2.1.2), and Y is obtained by a blowup of $\mathbb{P}^2$ at some points [Dol12, Theorem 8.1.15]. We have holds.

We prove (2). Since $K_Y^2 = K_X^2 \geq 2$, we have $\rho(Y) = 10 - K_Y^2 \leq 8$. Thus, the number of the (-2)-curves contracted by f is at most $8 - \rho(X) \leq 7$. Therefore, X does not have canonical singularities of E_8-type. Then X is strongly F-regular by [Har98b, Theorem 1.1] since $p > 3$ (see also [KT24a, Table 1]).

Next, we prove (3). Since $K_Y^2 = K_X^2 \geq 4$, we have $\rho(Y) = 10 - K_Y^2 \leq 6$. Thus, the number of the (-2)-curves contracted by f is at most $6 - \rho(X) \leq 5$. Therefore, X does not have canonical singularities of E-type. Then X is strongly F-regular by [Har98b, Theorem 1.1] since $p > 2$ (see also [KT24a, Table 1]).

Finally, we prove (4). Since $K_Y^2 = K_X^2 \geq 5$, we have $\rho(Y) = 10 - K_Y^2 \leq 5$. Thus, the number of the (-2)-curves contracted by f is at most $5 - \rho(X) \leq 4$. If $\rho(X) \geq 2$, then X has only A-type singularities, which are strongly F-regular [Har98b, Theorem 1.1]. If $\rho(X) = 1$, then X has only A-type singularities by [KN23, Theorem 1.1]. □

Proof of Theorem A. Let X be as in the statement of Theorem A. Taking the anticanonical model of X, we may assume that $-K_X$ is ample (Corollary 2.5). Then, by Theorem 1.5, it is enough to prove that X is strongly F-regular and F-split, which follow from Lemmas 3.9 and Theorem B, respectively. □

Acknowledgements. The authors express their gratitude to Burt Totaro and the anonymous referee for valuable comments.

Funding statement. Kawakami was supported by JSPS KAKENHI Grant number JP22KJ1771 and JP24K16897. Tanaka was supported by JSPS KAKENHI Grant number JP22H01112 and JP23K03028.

Competing interest. The authors have no competing interest to declare.

References

[BK05] M. Brion and S. Kumar, *Frobenius splitting methods in geometry and representation theory*. Progress in Mathematics, vol. 231. (Birkhäuser Boston, Inc., Boston, MA, 2005).

[BT22] F. Bernasconi and H. Tanaka, 'On del Pezzo fibrations in positive characteristic', *J. Inst. Math Jussieu* **21**(1) (2022), 197–239.

[CLS11] DA. Cox, JB. Little and HK Schenck. *Toric varieties*. Graduate Studies in Mathematics, vol. 124. (American Mathematical Society, Providence, RI, 2011).

[CTW17] P. Cascini, H. Tanaka and J. Witaszek, 'On log del Pezzo surfaces in large characteristic', *Compos. Math.* **153**(4) (2017), 820–50.

[CTW18] P. Cascini, H. Tanaka H and J. Witaszek, 'Klt del Pezzo surfaces which are not globally F-split', *Int. Math Res. Not. IMRN.* (**7**) (2017), 2135–55.

[Dol12] IV. Dolgachev, *Classical algebraic geometry*. (Cambridge University Press, Cambridge, 2012).

[Fuj07] O. Fujino O, 'Multiplication maps and vanishing theorems for toric varieties', *Math Z.* **257**(3) (2017), 631–41.

[Ful93] W. Fulton, *Introduction to toric varieties*. The William H. Roever Lectures in Geometry. Annals of Mathematics Studies, vol. 131. (Princeton University Press, Princeton, NJ, 1993),

[Har77] R. Hartshorne, *Algebraic geometry*. (Graduate Texts in Mathematics, No. 52). (Heidelberg: Springer-Verlag, New York, 1977).

[Har98a] N. Hara, 'A characterization of rational singularities in terms of injectivity of Frobenius maps', *Amer. J. Math.* **120**(5) (1998), 981–96.

[Har98b] N. Hara, 'Classification of two-dimensional F-regular and F-pure singularities', *Adv. Math.* **133**(1) (1998), 33–53.

[HX15] CD. Hacon and Xu ChenyangXu, 'On the three dimensional minimal model program in positive characteristic', *J. Amer. Math. Soc.* **28**(3) (2015), 711–44.

[Kaw22a] T. Kawakami, 'Bogomolov–Sommese vanishing and liftability for surface pairs in positive characteristic', *Adv. Math.* **409** (2022), Paper No. 108640.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press

[Kaw22b] T. Kawakami, 'Extendability of differential forms via Cartier operators', *J. Eur. Math Soc (JEMS)*. Preprint, 2022, arXiv:2207.13967.

[Kaw22c] T. Kawakami, 'On the Kawamata–Viehweg vanishing theorem for log Calabi–Yau surfaces in large characteristic', *Ann. Inst. Fourier*. Preprint, 2022, arXiv:2211.08751.

[KM98] J. Kollár, S. Mori, *Birational geometry of algebraic varieties*. With the collaboration of C. H. Clemens and A. Corti, Translated from Japanese. (Cambridge Tracts in Mathematics, No. 34). (Cambridge: Cambridge University Press, 1998).

[KN22] T. Kawakami, M. Nagaoka, 'Pathologies and liftability of Du Val del Pezzo surfaces in positive characteristic', *Math Z.* (**3**) (2022), 2975–3017.

[KN23] T. Kawakami, M. Nagaoka, 'Classification of Du Val del Pezzo surfaces of Picard rank one in characteristic two and three', *J. Algebra.* **636** (2023), 603–25.

[KT23] T. Kawakami T, B. Totaro. 'Endomorphisms of varieties and Bott vanishing', *J. Algebraic Geom.* Preprint, 2023, arXiv:2302.11921v5.

[KT24a] T. Kawakami, T. Takamatsu. 'On Frobenius liftability of surface singularities', 2024, arXiv:2402.08152.

[KT24b] T. Kawakami, H. Tanaka. Vanishing theorems for Fano threefolds in positive characteristic. 2024, arXiv:2404.04764.

[KT24c] T. Kawakami, H. Tanaka. 'Weak quasi-F-splitting and del Pezzo varieties', to appear in Journal of the London Mathematical Society.

[Lip65] J. Lipman. 'Free derivation modules on algebraic varieties', *Amer. J. Math.* 1965, **87**:874–98.

[Lip69] J. Lipman. 'Rational singularities, with applications to algebraic surfaces and unique factorization', *Inst. Hautes Études Sci. Publ. Math*. (36) (1969), 195–279.

[LMM21] C. Liedtke, G. Martin, Y. Matsumoto. 'Torsors over the rational double points in characteristic' (2021), arXiv:2110.03650.

[Mor75] S. Mori. 'On a generalization of complete intersections', *J. Math. Kyoto Univ.* **15**(3) (1975), 619–46.

[Sch08] K. Schwede. 'Generalized test ideals, sharp F-purity, and sharp test elements', *Math. Res. Lett.* **15**(6) (2008), 1251–61.

[SS10] K. Schwede, KE Smith. 'Globally F-regular and log Fano varieties', *Adv. Math.* **224**(3) (2010), 863–94.

[Tan15] H. Tanaka. 'The X-method for klt surfaces in positive characteristic', *J. Algebraic Geom.* **24**(4) (2015), 605–28.

[Tot24] B. Totaro. 'Bott vanishing for Fano threefolds', *Math. Z.* **307**(1) (2024), Paper No. 14, 31.

https://doi.org/10.1017/fms.2024.143 Published online by Cambridge University Press