Math. Struct. in Comp. Science (2005), *vol.* 15, *pp.* 149–166. © 2005 Cambridge University Press
DOI: 10.1017/S0960129504004402 Printed in the United Kingdom

Operations and equations for coalgebras

ALEXANDER KURZ[†] and JIŘÍ ROSICKÝ[‡]

[†]*Department of Computer Science, University of Leicester, UK.*
[‡]*Masaryk University, Janáčkovo nám. 2a, 662 95 Brno, Czech Republic*

Received 14 November 2002; revised 12 February 2004

We show how coalgebras can be presented by operations and equations. This is a special case of Linton's approach to algebras over a general base category $\mathscr{X}$, namely where $\mathscr{X}$ is taken as the dual of sets. Since the resulting equations generalise coalgebraic coequations to situations without cofree coalgebras, we call them coequations. We prove a general co-Birkhoff theorem describing covarieties of coalgebras by means of coequations. We argue that the resulting coequational logic generalises modal logic. This relies on the fact that coalgebraic operations respect an appropriate notion of bisimulation and can be considered as modal operators.

1. Introduction

Let us start by recalling that universal algebras are defined as sets equipped with operations subjected to equations. Operations can be infinitary. Given a set X, a mapping $f : A^X \to A$ is an X-ary operation on a set A. One is often working with Y-tuples $f_y : A^X \to A$, $y \in Y$, of X-ary operations on a set A. These Y-tuples correspond uniquely to mappings $f : A^X \to A^Y$. Starting with a set of operations, one always has free algebras. But there are important examples of universal algebras given by a class of operations that still have free algebras (complete semilattices, compact Hausdorff spaces). Linton showed in Linton (1966) that equationally defined universal algebras are, under the existence of free algebras, precisely the monadic categories over Set. Moreover, in Linton (1969), he generalised the result from sets to any base category $\mathscr{X}$. In that work, operations are still mappings

$$f : A^X \to A^Y$$

where A, X, and Y are objects in $\mathscr{X}$ and A^X is the set of all morphisms $X \to A$. Note that, in Set, A^X coincides with the X-fold product of A. In general, however, it is important to consider A^X as a set of morphisms because the other approach would be too special for a general base category $\mathscr{X}$. In particular, it would be too special for $\mathscr{X} = \mathsf{Set}^{\mathrm{op}}$. Davis (1972) used Linton's approach for introducing universal coalgebras over Set even without assuming the existence of cofree coalgebras (that is, free algebras over $\mathsf{Set}^{\mathrm{op}}$). The

[†] Partially supported by the Ministry of Education of the Czech Republic under the project MSM 143100009.
[‡] Supported by the Grant Agency of the Czech Republic under the grant 201/02/0148.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

second author then considered Linton's algebras over a general base category, without the existence of free algebras, in Rosický (1981).

There is another way of defining universal algebras over a general base category. One starts with an endofunctor $F : \mathscr{X} \to \mathscr{X}$ and defines F-algebras as objects A equipped with a morphism $\alpha : FA \to A$. These algebras are called F-dynamics in Manes (1976) and were extensively studied by Trnková and her students in Prague, *cf.* Adámek and Trnková (1990). Notably, Reiterman (1983) compared F-algebras with algebras given by operations and equations.

There is a revived interest in universal coalgebra motivated by its connections with the theory of systems, see Rutten (2000). Coalgebras are here understood as F-algebras over $\mathscr{X} = \mathsf{Set}^{\mathrm{op}}$, that is, as sets A equipped with a mapping $\alpha : A \to FA$ where $F : \mathsf{Set} \to \mathsf{Set}$. Our aim is to show the potential of defining coalgebras by means of operations and equations.

Outline of the Paper

We investigate *coequational categories*, that is, categories of coalgebras that are presented by operations and equations. It turns out that coequational categories subsume categories of coalgebras for a functor or for a comonad (Section 3). To understand the nature of coalgebraic operations, we introduce a novel notion of behavioural equivalence (generalising Aczel–Mendler bisimulation) and establish that coalgebraic operations are predicate transformers that respect behavioural equivalence (Section 4). We then introduce equations in implicit operations for coalgebras. Since they generalise previous concepts of coalgebraic coequations to situations without cofree coalgebras, we call them coequations. We prove a general co-Birkhoff theorem showing that covarieties of coalgebras are always definable by coequations (Section 5). Since our coequations on the one hand dualise equations for algebras and on the other hand can be understood as modal formulae, a new account of the duality of modal logic and equational logic arises (Section 6). The final section provides a full explanation of the theorem of Davis (1972) saying that any conceivable category of coalgebras can be given by operations and coequations provided that we allow the arities of the operations to be proper classes.

2. Preliminaries

We work in Gödel–Bernays set theory with the axiom of choice for classes. This means that all proper classes are isomorphic, and, in particular, to the class *Ord* of all ordinals. Categories are assumed to be locally small, which means that they have a class of objects and sets of morphisms between any two given objects. Occasionally we encounter categories that do not satisfy this requirement and we call them *illegitimate*.

Given an endofunctor $F : \mathscr{X} \to \mathscr{X}$, an *$F$-coalgebra* $\mathsf{A} = (A, \alpha)$ consists of an object $A \in \mathscr{X}$ and an arrow $\alpha : A \to FA$. F-coalgebras form a category $\mathsf{FCoalg}(F)$ where a *F-morphism* $f : (A, \alpha) \to (B, \beta)$ is an arrow $f : A \to B \in \mathscr{X}$ such that $Ff \circ \alpha = \beta \circ f$. The *forgetful functor* $U : \mathsf{FCoalg}(F) \to \mathscr{X}$ maps a coalgebra (A, α) to A and a morphism $f : (A, \alpha) \to (B, \beta)$ to the arrow $f : A \to B$ in $\mathscr{X}$. U creates, and hence preserves, colimits.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

A *comonad* (M, ε, δ) is given by a functor $M : \mathscr{X} \to \mathscr{X}$ and two natural transformations $\varepsilon : M \to Id$, $\delta : M \to MM$ satisfying $M\varepsilon_X \circ \delta_X = \varepsilon_{MX} \circ \delta_X = \mathrm{id}_{MX}$ and $M\delta_X \circ \delta_X = \delta_{MX} \circ \delta_X$. The category $\mathsf{CCoalg}(M)$ of *coalgebras for the comonad* M is the full subcategory of $\mathsf{FCoalg}(M)$ whose coalgebras $\xi : X \to MX$ satisfy $M\xi \circ \xi = \delta_X \circ \xi$ and $\varepsilon_X \circ \xi = \mathrm{id}_X$.

Set denotes the category of sets and functions and $\mathscr{P} : \mathsf{Set} \to \mathsf{Set}$ the covariant powerset functor. We also use the convention $2 = \{0, 1\}$ and call the elements of 2 truth values.

Given $A, X \in \mathscr{X}$, the set of arrows $A \to X$ is denoted by X^A or sometimes $\mathscr{X}(A, X)$. For $f : X \to Y$, the function $f^A : X^A \to Y^A$ is defined as $f^A(g : A \to X) = f \circ g$. For $f : A \to B$, the function $X^f : X^B \to X^A$ is defined as $X^f(g : B \to X) = g \circ f$. For instance, with $\mathscr{X} = \mathsf{Set}$, 2^A is the set of subsets of A and $2^f : 2^B \to 2^A$ is the inverse-image map of $f : A \to B$. The assignment $- \mapsto X^-$ gives rise to a functor $X^- : \mathscr{X}^{\mathrm{op}} \to \mathsf{Set}$. We write X^U for the functor $\mathscr{A}^{\mathrm{op}} \to \mathsf{Set}$ obtained by composing $U : \mathscr{A} \to \mathscr{X}$ and $X^- : \mathscr{X}^{\mathrm{op}} \to \mathsf{Set}$.

A *concrete category* $(\mathscr{A}, U)$ (over $\mathscr{X}$) is a faithful functor $U : \mathscr{A} \to \mathscr{X}$. A functor $F : \mathscr{A} \to \mathscr{A}'$ is a concrete functor between the concrete categories $U : \mathscr{A} \to \mathscr{X}$ and $U : \mathscr{A}' \to \mathscr{X}$ if $U'F = U$. Concrete categories are concretely isomorphic if the isomorphisms are concrete functors.

3. Coalgebras for a signature

Recall from the introduction that algebras over a category $\mathscr{X}$ are given by a carrier $A \in \mathscr{X}$ and operations $A^X \to A^Y$ where $A^X = \mathscr{X}(X, A)$ denotes the set of arrows $A \to X$. Since universal coalgebra over $\mathscr{X}$ is universal algebra over $\mathscr{X}^{\mathrm{op}}$ and $\mathscr{X}^{\mathrm{op}}(X, A) = \mathscr{X}(A, X)$, coalgebras over a category $\mathscr{X}$ can be introduced as carriers $A \in \mathscr{X}$ equipped with operations

$$f : X^A \to Y^A.$$

Definition 3.1 (Coalgebras for a signature). A signature Σ over a category $\mathscr{X}$ is a class of operation symbols σ each equipped with a pair (X, Y) of objects of $\mathscr{X}$. We call σ an (X, Y)-ary operation symbol. A Σ-coalgebra A is an object $A \in \mathscr{X}$ together with maps

$$\sigma_{\mathsf{A}} : X^A \to Y^A$$

for each (X, Y)-ary operation symbol $\sigma \in \Sigma$. A morphism of Σ-coalgebras is defined as an arrow $h : A \to B$ such that the square

$$\begin{array}{ccc} X^A & \xrightarrow{\sigma_{\mathsf{A}}} & Y^A \\ {\scriptstyle X^h}\uparrow & & \uparrow{\scriptstyle Y^h} \\ X^B & \xrightarrow{\sigma_{\mathsf{B}}} & Y^B \end{array}$$

commutes for all σ in Σ. The resulting (illegitimate) category of coalgebras is denoted by $\mathsf{SCoalg}(\Sigma)$.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

Example 3.2.

1 If $\mathscr{X} = \mathsf{Set}$ and Σ consists of a single $(1,2)$-ary operation symbol, then $\mathsf{SCoalg}(\Sigma)$ has as objects pairs $(A, A' \subseteq A)$ and morphisms $h : (A, A') \to (B, B')$ are functions $h : A \to B$ such that $h^{-1}(B') = A'$.
2 Let $\mathscr{X} = \mathsf{Set}$ and consider the signature Σ consisting of a single $(2,2)$-ary operation symbol $\Box$. Σ-coalgebras are predicate transformers $\Box_{\mathsf{A}} : 2^A \to 2^A$. We have $f : A \to B$ is a coalgebra morphism iff $a \in \Box_{\mathsf{A}}(f^{-1}(V)) \Leftrightarrow f(a) \in \Box_{\mathsf{B}}(V)$. We can think of Σ-coalgebras as transition systems via the concrete isomorphism $\mathsf{SCoalg}(\Sigma) \cong \mathsf{FCoalg}(2^{(2^-)})$ that associates to each $\Box_{\mathsf{A}} : 2^A \to 2^A$ the hypersystem (*cf.* Rutten (2000)) $\boxdot : A \to 2^{2^A}$ given by $U \in \boxdot_{\mathsf{A}}(a) \Leftrightarrow a \in \Box_{\mathsf{A}}(U)$. The condition on f to be a morphism can then be written more suggestively as $a \longrightarrow_{\mathsf{A}} f^{-1}(V) \Leftrightarrow f(a) \longrightarrow_{\mathsf{B}} V$.
3 Let $\mathscr{X}$ be a category of topological spaces containing the topological space $\mathfrak{2} = (2, \{\emptyset, \{1\}, \{0,1\}\})$. The elements of $\mathscr{X}(A, \mathfrak{2})$ are the open sets of A and a $(\mathfrak{2}, \mathfrak{2})$-ary operation $\Box$ is a topological predicate transformer, *cf.* Smyth (1983). Similarly, if 2 is the two-element discrete space, then the elements of $\mathscr{X}(A, 2)$ are the clopen (that is, closed and open) subsets of A.
4 If $\mathscr{X} = \mathsf{Set}^{\mathrm{op}}$, then $\mathsf{SCoalg}(\Sigma)$ is dually isomorphic to the category of algebras over Set for the signature Σ.

Proposition 3.3. Let Σ be a signature over Set consisting of a set of operation symbols. Then there exists a functor $F : \mathsf{Set} \to \mathsf{Set}$ such that the categories $\mathsf{SCoalg}(\Sigma)$ and $\mathsf{FCoalg}(F)$ are concretely isomorphic.

Proof. Let σ be an (X, Y)-ary operation symbol from Σ and A be a Σ coalgebra. Then the mapping

$$\sigma_{\mathsf{A}} : X^A \to Y^A$$

determines the mapping

$$\hat{\sigma}_{\mathsf{A}} : A \to Y^{X^A}.$$

Let F be the product of functors Y^{X^-} over all $\sigma \in \Sigma$. Then A determines an F-coalgebra (A, α) where $\alpha : A \to FA$ is induced by $\hat{\sigma}_{\mathsf{A}}$, $\sigma \in \Sigma$. Any F-coalgebra is given in this way. Let $h : A \to B$ be a morphism of Σ-coalgebras. Since $Y^h \circ \sigma_{\mathsf{B}}$ determines the composition

$$A \xrightarrow{h} B \xrightarrow{\hat{\sigma}_{\mathsf{B}}} Y^{X^B}$$

and $\sigma_{\mathsf{A}} \circ X^h$ determines

$$A \xrightarrow{\hat{\sigma}_{\mathsf{A}}} Y^{X^A} \xrightarrow{Y^{X^h}} Y^{X^B}.$$

h is a morphism of F-coalgebras. Again, any F-coalgebra morphism is given in this way. Hence $\mathsf{SCoalg}(\Sigma) \cong \mathsf{FCoalg}(F)$. □

As usual, a signature Σ gives rise to **terms**. Terms are also equipped with arities and are defined as follows:

1 every (X, Y)-ary operation symbol is an (X, Y)-ary term,

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

2 every mapping $f : X \to Y$ determines an (X, Y)-ary term x_f,
3 having an (X, Y)-ary term t_1 and a (Y, Z)-ary term t_2, we get an (X, Z)-ary term $t_2 \cdot t_1$.

For each (X, Y)-ary term t and a Σ-coalgebra A we get the mapping

$$t_{\mathsf{A}} : X^A \to Y^A$$

as follows:

2 $(x_f)_{\mathsf{A}}(v) = f \circ v$ for $v : A \to X$,
3 $(t_2 \cdot t_1)_{\mathsf{A}} = (t_2)_{\mathsf{A}} \circ (t_1)_{\mathsf{A}}$.

An equation is a pair (t_1, t_2) of (X, Y)-ary terms. We write $t_1 = t_2$. A Σ-coalgebra A satisfies this equation iff $(t_1)_{\mathsf{A}} = (t_2)_{\mathsf{A}}$. To emphasise that these equations are evaluated in coalgebras, we call them **coequations**. A **coequational theory** E is a class of coequations. The category of all Σ-coalgebras satisfying all coequations from E is denoted by $\mathsf{ECoalg}(E)$. This may be an illegitimate category. We are interested in legitimate categories $\mathsf{ECoalg}(E)$. Each such category is equipped with a forgetful functor $U : \mathsf{ECoalg}(E) \to \mathscr{X}$ and thus it is a concrete category.

Definition 3.4 (Coequational category). A concrete category will be called coequational if it is concretely isomorphic to $\mathsf{ECoalg}(E)$ for some coequational theory E.

When $\mathscr{X}$ has products, we can reduce the number of operations and equations needed for a coequational presentation by adding the possibility of forming new terms by pairing:

4 If I is a set and t_i are (X, Y_i)-ary terms, $i \in I$, then

$$\langle t_i \rangle$$

is an $(X, \prod_I Y_i)$-ary term that is interpreted by a Σ-coalgebra A as

$$\langle t_i \rangle_{\mathsf{A}} = X^A \xrightarrow{\langle (t_i)_{\mathsf{A}} \rangle} \prod_I \left(Y_i^A \right) \cong \left(\prod_I Y_i \right)^A .$$

This will be convenient in the following examples. Note that we can avoid pairing by adding instead new operation symbols $\langle t_i \rangle$ to the signature as well as, for all $j \in I$, new equations $x_{p_j} \cdot \langle t_i \rangle = t_j$ where $p_j : \prod_I Y_i \to Y_j$ is the j-th projection.

Example 3.5.

1 Let $\mathscr{X} = \mathsf{Set}$ and consider the signature Σ consisting of a single $(2, 2)$-ary operation symbol $\Box$. Using clause (2), we have $(2 \times 2, 2)$-ary terms p_1, p_2, $\wedge$, $\to$ induced by the maps $2 \times 2 \to 2$ known as first and second projection, conjunction and implication; similarly, for each set I, a $(\prod_I 2, 2)$-ary term $\bigwedge_I$; and denote $\bigwedge_\varnothing$ by *true*. Using clause (4), we obtain terms $(p_1 \to p_2) \to (\Box p_1 \to \Box p_2)$, $\Box(p_1 \wedge p_2) \leftrightarrow (\Box p_1 \wedge \Box p_2)$, $\bigwedge_I \Box p_i \leftrightarrow \Box \bigwedge_I p_i$. In the following examples, equations $t = \mathit{true}$ are abbreviated to t.
   (a) $\mathsf{ECoalg}(\{(p_1 \to p_2) \to (\Box p_1 \to \Box p_2)\})$ is concretely isomorphic to the full subcategory of those (2^{2^-})-coalgebras $g : A \to 2^{2^A}$ for which $g(a)$ is closed under supersets for all $a \in A$. These coalgebras are known in modal logic as monotone neighbourhood frames.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

(b) $\mathsf{ECoalg}(\{\Box \mathit{true}, \Box(p_1 \wedge p_2) \leftrightarrow (\Box p_1 \wedge \Box p_2)\})$ is concretely isomorphic to the full subcategory of those (2^{2^-})-coalgebras $g : A \to 2^{2^A}$ for which $g(a)$ is closed under supersets and finite intersections for all $a \in A$. These coalgebras are also known as coalgebras for the filter functor (Gumm 2001b) or as normal neighbourhood frames.

(c) The equations $E = \{\Box \mathit{true}, \Box(p_1 \wedge p_2) \leftrightarrow (\Box p_1 \wedge \Box p_2), \Box p_1 \to p_1, \Box p_1 \to \Box\Box p_1\}$ are the usual axioms of an interior operator. $\mathsf{ECoalg}(E)$ is concretely isomorphic to the category of topological spaces with open and continuous maps as morphisms.

(d) $\mathsf{ECoalg}(\{\bigwedge_I \Box p_i \leftrightarrow \Box \bigwedge_I p_i \mid I \text{ a set }\})$ is concretely isomorphic to the category of Kripke frames, that is, $\mathsf{FCoalg}(\mathscr{P})$.

2 If $\mathscr{X}$ is the category of Stone spaces (Johnstone 1982), 2 the (discrete) two-element Stone space, and Σ consists of a single $(2,2)$-ary operation symbol $\Box$, then $\mathsf{ECoalg}(\{\Box \mathit{true}, \Box(p_1 \wedge p_2) \leftrightarrow (\Box p_1 \wedge \Box p_2)\})$ is isomorphic to the category of descriptive general frames (Goldblatt 1976).

We will show that $\mathsf{FCoalg}(F)$ is always coequational. This result is due to Reiterman. His paper, Reiterman (1983, page 62), formulates it, without proof, over $\mathsf{Set}^{\mathrm{op}}$ only (that is, for algebras over Set) and thus we present Reiterman's proof as sent to the second author in the late 1970s.

Proposition 3.6 (Reiterman). Let $F : \mathscr{X} \to \mathscr{X}$ be a functor. Then $\mathsf{FCoalg}(F)$ is coequational.

Proof. Let Σ be a signature consisting of (X, FX)-ary operation symbols σ^X for every set X and let E consist of coequations

$$\sigma^Y \cdot x_f = x_{Ff} \cdot \sigma^X \tag{1}$$

for every mapping $f : X \to Y$. There is a concrete functor $G : \mathsf{FCoalg}(F) \to \mathsf{ECoalg}(E)$ given as follows: for $\mathsf{A} = (A, \alpha)$, $G(\mathsf{A})$ is the E-coalgebra $(A, \sigma^X_{G\mathsf{A}})$ where

$$\sigma^X_{G\mathsf{A}}(v) = Fv \circ \alpha$$

for $v : A \to X$. On the other hand, there is a concrete functor $H : \mathsf{ECoalg}(E) \to \mathsf{FCoalg}(F)$ sending an E-coalgebra $\mathsf{A} = (A, \sigma^X_{\mathsf{A}})$ to the F-coalgebra $(A, \sigma^A_{\mathsf{A}}(\mathrm{id}_A))$.

We have $HG = Id$ because, for each F-coalgebra $\mathsf{A} = (A, \alpha)$ we have $\sigma^A_{G\mathsf{A}}(\mathrm{id}_A) = F\mathrm{id}_A \circ \alpha = \alpha$. Conversely, $GH = Id$ because, for each E-coalgebra $\mathsf{A} = (A, \sigma^X_{\mathsf{A}})$, we have $GH\mathsf{A} = G(A, \sigma^A_{\mathsf{A}}(\mathrm{id}_A)) = (A, F(-) \circ \sigma^A_{\mathsf{A}}(\mathrm{id}_A))$, which equals $(A, \sigma^X_{\mathsf{A}})$ because for all $v : A \to X$

$$\begin{array}{ccc} A^A & \xrightarrow{\sigma^A_{\mathsf{A}}} & (FA)^A \\ {\scriptstyle (x_v)_{\mathsf{A}}}\downarrow & & \downarrow{\scriptstyle (x_{Fv})_{\mathsf{A}}} \\ X^A & \xrightarrow{\sigma^X_{\mathsf{A}}} & (FX)^A \end{array}$$

commutes (due to the coequation $\sigma^X \cdot x_v = x_{Fv} \cdot \sigma^A$), and thus $Fv \circ \sigma^A_{\mathsf{A}}(\mathrm{id}_A) = \sigma^X_{\mathsf{A}}(v \circ \mathrm{id}_A) = \sigma^X_{\mathsf{A}}(v)$. □

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

The procedure just described needs a proper class of operation symbols. For $\mathscr{X} = \mathsf{Set}$, this can be avoided for functors preserving cofiltered limits if we exclude the existence of arbitrarily large measurable cardinals (which is consistent with ZFC). Recall that a *measurable cardinal* κ is a cardinal on which a non-principal κ-complete ultrafilter exists. An ultrafilter is κ-complete if it is closed under intersections of cardinality $< \kappa$, and it is non-principal if it does not contain a singleton-set. Let (M) denote the following set-theoretic statement, (see Adámek and Rosický (1994) for more information).

There do not exist arbitrarily large measurable cardinals. (M)

Proposition 3.7. Assume (M) and let $F : \mathsf{Set} \to \mathsf{Set}$ be a functor preserving cofiltered limits. Then $\mathsf{FCoalg}(F)$ is coequational in a signature consisting of a single operation symbol.

Proof. Let P be an infinite set whose cardinality is greater than any measurable cardinal. Then the full subcategory P of Set having a single object P is codense in Set (see Adámek and Rosický (1994, A.5). Let Σ consist of a single (P, FP)-ary operation symbol σ and E consist of coequations (1) for $f : P \to P$ (where $\sigma = \sigma^P$). Analogously to Proposition 3.6, we get a functor $G : \mathsf{FCoalg}(F) \to \mathsf{ECoalg}(E)$ and our task is to define $H : \mathsf{ECoalg}(E) \to \mathsf{FCoalg}(F)$. Let $(A, \sigma_{\mathsf{A}}) \in \mathsf{ECoalg}(E)$. Since P is infinite, the comma-category $(A{\downarrow}\mathsf{P})$ is cofiltered ($\langle u, v\rangle$ serves as a lower bound for $u, v : A \to P$). The codensity of P means that $f : A \to P$ forms a limit cone to the projection $Q : (A{\downarrow}\mathsf{P}) \to \mathsf{Set}$. Since F preserves cofiltered limits, $Ff : FA \to FP$ forms a limit cone to FQ. Coequations in E say that $\sigma_{\mathsf{A}}(f) : A \to FP$ is a cone to FQ. Thus there is a unique mapping $\alpha : A \to FA$ such that $Ff \circ \alpha = \sigma_{\mathsf{A}}(f)$ for each $f : A \to P$. We put $H\mathsf{A} = (A, \alpha)$. The rest of the proof is analogous to the proof of Proposition 3.6. □

Remark 3.8.

1 We can assume that there are no measurable cardinals. If V is a model of ZFC in which measurable cardinals exist, let κ be the smallest such. Then V_κ, the restriction of V to sets of rank $< \kappa$, is a model of ZFC that contains no measurable cardinals.
2 If there are no measurable cardinals, we can take a countable set for P.
3 Since cofiltered limits are connected, Proposition 3.7 applies to polynomial functors $F : \mathsf{Set} \to \mathsf{Set}$.

We have seen that categories of coalgebras for a functor are coequational. But one is often more interested in covarieties of these categories. Following Manes (1976), a full subcategory $\mathscr{C}$ of a coequational category $(\mathscr{A}, U)$ is called a **covariety** if $\mathscr{C}$ is closed under (existing) colimits, quotients and U-split subobjects (this is a subobject $m : \mathsf{A} \to \mathsf{B}$ such that Um is a split monomorphism). Over Set covarieties correspond to the usual notion of full subcategories closed under coproducts, quotients and subobjects.

Most of the covarieties of coequational categories have cofree coalgebras. The forgetful functor is comonadic in these cases and therefore, following Linton (1969), coequational. We show that this easily follows from Reiterman's proposition. Following

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

Mac Lane (1971), we say that a concrete category $(\mathscr{C}, U)$ is *comonadic* if U has a right adjoint R and the canonical functor from $\mathscr{C}$ to the category of all coalgebras for the comonad UR is an isomorphism.

Proposition 3.9 (Linton). Every comonadic category is coequational.

Proof. Let $F : \mathscr{X} \to \mathscr{X}$ be a comonad with counit $\varepsilon : F \to Id$ and comultiplication $\delta : F \to FF$. Then coalgebras for the comonad F are specified in $\mathsf{FCoalg}(F)$ by

$$x_{\varepsilon_X} \cdot \sigma^X = x_{id_X},$$

$$\sigma^{FX} \cdot \sigma^X = x_{\delta_X} \cdot \sigma_X.$$

□

Moreover, comonadic categories coincide with coequational categories having cofree coalgebras. Let $U : \mathscr{A} \to \mathscr{X}$ be a functor. A *U-split equaliser* is an equaliser in $\mathscr{A}$

$$\mathsf{A} \xrightarrow{\ e\ } \mathsf{B} \underset{g}{\overset{f}{\rightrightarrows}} \mathsf{C}$$

such that its U-image splits, that is, it is equipped with $t : U\mathsf{C} \to U\mathsf{B}$ and $s : U\mathsf{B} \to U\mathsf{A}$ such that $s \circ Ue = \mathrm{id}_{U\mathsf{A}}$, $t \circ Uf = \mathrm{id}_{U\mathsf{B}}$, and $t \circ Ug = Ue \circ s$. U *creates U-split equalisers* if it creates equalisers of pairs f, g for which Uf, Ug has a split equaliser in $\mathscr{X}$. Beck's theorem (Mac Lane 1971) says that U is comonadic iff it has a right-adjoint and creates U-split equalisers. We will say that a concrete category $\mathscr{A}$ is **co-Beck** if U creates colimits and U-split equalisers.

Proposition 3.10. Each covariety of a coequational category is co-Beck.

Proof. The proof is straightforward, *cf.* Rosický (1981). □

Corollary 3.11 (Linton). Every coequational category with cofree coalgebras is comonadic.

Proof. The proof follows from Proposition 3.10 and Beck's theorem. □

We have seen that every covariety (of a coequational category) with cofree coalgebras is coequational. Without the assumption, this is an open problem.

Problem 3.12. Is every covariety in $\mathsf{FCoalg}(F)$, where $F : \mathsf{Set} \to \mathsf{Set}$, coequational? More generally, is every covariety of a coequational category coequational?

Although important, the problem is not crucial for our approach. We will see in Section 5 that every covariety (of a coequational category over Set) can be defined by coequations in implicit operations.

4. Behavioural equivalence

In the case of algebras over Set, we think of operations as functions allowing us to construct new elements of the carrier. We show that in the case of coalgebras over Set, operations can be thought of as predicate transformers that respect behavioural equivalence. They will appear as modal operators in Section 6.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

We first introduce a novel notion of behavioural equivalence. It formalises the idea that behavioural equivalence is the smallest equivalence relation that is invariant under morphisms. We define two notions, the second one taking 'colourings' into account: given a functor $U : \mathscr{A} \to \mathsf{Set}$ and a set of 'colours' X, a *colouring* v for an object $\mathsf{A} \in \mathscr{A}$ is a map $U\mathsf{A} \to X$.

Definition 4.1 (Behavioural equivalence). Consider a functor $U : \mathscr{A} \to \mathsf{Set}$.

1 $\sim$ is the smallest equivalence relation on the class of all pairs (A, a), $\mathsf{A} \in \mathscr{A}$, $a \in U\mathsf{A}$, satisfying

$$(\mathsf{A}_1, a_1) \sim (\mathsf{A}_2, Uf(a_1)) \quad \text{for all } f : \mathsf{A}_1 \to \mathsf{A}_2.$$

2 Given $X \in \mathsf{Set}$, we define $\sim_X$ to be the smallest equivalence relation on the class of all triples (A, v, a), $\mathsf{A} \in \mathscr{A}$, $v : U\mathsf{A} \to X$, $a \in U\mathsf{A}$, satisfying

$$(\mathsf{A}_1, v_2 \circ Uf, a_1) \sim_X (\mathsf{A}_2, v_2, Uf(a_1)) \quad \text{for all } v_2 : U\mathsf{A}_2 \to X,\ f : \mathsf{A}_1 \to \mathsf{A}_2.$$

If $(\mathsf{A}, a) \sim (\mathsf{B}, b)$, we say that a and b are behaviourally equivalent. If $(\mathsf{A}, v, a) \sim_X (\mathsf{B}, w, b)$, we say that a and b are X-behaviourally equivalent.

Categorically speaking, the equivalence classes of behavioural equivalence are the components of the category of elements of the forgetful functor; the equivalence classes of X-behavioural equivalence are the components of the category of elements of the functor $(U{\downarrow}X) \to \mathsf{Set}$, $(\mathsf{A}, U\mathsf{A} \to X) \mapsto U\mathsf{A}$.

Example 4.2. If $\mathscr{A} = \mathsf{FCoalg}(F)$, $F : \mathsf{Set} \to \mathsf{Set}$, then $(\mathsf{A}, a) \sim (\mathsf{B}, b)$ iff a and b are identified by the morphisms into the final coalgebra. As is well known, this means that if F preserves weak pullbacks, then $(\mathsf{A}, a) \sim (\mathsf{B}, b)$ iff a and b are related by a bisimulation in the sense of Aczel and Mendler (1989). In particular, in the case of $\mathsf{FCoalg}(\mathscr{P})$ or $\mathsf{FCoalg}(\mathscr{P}(- \times L))$, behavioural equivalence yields the familiar notion of bisimilarity for (labelled) transition systems.

The following gives an alternative characterisation of behavioural equivalence (the requirement on pushouts is satisfied by coequational categories).

Proposition 4.3. Suppose that $\mathscr{A}$ has pushouts and that $U : \mathscr{A} \to \mathsf{Set}$ preserves them. $(\mathsf{A}_1, v_1, a_1) \sim_X (\mathsf{A}_2, v_2, a_2)$ iff there are B, w and morphisms $f_i : \mathsf{A}_i \to \mathsf{B}$ such that

$$\begin{array}{ccccc} & & X & & \\ & {\scriptstyle v_1}\nearrow & \uparrow{\scriptstyle w} & \nwarrow{\scriptstyle v_2} & \\ U\mathsf{A}_1 & \xrightarrow[Uf_1]{} & U\mathsf{B} & \xleftarrow[Uf_2]{} & U\mathsf{A}_2 \end{array} \tag{2}$$

commutes and $(Uf_1)(a_1) = (Uf_2)(a_2)$.

Proof. Let $\approx$ denote the relation defined by condition (2). $\approx \subseteq \sim$ is immediate. For the converse, note that $\approx$ contains the generating pairs of $\sim$ and is reflexive and symmetric. $\approx$ is transitive, since $\mathscr{A}$ has pushouts and U preserves them. □

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

An (X, Y)-valued predicate transformer P_{A} maps X-valued predicates to Y-valued predicates, that is, $P_{\mathsf{A}} : X^{U\mathsf{A}} \to Y^{U\mathsf{A}}$. It is called behavioural if it respects behavioural equivalence.

Definition 4.4 (Behavioural predicate transformers). Consider a functor $U : \mathscr{A} \to \mathsf{Set}$. An (X, Y)-valued predicate transformer P is an operation that determines for each $\mathsf{A} \in \mathscr{A}$, $v : U\mathsf{A} \to X$, $a \in U\mathsf{A}$ a value

$$P_{\mathsf{A}}(v, a) \in Y. \tag{3}$$

P is called a *behavioural* predicate transformer iff

$$(\mathsf{A}, v, a) \sim_X (\mathsf{B}, w, b) \;\Rightarrow\; P_{\mathsf{A}}(v, a) = P_{\mathsf{B}}(w, b) \tag{4}$$

for all $w : U\mathsf{B} \to X$ and $a \in U\mathsf{A}$. When $X = 1$ and $Y = 2$, we call P a *behavioural predicate*.

Note that behavioural predicates are precisely those predicates that are invariant under behavioural equivalence. The following is immediate from the respective definitions.

Lemma 4.5. An operation P that determines for each $\mathsf{A} \in \mathscr{A}$, $v : U\mathsf{A} \to X$, $a \in U\mathsf{A}$ a value $P_{\mathsf{A}}(v, a) \in Y$ is a behavioural predicate transformer iff

$$P_{\mathsf{A}}(w \circ Uf, a) = P_{\mathsf{B}}(w, Uf(a)) \tag{5}$$

for all morphisms $f : \mathsf{A} \to \mathsf{B}$, all $w : U\mathsf{B} \to X$, and all elements a of A.

We now show that invariance of a predicate transformer $X^U \to Y^U$ under X-behavioural equivalence is equivalent to the naturality of $X^U \to Y^U$.

Theorem 4.6. Consider a functor $U : \mathscr{A} \to \mathsf{Set}$. An operation P that determines for each $\mathsf{A} \in \mathscr{A}$, $v : U\mathsf{A} \to X$, $a \in U\mathsf{A}$ a truth value $P_{\mathsf{A}}(v, a) \in Y$ is a behavioural predicate transformer iff

$$P_{\mathsf{A}} : X^{U\mathsf{A}} \longrightarrow Y^{U\mathsf{A}}$$

is a natural transformation.

Proof. Naturality of P means that for any morphism $f : \mathsf{A} \to \mathsf{B}$

$$\begin{array}{ccc} X^{U\mathsf{A}} & \xrightarrow{P_{\mathsf{A}}} & 2^{U\mathsf{A}} \\ {\scriptstyle X^{Uf}}\uparrow & & \uparrow{\scriptstyle 2^{Uf}} \\ X^{U\mathsf{B}} & \xrightarrow{P_{\mathsf{B}}} & 2^{U\mathsf{B}} \end{array}$$

commutes. Given $w : U\mathsf{B} \to X$ and spelling out the definition of the vertical arrows, we obtain $P_{\mathsf{A}}(w \circ Uf) = P_{\mathsf{B}}(w) \circ Uf$, that is, $P_{\mathsf{A}}(w \circ Uf, a) = P_{\mathsf{B}}(w, Uf(a))$ for all $a \in U\mathsf{A}$, which yields condition (5) in Lemma 4.5. □

The theorem is important to us for the following reasons. First, it gives an abstract characterisation of invariance under behavioural equivalence as a naturality condition.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

For example, for any functor $F : \mathsf{Set} \to \mathsf{Set}$, the behavioural predicates of $\mathsf{FCoalg}(F)$ are precisely the natural transformations $U \to 2$ where $U : \mathsf{FCoalg}(F) \to \mathsf{Set}$ is the forgetful functor. Similarly, modal operators can be characterised as natural transformations, as explained in detail in Section 6.

Second, since every (X, Y)-ary operation of a signature for coalgebras gives rise to a natural transformation $X^U \to Y^U$, it shows that, in the case of coalgebras over Set, coalgebraic operations are predicate transformers that respect behavioural equivalence.

The next section is devoted to a detailed study of natural transformations $X^U \to Y^U$.

5. Implicit operations

In universal algebra, an implicit operation is a natural transformation $A^X \to A$. If free algebras exist, each implicit operation is explicit (that is, given by a term). If free algebras do not exist, implicit operations are more powerful. For example, in the case of algebras over finite sets, Reiterman (1982) showed that every variety is definable by equations in implicit operations (but not by equations in explicit operations). This section shows how to define covarieties of coequational categories using implicit operations.

If Σ is a signature and $U : \mathsf{SCoalg}(\Sigma) \to \mathscr{X}$ the forgetful functor, then each (X, Y)-ary operation symbol $\sigma \in \Sigma$ determines a natural transformation

$$\sigma : X^U \to Y^U.$$

So does each (X, Y)-ary term. This leads us to define (X, Y)-ary **implicit operations**, for every functor $U : \mathscr{A} \to \mathscr{X}$, as natural transformations

$$X^U \to Y^U.$$

Definition 5.1 (Coequations in implicit operations). Let $U : \mathscr{A} \to \mathscr{X}$ be a faithful functor and $X, Y \in \mathscr{X}$. Having two (X, Y)-ary implicit operations σ_1 and σ_2 in $\mathscr{A}$, we say that an object $\mathsf{A} \in \mathscr{A}$ satisfies the coequation $\sigma_1 = \sigma_2$ and write $\mathsf{A} \models \sigma_1 = \sigma_2$ iff $(\sigma_1)_{\mathsf{A}} = (\sigma_2)_{\mathsf{A}}$.

Remark 5.2.

1 If the functor U has a right adjoint R, then (X, Y)-ary implicit operations correspond to natural transformations

$$\mathscr{A}(-, RX) \to \mathscr{A}(-, RY),$$

that is, to morphisms $RX \to RY$. Consider implicit operations σ_1, σ_2 represented by $s_1, s_2 : RX \to RY$, respectively. Then an object A satisfies the coequation $\sigma_1 = \sigma_2$ iff every morphism $h : A \to RX$ is equalised by s_1 and s_2, that is, iff h factors through an equaliser

$$\mathsf{S} \longrightarrow RX \underset{s_2}{\overset{s_1}{\rightrightarrows}} RY$$

(provided that the latter exists). This notion of a coequation as a subobject S of a cofree object RX is a special case of Manes (1976, Theorem 3.4, page 227). It was further investigated in Rutten (2000), Gumm (2001a), Roşu (2001), Kurz (2000)

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

and Hughes (2001). Conversely, for any subobject $\mathsf{S} \to RX$, take the cokernel pair $f, g : RX \to \mathsf{A}$ and compose it with $\eta_{\mathsf{A}} : \mathsf{A} \to RU\mathsf{A}$ given by the unit η of the adjunction $U \dashv R$. Then the pair $\eta_{\mathsf{A}} \circ f, \eta_{\mathsf{A}} \circ g$ produces the pair of natural transformations $X^U \to Y^U$ in our sense. Thus, in the presence of cofree coalgebras, our approach is equivalent to the coequations-as-subobjects-of-cofree-objects approach.

2 Without cofree coalgebras, there are related concepts of a coequation in Adámek and Porst (2001) and Reiterman (1983). They are subsumed by coequations in implicit operations.

The following fact is a consequence of Rosický (1981, 5.3). Covarieties are defined in the paragraph following Remark 3.8.

Proposition 5.3. Suppose that $U : \mathscr{A} \to \mathscr{X}$ is coequational. Then, for a collection E of coequations in implicit operations, the full subcategory of all objects satisfying each coequation from E is a covariety in $\mathscr{A}$.

Proof. The closedness under colimits is clear. If $m : \mathsf{B} \to \mathsf{A}$ is a U-split subobject of A, then $X^{Um} : X^{U\mathsf{A}} \to X^{U\mathsf{B}}$ is epi, which yields closedness under U-split subobjects. □

We now show that over Set, the converse holds. This is a co-Birkhoff theorem not relying on the existence of cofree coalgebras. The proof uses certain implicit operations defined in terms of the behavioural equivalence relations $\sim_X$ introduced in Definition 4.1.

Theorem 5.4. Let E be a coequational theory over Set. Then every covariety $\mathscr{C}$ in $\mathsf{ECoalg}(E)$ is definable by coequations in implicit operations.

Proof. For a set X and an E-coalgebra A, define


$$\varphi^X_{\mathsf{A}} : X^{U\mathsf{A}} \to 2^{U\mathsf{A}}$$

as follows: for $v : U\mathsf{A} \to X$ and $a \in U\mathsf{A}$, let $\varphi^X_{\mathsf{A}}(v)(a) = 1$ iff there are $\mathsf{C} \in \mathscr{C}$, $u : U\mathsf{C} \to X$, $c \in U\mathsf{C}$, such that $(\mathsf{A}, v, a) \sim_X (\mathsf{C}, u, c)$ (see Definition 4.1). φ^X is an implicit operation by Theorem 4.6.

Consider coequations


$$\varphi^X = \mathit{true} \tag{6}$$

where *true* is the implicit operation $\mathit{true}_{\mathsf{A}} : X^{U\mathsf{A}} \to 2^{U\mathsf{A}}$ induced by the constant function $X \to 2, x \mapsto 1$. Each $\mathsf{C} \in \mathscr{C}$ satisfies (6). Conversely, assume that A satisfies all coequations (6). Then, by Proposition 4.3, for each $a \in U\mathsf{A}$, there are $\mathsf{B}_a \in \mathsf{ECoalg}(E)$, $\mathsf{C}_a \in \mathscr{C}$, and homomorphisms f_a, g_a and mappings w_a, u_a such that




https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

commutes and $(Uf_a)(a) \in (Ug_a)(U\mathsf{C}_a)$. Using a multiple pushout of the f_a



we get



with $(Uf)(U\mathsf{A}) \subseteq \bigcup\{U(f'_a \circ g_a)(U\mathsf{C}_a) : a \in U\mathsf{A}\}$. Note that $\bigcup\{U(f'_a \circ g_a)(U\mathsf{C}_a) : a \in U\mathsf{A}\}$ is the carrier of an E-coalgebra, which is in $\mathscr{C}$ due to closure under coproducts and quotients. Since f is injective and $\mathscr{C}$ is closed under subobjects, it follows that $\mathsf{A} \in \mathscr{C}$. □

Remark 5.5.

1 If $\mathsf{ECoalg}(E)$ has cofree coalgebras, the implicit operation φ^X is induced by a morphism $h : RX \to R2$, $U \dashv R$, or, equivalently, by a mapping $\tilde{h} : URX \to 2$. Then, for $v : U\mathsf{A} \to X$,

$$\varphi^X_{\mathsf{A}}(v) = \tilde{h} \circ Uv^\sharp,$$

with $v^\sharp : \mathsf{A} \to RX$ being the transpose of v.

2 Our proof works in the universe of finite sets, that is, every covariety of finite coalgebras is given by coequations in implicit operations. This is the 'Reiterman Theorem' (Reiterman 1982) for coalgebras.

3 Note that it does not follow from the theorem that $\mathscr{C}$ is coequational. Hence the theorem does not solve Problem 3.12. The reason is that the interpretations of implicit operations do not need to be coequationally determined by equations from E.

6. Modal logic

We associate to each category $\mathscr{A}$ equipped with a functor $U : \mathscr{A} \to \mathsf{Set}$ a modal logic, called the internal modal language of $\mathscr{A}$, and show that modal formulae and coequations have the same expressive power. Since coequations for coalgebras dualise equations for algebras, this leads to a new formalisation of the statement that modal logic for coalgebras is dual to equational logic for algebras.

To start with, let us take a brief look at the classical modal logic of Kripke frames, see, for example, Blackburn *et al.* (2001) for details. We use KF to denote the category of Kripke frames and bounded morphisms (a bounded morphism is a function whose graph is a bisimulation), that is, $\mathsf{KF} = \mathsf{FCoalg}(\mathscr{P})$. Writing $U : \mathsf{KF} \to \mathsf{Set}$ for the forgetful functor mapping Kripke frames to their carriers, the semantics $[\![\varphi]\!]$ of a modal formula φ in propositional variables $\{p_i \mid i \in I\}$ can be understood as a KF-indexed class of operations

$$[\![\varphi]\!]_{\mathsf{A}} : \left(\prod_I 2\right)^{U\mathsf{A}} \to 2^{U\mathsf{A}}, \quad \mathsf{A} \in \mathsf{KF},$$

that is, each $[\![\varphi]\!]_{\mathsf{A}}$ maps valuations $v \in (\prod_I 2)^{U\mathsf{A}}$ and elements $a \in U\mathsf{A}$ to truth values $[\![\varphi]\!]_{\mathsf{A}}(v, a) \in 2 = \{0, 1\}$. A central feature of modal logic is that formulae are invariant under bisimulation. That is, for a modal formula φ and two Kripke models (A, v), (B, w),

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

and $a \in A$, $b \in B$ we have

$$a, b \text{ bisimilar } \Rightarrow [\![\varphi]\!]_{\mathsf{A}}(v, a) = [\![\varphi]\!]_{\mathsf{B}}(w, b) \tag{7}$$

But we have seen in Theorem 4.6 that (7) is equivalent to $[\![\varphi]\!]_{\mathsf{A}}$ being natural in A. If we take invariance under bisimulation as the essence of modal logic, it therefore makes sense to consider any natural transformation $(\prod_I 2)^U \to 2^U$ as a modal operator and any category $\mathscr{A}$ equipped with a functor $U : \mathscr{A} \to \mathsf{Set}$ as a semantic domain for modal logic. This leads us to the following definition.

Definition 6.1. The *internal modal language* of a functor $U : \mathscr{A} \to \mathsf{Set}$ is given as follows. For each set I and each $i \in I$ the propositional variables p_i^I are I-ary formulae. For each natural transformation $\mu : (\prod_I 2)^U \longrightarrow 2^U$ there is an I-ary modal operator $\Box_\mu$. If φ_i are J-ary formulae, then $\Box_\mu\langle\varphi_i\rangle$ is a J-ary formula. The semantics of formulae is given by

$$[\![\Box_\mu\langle\varphi_i\rangle]\!] = \left(\prod_J 2\right)^U \xrightarrow{\langle[\![\varphi_i]\!]\rangle} \prod_I (2^U) \cong \left(\prod_I 2\right)^U \xrightarrow{\mu} 2^U.$$

Given an I-ary formula φ and $v : U\mathsf{A} \to \prod_I 2$, we write $\mathsf{A}, v, a \models \varphi$ iff $[\![\varphi]\!]_{\mathsf{A}}(v, a) = 2$ and $\mathsf{A} \models \varphi$ iff $[\![\varphi]\!]_{\mathsf{A}}(v) = U\mathsf{A}$ for all $v : U\mathsf{A} \to \prod_I 2$. For 0-ary formulae we drop v from the notation.

Remark 6.2.

1 Behavioural equivalence is characterised by formulae of the language, that is, for each $\mathsf{A} \in \mathscr{A}$ and $a \in U\mathsf{A}$ there is a (0-ary) formula φ_a such that $\mathsf{B}, b \models \varphi_a \Leftrightarrow a \sim b$.

2 The restriction of sets of colours to products of 2 allows for a notion of substitution of formulae for propositional variables. As usual in categorical logic, the semantics of substitution is given by composition.

3 The basic calculus for this modal logic is given by the axioms of infinitary propositional logic plus the rule 'from $\varphi \leftrightarrow \psi$ derive $\Box\varphi \leftrightarrow \Box\psi$' for each modal operator $\Box$. If, for example, $\Box$ is a normal modal operator, that is, the axioms $\Box \mathit{true} = \mathit{true}$ and $\Box(p_1 \wedge p_2) \leftrightarrow \Box p_1 \wedge \Box p_2$ are added, then we obtain a calculus that is equivalent to the basic modal logic **K**.

4 The definition of the internal modal language makes sense over other categories besides the category Set. For example, if we replace Set by the category of Stone spaces and let 2 be the discrete two-element space, then the continuity of the valuations $v : U\mathsf{A} \to \prod_I 2$ expresses the fact that the extensions of the propositions p_i have to be admissible (that is, clopen).

Example 6.3. The following examples illustrate our notion of modal operator as a natural transformation $(\prod_I 2)^U \to 2^U$.

1 An I-ary *boolean operator* is a modal operator $f^U : (\prod_I 2)^U \longrightarrow 2^U$ given by functions $f : \prod_I 2 \to 2$ such as, for example, *true*, $\to$, $\bigwedge_I$: compare this with Example 3.5.

2 *Atomic propositions* are 0-ary modal operators and are given by natural transformations $1 \to 2^U$ (or also $U \to 2$).

3 All commonly considered modal operators are examples. This includes non-normal modal operators, polymodal operators, modal operators given by natural

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

relations (Pattinson 2001) or predicate liftings (Pattinson (to appear)), and recursively defined modalities as in dynamic logic or the μ-calculus.

4 Examples of modalities that are not covered by Definition 6.1 can be obtained by definitions that require a 'change of structure'. For instance, consider $\mathsf{A} = (A, \alpha) \in \mathsf{FCoalg}(\mathscr{P})$ and define

$$\mathsf{A}, v, a \models \Box(\varphi, \psi) \;=\; \begin{cases} \mathsf{A}^\varphi, v^\varphi, a \models \psi & \text{if } \mathsf{A}, v, a \models \varphi \\ \mathit{true} & \text{otherwise} \end{cases}$$

where $\mathsf{A}^\varphi = (A^\varphi, \alpha^\varphi)$ is given by $A^\varphi = A \setminus \{a : \mathsf{A}, v, a \not\models \varphi\}$ and $\alpha^\varphi, v^\varphi$ are the restriction of α, v to A^φ. Modalities of this kind, often denoted by $[!\varphi]\psi$, arise in epistemic logic, see Baltag *et al.* (1999).

We can now translate results for implicit operations into results for modal logic. For example, the following statement is a corollary of Theorem 5.4 and Remark 5.5.2. To prove it, we just have to determine the modal language that is expressive enough to describe the formulas φ^X appearing in the proof of Theorem 5.4.

Proposition 6.4 (A Reiterman theorem for modal logic). If a class $\mathscr{K}$ of finite Kripke frames is closed under finite coproducts, quotients and embeddings, then $\mathscr{K}$ is definable by a set $\{\varphi^X \mid X = 2^n, n \in \mathbb{N}\}$ of formulae φ^X, each of which is a (countable) disjunction

$$\varphi^X \;=\; \bigvee_{\mathsf{B} \in \mathscr{K}, w : B \to X, b \in B} \psi^{(\mathsf{B}, w, b)}$$

of finitary formulae $\psi^{(\mathsf{B}, w, b)}$.

Finally, let us make precise the relationship between coequations in implicit operations and modal logic.

Proposition 6.5. Let $U : \mathscr{A} \to \mathsf{Set}$ be a functor. Coequations in implicit operations and formulae of the internal modal language have the same expressive power.

Proof. Each modal formula φ is logically equivalent to the coequation $\varphi = \mathit{true}$, which is to say that $\mathsf{A} \models \varphi$ (Definition 6.1) iff $\mathsf{A} \models \varphi = \mathit{true}$ (Definition 5.1) for all $\mathsf{A} \in \mathscr{A}$. Conversely, for a coequation $t = s$ with $t, s : X^U \to Y^U$, we find a set I and a surjective function $e : \prod_I 2 \to X$ and a modal formula $t \circ e \leftrightarrow s \circ e : (\prod_I 2)^U \to 2^U$ such that $\mathsf{A} \models t = s \;\Leftrightarrow\; \mathsf{A} \models t \circ e \leftrightarrow s \circ e$ for all $\mathsf{A} \in \mathscr{A}$. □

Remark 6.6 (Duality of modal and equational logic). Consider $U : \mathscr{A} \to \mathsf{Set}$. A (coalgebraic) implicit operation for U is the same as an (algebraic) implicit operation for U^{op}. $A^{\mathrm{op}} \in \mathscr{A}$ satisfies the (algebraic) equation $\sigma_1 = \sigma_2$ iff A satisfies the modal formula $\sigma_1 \leftrightarrow \sigma_2$. A satisfies the modal formula φ iff A^{op} satisfies the equation $\varphi = \mathit{true}$.

7. Davis's theorem

We may allow signatures with (X, Y)-ary operation symbols where X and Y are classes. We call them **meta-signatures**. In the same way as signatures yield coequational categories, meta-signatures lead to **meta-coequational categories**. Every meta-coequational category

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press


is co-Beck (see the paragraph before Proposition 3.10). Davis (1972) proved the converse. However, he overstated his result by claiming that every co-Beck category is coequational, which is not true as Example 7.3 shows. The second author observed Davis's mistake in Rosický (1981); here we give a full explanation. First, we sketch an argument proving Davis's theorem.

Theorem 7.1 (Davis). A concrete category is meta-coequational iff it is co-Beck.


Proof. Let $U : \mathscr{C} \to \mathsf{Set}$ create colimits and U-split equalisers. Let $R : \mathsf{Class} \to \mathsf{Class}$ be the density comonad of

$$\bar{U} : \mathscr{C} \xrightarrow{U} \mathsf{Set} \hookrightarrow \mathsf{Class}.$$


It means that RX is the colimit of the canonical diagram $(\bar{U}\downarrow X) \to \mathsf{Class}$



where ε_X is induced by the cone given by v's. Since $\bar{U}$ creates colimits, $(\bar{U}\downarrow X)$ is ∞-filtered (= every small subcategory of $(\bar{U}\downarrow X)$ has an upper bound). Since the (illegitimate) category $\mathsf{CCoalg}(R)$ of coalgebras for the comonad R is coequational over Class, it suffices to prove that the image of the comparison functor



consists precisely of the R-coalgebras (X, ξ) with X a set. Any such coalgebra is given by a V-split equaliser



where δ is the comultiplication. Since RRX and RX are given by ∞-filtered colimits, this equaliser is an ∞-filtered colimit of U-split equalisers in $\mathscr{C}$ (we are also using the fact that U creates U-split equalisers). Since X is a set, some R-coalgebra homomorphism $c_i : \mathsf{C}_i \to (X, \xi)$ splits, that is, $c_i \circ s = \mathrm{id}_X$ for some $s : (X, \xi) \to \mathsf{C}_i$. Hence (X, ξ) is isomorphic to some $\mathscr{C}$-object. □

Remark 7.2. Let $\mathscr{C}$ be a concrete category and $\sigma : X^U \to Y^U$ be an implicit operation where X, Y are classes. Take mappings $f : X_1 \to X$ and $g : Y \to Y_1$ where X_1, Y_1 are sets. We get an implicit operation $g^U \circ \sigma \circ f^U$ where arities are sets. If sets are

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

codense in classes, then Y is a canonical limit of the canonical diagram $(Y\downarrow\mathsf{Set}) \to \mathsf{Class}$ consisting of $g : Y \to Y_1$ where Y_1 is a set. Hence, the implicit operation $\sigma \circ f^U$ is determined by implicit operations $g^U \circ \sigma \circ f^U$. Moreover, since $U\mathsf{C}$ are sets, σ is determined by implicit operations $g^U \circ \sigma \circ f^U$. This is what Davis claimed. However, it does not mean that coequations of implicit operations whose arities are classes can be replaced by coequations of implicit operations whose arities are sets. There is a problem with compositions $X^U \xrightarrow{\sigma_1} Z^U \xrightarrow{\sigma_2} Y^U$ where Z is a proper class. The precise result is Proposition 5.5 in Rosický (1981).

As before (see the proof of Proposition 3.7), sets are codense in classes iff *Ord* is not measurable, that is, iff each *Ord*-complete ultrafilter is principal. A model of such a set theory is V_α where α is inaccessible but not measurable. On the other hand, in the theory of finite sets, that is, in V_ω, is $Ord = \omega$ measurable.

Example 7.3. Let Σ consist of a single $(1, Ord)$-ary operation symbol σ. Then Σ-coalgebras A are sets A equipped with an operation $\sigma_\mathsf{A} : 1^A \to Ord^A$, that is, with a mapping $\alpha : A \to Ord$. Homomorphisms $h : (A, \alpha) \to (B, \beta)$ are mappings $h : A \to B$ such that $\beta \circ h = \alpha$. $\mathsf{SCoalg}(\Sigma)$ is a (legitimate) meta-coequational category. It cannot be isomorphic to any full subcategory of $\mathsf{FCoalg}(F)$ for any functor $F : \mathsf{Set} \to \mathsf{Set}$ because it contains a proper class of one-element coalgebras.

Each mapping $f : Ord \to m$ gives a $(1, m)$-ary term (that is, a $(1, m)$-ary implicit operation) $x_f \cdot \sigma$. In fact, every implicit operation $\varphi : 1^U \to m^U$ is of that kind. It suffices to take $f : Ord \to m$ given as $f(p) = \varphi_\mathsf{P}(\mathrm{id}_1)$ where P is the one-element Σ-coalgebra with σ_P taking the value p (see Rosický (1981, 7.2) for the easy calculation that $\varphi = x_f \cdot \sigma$.)

Let Σ_1 be the collection of all $(1, m)$-ary operation symbols $\sigma_f, f : Ord \to m$. Σ_1 is not a signature because it is larger than a class. If *Ord* is not measurable, then, following Rosický (1981, 5.5), $\mathsf{SCoalg}(\Sigma)$ is described in Σ_1 by coequations $x_g \cdot \sigma_f = \sigma_{g \circ f}$, for $f : Ord \to m$ and $g : m \to k$. In fact, having a Σ_1-coalgebra $(A, (\sigma_f)_\mathsf{A})$ satisfying these equations, we get a cone $(\sigma_f)_\mathsf{A} : A \to m$ of the canonical diagram $(Ord\downarrow\mathsf{Set})$ and, therefore, the induced mapping $\alpha : A \to Ord$. (A, α) is the Σ-coalgebra determined by $(A, (\sigma_f)_\mathsf{A})$.

If *Ord* is measurable, then $\mathsf{SCoalg}(\Sigma)$ is not coequational. In fact, it is shown in Rosický (1981, 7.2) that *Ord*-complete ultrafilters provide one-element coalgebras living in all coequational categories containing $\mathsf{SCoalg}(\Sigma)$.

Acknowledgements

The first author thanks Alexandru Baltag for pointing out an example of a modal operator that is not an implicit operation and Yde Venema for the observation that the morphisms of monotone neighbourhood frames coincide with the morphisms of (2^{2^-})-coalgebras. The second author has profited from discussions with H. P. Gumm. Diagrams were produced using Paul Taylor's macro package.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press

References

Aczel, P. and Mendler, N. P. (1989) A final coalgebra theorem. In: Pitt, D. H., Rydeheard, D. E., Dybjer, P., Pitts, A. M. and Poigné, A. (eds.) Category Theory and Computer Science. *Springer-Verlag Lecture Notes in Computer Science* **389** 357–365.

Adámek, J. and Porst, H.-E. (2001) From varieties of algebras to covarieties of coalgebras. In: Corradini, A., Lenisa, M. and Montanari, U. (eds.) Coalgebraic Methods in Computer Science (CMCS'01). *ENTCS* **44.1**.

Adámek, J. and Rosický, J. (1994) *Locally Presentable and Accessible Categories*, London Mathematical Society Lecture Notes Series **189**, Cambridge University Press.

Adámek, J. and Trnková, V. (1990) *Automata and Algebras in Categories*, Kluwer Academic Publishers.

Baltag, A., Moss, L. and Solecki, S. (1999) The logic of public announcements, common knowledge, and private suspicions. Technical Report SEN-R9922, CWI, Amsterdam.

Blackburn, P., de Rijke, M. and Venema, Y. (2001) *Modal Logic*, Cambridge University Press.

Davis, R. (1972) Quasi-cotripleable categories. *Proceedings of the American Mathematical Society* **35** 43–38.

Goldblatt, R. (1976) Metamathematics of modal logic I. *Reports on Mathematical Logic* **6** 41–78.

Gumm, H. P. (2001a) Equational and implicational classes of colgebras. *Theoretical Computer Science* **260**.

Gumm, H. P. (2001b) Functors for coalgebras. *Algebra Universalis* **45** 135–147.

Hughes, J. (2001) *A Study of Categories of Algebras and Coalgebras*, Ph.D. thesis, Carnegie Mellon University, Pittsburgh.

Johnstone, P. (1982) *Stone Spaces*, Cambridge University Press.

Kurz, A. (2000) *Logics for Coalgebras and Applications to Computer Science*, Ph.D. thesis, Ludwig-Maximilians-Universität München.

Linton, F. (1966) Some aspects of equational categories. In: Eilenberg, S., Harrison, D., Lane, S. M. and Röhrl, H. (eds.) *Proceedings of the Conference on Categorical Algebra, La Jolla, 1965*, Springer 84–91.

Linton, F. (1969) An outline of functorial semantics. In: Eckmann, B. (ed.) Seminar on triples and categorical homology theory. *Springer-Verlag Lecture Notes in Mathematics* **80** 7–52.

Mac Lane, S. (1971) *Category Theory for the Working Mathematician*, Springer.

Manes, E. G. (1976) *Algebraic Theories*, Springer.

Pattinson, D. (to appear) Coalgebraic modal logic: Soundness, completeness and decidability of local consequence. *Theoretical Computer Science*.

Pattinson, D. (2001) Semantical principles in the modal logic of coalgebras. In: Proceedings 18th International Symposium on Theoretical Aspects of Computer Science (STACS 2001). *Springer-Verlag Lecture Notes in Computer Science* **2010**.

Reiterman, J. (1982) The Birkhoff theorem for finite algebras. *Algebra Universalis* **14** 1–10.

Reiterman, J. (1983) Algebraic theories and varieties of functor algebras. *Fundamenta Mathematicae* **118** 59–68.

Roşu, G. (2001) Equational axiomatizability for coalgebra. *Theoretical Computer Science* **260** 229–247.

Rosický, J. (1981) On algebraic categories. In: Universal Algebra (Proc. Coll. Esztergom 1977). *Colloq. Math. Soc. J. Bolyai* **29** 662–690.

Rutten, J. (2000) Universal coalgebra: A theory of systems. *Theoretical Computer Science* **249** 3–80.

Smyth, M. (1983) Power domains and predicate transformers: a topological view. In: Diaz, J. (ed.) Proceedings 10th International Colloquium on Automata, Languages and Programming. *Springer-Verlag Lecture Notes in Computer Science* **154** 662–675.

https://doi.org/10.1017/S0960129504004402 Published online by Cambridge University Press