Combinatorics, Probability and Computing (2003) 12, 515–545. © 2003 Cambridge University Press
DOI: 10.1017/S0963548303005881 Printed in the United Kingdom

Ramsey Games Against a One-Armed Bandit

EHUD FRIEDGUT,[1] YOSHIHARU KOHAYAKAWA,[2†]
VOJTĚCH RÖDL,[3‡] ANDRZEJ RUCIŃSKI[4§] and PRASAD TETALI[5¶]

[1]Einstein Institute of Mathematics, The Hebrew University of Jerusalem,
Jerusalem, 91904, Israel
(e-mail: ehudf@math.huji.ac.il)

[2]Instituto de Matemática e Estatística, Universidade de São Paulo,
Rua do Matão 1010, 05508-900 São Paulo, Brazil
(e-mail: yoshi@ime.usp.br)

[3]Department of Mathematics and Computer Science,
Emory University, Atlanta, GA 30322, USA
(e-mail: rodl@mathcs.emory.edu)

[4]Faculty of Mathematics and Computer Science,
Adam Mickiewicz University, Poznań, Poland,
and
Department of Mathematics and Computer Science,
Emory University, Atlanta, GA 30322, USA
(e-mail: rucinski@amu.edu.pl)

[5]School of Mathematics, Georgia Institute of Technology,
Atlanta, GA, 30332, USA
(e-mail: tetali@math.gatech.edu)

Received 20 June 2002; revised 8 July 2003

We study the following one-person game against a random graph process: the Player's goal is to 2-colour a random sequence of edges $e_1, e_2, \ldots$ of a complete graph on n vertices, avoiding a monochromatic triangle for as long as possible. The game is over when a monochromatic triangle is created. The online version of the game requires that the Player should colour each edge as it comes, before looking at the next edge.

† Partially supported by MCT/CNPq through ProNEx Programme (Proc. CNPq 664107/1997–4) and by CNPq (Proc. 300334/93–1 and 468516/2000–0).
‡ Partially supported by NSF Grant 0071261. The collaboration between the second and third authors is supported by a CNPq/NSF cooperative grant.
§ Supported by KBN grant 2 PO3A 015 23.
¶ Partially supported by NSF grant DMS-0100298.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

While it is not hard to prove that the expected length of this game is about $n^{4/3}$, the proof of the upper bound suggests the following relaxation: instead of colouring online, the random graph is generated in only two rounds, and the Player colours the edges of the first round before the edges of the second round are thrown in. Given the size of the first round, how many edges can there be in the second round for the Player to be likely to win? In the extreme case, when the first round consists of a random graph with $cn^{3/2}$ edges, where c is a positive constant, we show that the Player can win with high probability only if *constantly* many edges are generated in the second round.

1. Introduction

In this paper, we study *Ramsey one-person triangle avoidance games against a random graph process*. In all versions of our game, the goal of the player, called throughout the *Painter*, is to colour randomly generated edges, using a given number of colours, without creating a monochromatic triangle. The object of this paper is to study how long the Painter can 'survive', that is, stay in the game. We distinguish three versions of the game: *online*, *two-round*, and *offline*.

1.1. The three games

Let us discuss the variants of our game separately.

1.1.1. The online game. In the *online* version of our game, the Painter receives a random sequence of edges, and she has to colour each edge as it comes, avoiding monochromatic triangles as long as possible.

It turns out that, when the number of colours available to the Painter is $r = 2$, this game is quite easy to analyse; indeed, a fairly simple 'local' argument shows that this game typically lasts for $\Theta(n^{4/3})$ steps.

We observe that the length of the game is not so sharply determined. When the Painter has seen only $o(n^{4/3})$ edges, she is essentially certain to be safe; she is most unlikely to survive up to $\omega n^{4/3}$ edges, for any ω with $\omega = \omega(n) \to \infty$ as $n \to \infty$; but, for any $c > 0$, she has probability bounded away from 0 and 1 of surviving up to $cn^{4/3}$ edges.

1.1.2. The two-round game. What happens if the Painter has a *mercy period*? That is, suppose she is allowed *not* to colour the first, say, N_0 edges online, but may wait to see *all of them* before committing herself to the colour of those edges. Clearly, she should be able to survive longer, because her initial colouring can be cleverer. Moreover, instead of switching back to the online version after the mercy period of N_0 edges, let us suppose that another N_1 random edges are now presented to the Painter. If she is able to extend her initial colouring of the N_0 edges to the union of the N_0 and the N_1 edges still avoiding monochromatic triangles, she wins; otherwise, she loses. This is the *two-round* version of our game.

We shall analyse the two-round game with $N_0 = cn^{3/2}$, for c a small positive constant. (We shall see in a moment why this is the interesting choice for N_0.) The result here is that, with such a value of N_0, if the Painter has $r = 2$ colours available, then typically she can win only if $N_1 = O(1)$: for any $N_1 = N_1(n)$ that tends to infinity, the probability that the Painter wins tends to 0.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

We shall also show that, if $N_0 = cn^{3/2}$ and $r = 3$ colours are available, then the breakpoint for N_1 is of order n: if $N_1 = o(n)$, then the Painter typically wins; if $N_1 = \omega n$, where $\omega = \omega(n)$ tends to infinity, then the Painter typically loses. If $N_1 = \Theta(n)$, then she wins with probability bounded away from 0 and 1. Note that, as in the online game, the transition is not sharp.

1.1.3. The offline game. In the offline game, an n-vertex graph $G(n, N)$ with a fixed number of edges $N = N(n)$ is generated uniformly at random and is presented to the Painter; her task is to colour the edges of $G(n, N)$ avoiding monochromatic triangles.

The reader familiar with the theory of random graphs will immediately see that the analysis of this game amounts exactly to determining the threshold for the Ramsey property $\mathscr{R}_r$, which consists of all graphs such that every r-colouring of their edges results in a monochromatic triangle.

The threshold for the property $\mathscr{R}_r$ has been investigated for some time now. In particular, it has been proved that there exist constants c_r and C_r such that

$$\lim_{n\to\infty} \mathbb{P}[G(n, N) \in \mathscr{R}_r] = \begin{cases} 1 & \text{if } N > C_r n^{3/2}, \\ 0 & \text{if } N < c_r n^{3/2}. \end{cases} \tag{1.1}$$

The above threshold of $n^{3/2}$ was verified for $r = 2$ in [17] (see also [8]), and then proved for arbitrary $r \geqslant 2$ in [18]. It is interesting to note that the dependence on r is not present in the exponent of n, but only in the constants.

Returning to the case $r = 2$, the current proofs leave a wide gap between c_2 and C_2. Using a criterion from [9] for the existence of sharp thresholds, that gap was recently removed in [10]. In the usual parlance, the Ramsey property $\mathscr{R}_2$ admits a *sharp threshold*. We shall come back to this in Section 2. Note that the existence of a sharp threshold for $\mathscr{R}_2$ means that the offline game for $r = 2$ colours has a sharply determined value, unlike the online and the two-round games.

1.1.4. Further remarks. It is the result in (1.1) that suggests the investigation of the two-round game with N_0 of order $n^{3/2}$. Note that the game would be trivial if $N_0 > C_r n^{3/2}$, for in that case the chances of the Painter surviving even the first round are rather slim. On the other hand, if $r = 2$ and $N_0 = cn^{3/2}$ with $c < c_r$, then it first seems that the Painter should be happy, because she has a very good chance of being able to colour the edges of the first round. However, our result (Theorem 3.3 below) shows that, at least for $r = 2$, her happiness is short-lived: *regardless of how small c is*, the fact that she has to colour those N_0 edges *before seeing the final graph* makes it very unlikely that she will succeed in colouring even as few as, say, $N_1 = \log\log\log n$ further edges! This is in contrast to the fact that, if $N_0 + N_1 < c_r n^{3/2}$, then, typically, $G(n, N_0 + N_1)$ may be 2-edge-coloured avoiding monochromatic triangles (see (1.1)).

We mention in passing that in Section 2 we shall attempt to explain how the two-round version of our game naturally arose from a technical part of the proof of the existence of a sharp threshold for $\mathscr{R}_2$.

Interestingly enough, both the two-round and online versions of our Ramsey game are much more sensitive to the number of colours available to the Painter than the offline one is. Recall that, for example, in the two-round game, while for $r = 2$ colours the survival

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

time is $O(1)$, for $r = 3$ colours the survival time is of order n. For the online, 3-colour game, we can easily deduce a lower bound of roughly $n^{7/5}$ for its length, which is much larger than for two colours. At this point, we do not have a matching upper bound for the 3-colour game. We will expand on this a bit in Section 3.1.

1.2. Organization

This paper is organized as follows. In Section 2 we briefly discuss the proof of the existence of a sharp threshold for the property $\mathscr{R}_2$, and its relation to this paper. In Section 3 we state and prove our results on Ramsey games. As it turns out, most of the work will go into proving a certain delicate lemma (Lemma 3.2); Section 4 will be entirely devoted to proving this lemma. In Section 5 we prove some additional auxiliary lemmas. In Section 6 we briefly discuss some open problems.

1.3. Notation and terminology

Our notation is fairly standard. We let $[n] = \{1, \dots, n\}$. We write $[X]^k$ for the family of k-element subsets of X. Let $G = (V(G), E(G))$ be a graph. If x is a vertex of G, then $N_G(x) = N(G; x)$ denotes the neighbourhood of x in G. We write $\deg_G(x) = |N_G(x)|$ and $\Delta(G) = \max\{\deg_G(x) : x \in V(G)\}$. If $U \subseteq V(G)$, we write $G[U]$ for the subgraph of G induced by U. Thus, $E(G[U]) = E(G) \cap [U]^2$. We put $e_G(U)$ for $|E(G[U])|$, the number of edges induced by U in G, and $N_G(U) = \bigcup_{x \in U} N_G(x)$.

If $U, W \subseteq V(G)$ are disjoint, then $G[U, W]$ stands for the bipartite graph with vertex classes U and W naturally induced by U and W in G. We write $E_G(U, W)$ for the set of edges of G with one endvertex in U and the other in W. That is, $E_G(U, W) = E(G[U, W])$. We let $e_G(U, W) = |E_G(U, W)|$. We sometimes identify a graph with its edge set.

We define, for a graph F, the *base graph* $\mathrm{Base}(F)$ on the vertex set $V(F)$ as follows: a pair e of vertices of F is an edge of $\mathrm{Base}(F)$ if it forms a triangle with two edges of F. A colouring of the edges of a graph will be called *proper* if it does not contain a monochromatic triangle.

We shall consider the three standard models for random graphs: the binomial random graphs $G(n, p)$, the uniform random graphs $G(n, N)$, and the random graph process $G_0 \subseteq \dots \subseteq G_{\binom{n}{2}}$, which naturally corresponds to random permutations $e_1, \dots, e_{\binom{n}{2}}$ of the edges of the complete graph on $[n]$. In particular, if we stop a random graph process at time t, we obtain a random graph in the uniform model $G(n, t)$. When convenient, we switch between these models without discussion. A rule of thumb is that most results can be 'translated' back and forth between the binomial and uniform models, provided $N \sim p\binom{n}{2}$. For more on the (asymptotic) equivalence of these models and on random graphs in general, we refer the reader to [6] and [12]. Another feature we will use is the monotonicity. If a graph property Q is increasing (preserved under edge addition), and $p_1 \leqslant p_2$, then $\mathbb{P}(G(n, p_1) \in Q) \leqslant \mathbb{P}(G(n, p_2) \in Q)$.

Most of the time we have a parameter n, and we are interested in the case in which $n \to \infty$. As usual, we use the term 'asymptotically almost surely' (a.a.s.) to mean 'with probability tending to 1 as $n \to \infty$'. Therefore, we often tacitly assume that n is large enough (for instance, for some inequalities to hold). Also, we drop the $\lfloor\ \rfloor$ and $\lceil\ \rceil$ from our formulae when they are not important. This will simplify the exposition considerably.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Our logarithms are natural logarithms.

2. Sharp thresholds

The questions considered in this paper arose naturally during the work on [10], which studies the notion of sharp threshold for the property $\mathscr{R}_2$. We briefly outline the ideas that arise in that paper in order to explain the background to the present work. However, the remainder of this paper does *not* depend on this section.

Recall that $\mathscr{R}_2$ consists of all graphs such that every 2-colouring of their edges results in a monochromatic triangle. Below, we refer to the graphs in $\mathscr{R}_2$ as *Ramsey graphs*. The main result in [10] is the following theorem, asserting that property $\mathscr{R}_2$ admits a sharp threshold.

Theorem 2.1. *There exists a function $b = b(n) = \Theta(1)$ such that for all $\varepsilon > 0$ we have*

$$\lim_{n\to\infty} \mathbb{P}[G(n,p) \in \mathscr{R}_2] = \begin{cases} 1 & \text{if } p > (1+\varepsilon)b(n)/\sqrt{n}, \\ 0 & \text{if } p < (1-\varepsilon)b(n)/\sqrt{n}. \end{cases}$$

The proof of Theorem 2.1 is based on a consequence of Friedgut's criterion from [9], which we now discuss. Let M be an arbitrary, fixed balanced graph with average degree 4, and let M^* be a random copy of M in K_n. It follows from [9] that, in order for $\mathscr{R}_2$ to have a sharp threshold, it is sufficient to prove the following:

(*) For any choice of the constants c, C, α, and $\xi > 0$, and for any M as above, there is a graph property $\mathscr{G}$ such that

(a) if $p = p(n)$ satisfies $c/\sqrt{n} \leqslant p \leqslant C/\sqrt{n}$, then $G(n,p) \in \mathscr{G}$ a.a.s.,

and, moreover, such that, for all n-vertex graphs $G \in \mathscr{G}$ with sufficiently large n,

(b) if $\mathbb{P}[G \cup M^* \in \mathscr{R}_2] > \alpha$, then

$$\mathbb{P}[G \cup G(n, \xi p) \in \mathscr{R}_2] \to 1 \text{ as } n \to \infty.$$

Note that, roughly speaking, the implication in (b) means that every $G \in \mathscr{G}$ is such that, if G may be turned with positive probability into a Ramsey graph by adding a random copy of M, then G turns a.a.s. into a Ramsey graph when $(1+o(1))\xi p\binom{n}{2}$ edges are sprinkled randomly.

We recall that M is a small graph, and hence the addition of M^* to G is a 'local' change, whereas adding $G(n, \xi p\binom{n}{2})$ is 'global'. Moreover, observe that (b) is nontrivial only when $G \notin \mathscr{R}_2$. Also, because of our conditions on M, it can be proved that $M \notin \mathscr{R}_2$ (see [10, 16]).

Let us now say a few words about the verification of Friedgut's condition (*). We take for $\mathscr{G}$ a certain well-defined pseudorandom property of graphs with n vertices and $(1+o(1))p\binom{n}{2}$ edges (see [10, Definition 6.1]) and we assume that the assumption in implication (b) holds. As observed above, we may assume that $G \notin \mathscr{R}_2$ and we have $M \notin \mathscr{R}_2$. A fairly complicated and lengthy argument, involving a tailor-made regularity lemma (see [10]), leads to the conclusion that, for some $\lambda > 0$, there exists a 'small' family $\mathscr{K}$ of 'large' subgraphs of G such that there is always a monochromatic copy of a member

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

of $\mathscr{K}$ in every proper colouring of the edges of G. More precisely, the family $\mathscr{K}$ is such that

(i) $|\mathscr{K}| = 2^{o(n^{3/2})}$,
(ii) for all $K \in \mathscr{K}$, we have $|E(K)| > \lambda |E(G)|$,

and, most importantly,

(iii) for every proper 2-colouring of the edges of G, there is a $K \in \mathscr{K}$ that is monochromatic.

Now we observe that, in order to prove the conclusion of (b), it is enough to find, for every $K \in \mathscr{K}$, a triangle $T = T(K) \subseteq G(n, \xi p)$, such that $T \subseteq \mathrm{Base}(K)$, *i.e.*, a triangle T such that each of the edges of T forms a triangle together with a path of length two in K. (Indeed, in this case, if K is monochromatic, there is no way to extend the colouring of G to a proper colouring of $G \cup G(n, \xi p)$.)

To establish the existence of $T = T(K)$ above, we first define the following property of a graph G:

(†) Given two real numbers $0 < \lambda < 1$ and $0 < a < 1/6$, we say that G has property $\mathscr{T}(\lambda, a)$ if, for any subgraph F of G with at least $\lambda |E(G)|$ edges, the graph $\mathrm{Base}(F)$ contains at least $a|V(G)|^3$ triangles.

Since the $K \in \mathscr{K}$ has at least $\lambda |E(G)|$ edges (it satisfies (ii) above), if $\mathscr{T}(\lambda, a)$ holds for $G \in \mathscr{G}$, then we may conclude that every $K \in \mathscr{K}$ is such that $\mathrm{Base}(K)$ contains at least $a|V(G)|^3$ triangles. As it turns out (see the following lemma), we may *define* $\mathscr{G}$ such that (a) holds and $\mathscr{T}(\lambda, a)$ holds for every $G \in \mathscr{G}$.

Lemma 2.2. *For any $c > 0$ and $\lambda > 0$, there exists $a > 0$ such that if $p \geqslant c/\sqrt{n}$ then a.a.s. the random graph $G(n, p)$ has property $\mathscr{T}(\lambda, a)$.*

Lemma 2.2 allows us to require that $\mathscr{T}(\lambda, a)$ should hold for all $G \in \mathscr{G}$.

Now we are ready to establish the existence of $T = T(K)$ for all $K \in \mathscr{K}$. Indeed, for any fixed $K \in \mathscr{K}$, the fact that $\mathrm{Base}(K)$ contains at least $a|V(G)|^3$ triangles implies, via Janson's inequality [11], that, with probability $1 - 2^{-\Theta(n^{3/2})}$, there is a triangle $T(K)$ in $\mathrm{Base}(K) \cap G(n, \xi p)$. Since (i) holds, we have that a suitable triangle $T = T(K)$ does exist a.a.s. for every $K \in \mathscr{K}$. The conclusion of (b) now follows from (iii).

To summarize, the convergence

$$\mathbb{P}[G \cup G(n, \xi p) \in \mathscr{R}_2] \to 1$$

follows indirectly from a statement about the two-round Ramsey game against a random graph: the proof involves extending proper colourings of graphs of size $\Theta(n^{3/2})$. This explains why Lemma 2.2 and a related result, Lemma 3.2 given below, are useful tools in studying such games.

3. Ramsey triangle avoidance games

This section contains our main results. In Section 3.1 below, we prove a relatively simple result for the online version of the Ramsey game. In Section 3.2, we state two results

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

on the two-round game, together with their proofs, pending a major technical lemma, Lemma 3.2, which will be proved in Section 4.

3.1. The online game

Consider the random graph process

$$e_1, \dots, e_{\binom{n}{2}}$$

on n vertices (see Section 1.3), revealing its edges one by one, and the following one-person game related to it. The Painter's task is to 2-colour the edges online, with colours red and blue, say, and not to create a monochromatic triangle for as long as she can. Here *online* means that the Painter has to decide on the colour of e_i before e_{i+1} is generated. The game is over when a monochromatic triangle is created (this does happen eventually if $n \geqslant 6$). For how long can she stay in the game with probability approaching 1 as $n \to \infty$?

More formally, for each strategy π of the Painter, let X_π be the length of the game, *i.e.*, X_π is the largest index N such that $e_1, \dots, e_N$ contains no monochromatic triangle. The goal of the game is to choose a strategy that maximizes the expectation $\mathbb{E}(X_\pi)$. By a *strategy* we mean a complete set of rules assigning a colour to the next edge.

Since $e_1, \dots, e_N$ is exactly the random graph $G(n, N)$, clearly, by (1.1), an upper bound on the expected duration of the game is $C_2 n^{3/2}$. We shall soon show that the threshold for the length of the online game is in fact much smaller.

A crucial tool in our investigation is the base graph $\mathrm{Base}(F)$ of a graph F. Recall that a pair e of vertices of F is an edge in $\mathrm{Base}(F)$ if it forms a triangle with two edges of F. Now consider an online 2-colouring of the edges $e_1, \dots, e_{\binom{n}{2}}$. Let R_i and B_i be the sets of the edges e_j $(j \leqslant i)$ coloured so far with colours red and blue, respectively. The Painter is stuck if and only if

$$e_{i+1} \in \mathrm{Base}(R_i) \cap \mathrm{Base}(B_i).$$

Similarly, the game is bound to be over by time j if for some $i < j$, the intersection of the set of edges $\{e_{i+1}, \dots, e_j\}$ with $\mathrm{Base}(R_i)$ (or with $\mathrm{Base}(B_i)$) contains a triangle.

Let us begin with a trivial lower bound. As long as $N = o(n^{6/5})$, there is a.a.s. no copy of a diamond in $e_1, \dots, e_N$, that is, a K_4 with an edge omitted, and the Painter never gets stuck until the Nth edge, since a.a.s.

$$e_{i+1} \notin \mathrm{Base}(R_i) \cap \mathrm{Base}(B_i)$$

for each $i < N$.

But she can do better: as long as $N = o(n^{4/3})$ a.a.s. neither a copy of K_4 nor a copy of the pyramid graph (see Figure 1) has emerged, and so $\mathrm{Base}(G(n, N)) \cap G(n, N)$ does not contain a triangle. Hence the greedy strategy π_0 of colouring each edge red, unless such a move would create a red K_3, is successful. Indeed, $\mathrm{Base}(R_N)$, which is a subgraph of $\mathrm{Base}(G(n, N))$, is triangle-free, and no blue triangle will be created. We have just proved that a.a.s. $X_{\pi_0} > n^{4/3}/\omega$ for any $\omega = \omega(n)$ with $\omega \to \infty$.

Quite surprisingly, $n^{4/3}$ is the right threshold for the length of the random online triangle avoidance game.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press




Figure 1. The pyramid, the enhanced pyramid, and the double pyramid

Theorem 3.1. *For every ω with $\omega = \omega(n) \to \infty$ as $n \to \infty$ and for every strategy π of the Painter, a.a.s. $X_\pi < \omega n^{4/3}$.*

Proof. We relax the online regime by allowing a *mercy period* of $N_0 = n^{4/3}$ steps; *i.e.*, only after N_0 edges are generated do they have to be coloured by the Painter. We shall argue that no matter how they are coloured, the Painter loses a.a.s. after another $(\omega - 1)n^{4/3}$ edges. This is nothing else but a two-round exposure technique (see [12]), with a colouring in between.

The aim is to show that a.a.s. every subgraph of $G(n, N_0)$ with at least half of the edges (the majority colour) has at least $\Omega(n^2)$ triangles in its base. Then, by the second moment method, any $(\omega - 1)n^{4/3}$ additional random edges will a.a.s. contain at least one of these triangles and the game will be over.

A.a.s. the number of copies of $K_{2,3}$ in $G(n, N_0)$, denoted by $X(K_{2,3})$, is smaller than, say, $n \log n$. Let R be the majority colour in $G(n, N_0)$ and let $d_1, \ldots, d_n$ be the degree sequence in R. Then the number of triangles in the base of R is at least

$$\sum_{i=1}^{n} \binom{d_i}{3} - X(K_{2,3}) \geqslant n \binom{N_0/n}{3} - X(K_{2,3}) = \Omega(n^2),$$

where the last equation holds a.a.s. □

The reader will have no difficulty in verifying that the proofs above show that, for any $c > 0$, the Painter has probability bounded away from 0 and 1 of surviving up to $cn^{4/3}$ edges.

With three colours at hand the Painter can survive quite a bit longer. Sadly, we do not know the precise threshold for the 3-colour case, as we cannot match our lower bounds with a corresponding upper bound. In fact the only upper bound we have is the trivial $n^{3/2}$ coming from the offline case. The analysis below is intended mainly to show that, unlike the offline case, the power of n in the online threshold depends on the number of colours.

For any such online Ramsey game, where the Painter tries to avoid a monochromatic copy of H using r colours, define the following simple ordered–greedy strategy: the Painter orders her palette of colours arbitrarily $c_1, c_2, \ldots, c_r$ and then always uses the highest ranking colour that does not complete a monochromatic copy of H. It would interesting to see an example where this is *not* asymptotically optimal.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Returning to the case of 3-colour online triangle avoidance, we now argue that the Painter, using the ordered–greedy strategy, can survive at least $\Theta(n^{7/5})$ rounds. The meticulous referee noted that in fact our argument can give $n^{13/9}$, since in what follows below we only use the fact that red and blue are greedily selected over yellow, but not the fact that red is greedily selected over blue. However, we do not include the full, more complicated, analysis of this subtlety.

Let the Painter use the ordered–greedy strategy, say π_1, which colours an edge red if possible, otherwise blue if possible, and finally yellow if none of red and blue will do. Using π_1, the Painter gets stuck only if a copy of the double pyramid, or of the enhanced pyramid (see Figure 1), or of K_5 with an edge omitted has emerged. Since such subgraphs a.a.s. do not appear in the random graph with $o(n^{7/5})$ edges, we obtain that $\mathbb{P}(X_{\pi_1} > n^{7/5}/\omega) \to 1$. As noted above, using the full power of the ordered–greedy assumption gives a longer, more complex, list of graphs which are not ordered–greedy-colourable.

3.2. The two-round game

The proof of Theorem 3.1 bridges the online and the two-round game, which we now define formally. Given two integers N_0 and N_1, first, N_0 edges are generated randomly (let $G_0 = G(n, N_0)$ be the graph defined by these edges), and the Painter colours them *properly* with r colours (if possible), that is, avoiding monochromatic triangles. Then a second round of N_1 random edges (graph G_1) is thrown in and the Painter wins if her colouring of G_0 has an extension to a proper colouring of $G_0 \cup G_1$ (if G_0 cannot be properly coloured, the Painter has already lost at that stage).

For a given strategy π of the Painter, let $P_\pi(r, N_0, N_1)$ be the probability that the Painter wins the game when using π. Here, by a *strategy*, we mean an assignment of an r-edge-colouring to every graph with N_0 edges. Thus, the goal of the Painter is to find a strategy π that maximizes $P_\pi(r, N_0, N_1)$. Let

$$P^*(r, N_0, N_1) = \max_\pi P_\pi(r, N_0, N_1).$$

The above proof of Theorem 3.1 gives that $P^*(2, n^{4/3}, \omega n^{4/3}) = o(1)$, because the base of the majority colour in $G_0 = G(n, n^{4/3})$ contains $\Theta(n^2)$ triangles. This simple argument may be extended to a wider range of N_0 (for $r = 2$), namely,

$$n^{7/6} \leqslant N_0 \leqslant cn^{3/2},$$

where here and below, unless stated otherwise, c is always assumed to be so small that

$$\mathbb{P}(G(n, cn^{3/2}) \in \mathscr{R}_2) = o(1).$$

Indeed, along similar lines, one may easily prove that there are $\Theta(N_0^3/n^2)$ triangles in the base graph of the majority colour of $G_0 = G(n, N_0)$, and consequently, arguing as in the proof of Theorem 3.1, we have

$$P^*(2, N_0, \omega n^{8/3}/N_0) = o(1).$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

In particular, in the most extreme case, we obtain that the number of triangles in the base graph of the majority colour of $G_0 = G(n, cn^{3/2})$ is $\Theta(n^{5/2})$, and thus $P^*(2, cn^{3/2}, \omega n^{7/6}) = o(1)$.

Using Lemma 2.2 with $\lambda = 1/2$ and with F consisting of the edges coloured by the majority colour, we can improve the above and conclude that in fact a.a.s., for any 2-edge-colouring of $G(n, cn^{3/2})$, the number of triangles in the base graph of the majority colour is of maximal possible order $\Theta(n^3)$. By the second moment method this yields that

$$P^*\big(2, cn^{3/2}, \omega n\big) = o(1)$$

(a triangle is needed in the base of one colour). To improve this any further we need to find a stronger tool.

As we have just seen, Lemma 2.2, put in the colouring context, says that the base of the majority colour is rich in triangles. A powerful extension of that result, Lemma 3.2 below, which is the main technical result of this paper, implies that the same is true for the intersection of the bases of two colours, provided the colouring is proper. This result not only settles the two-round game for $r = 2$, but also for $r = 3$, the other interesting case.

Lemma 3.2. *For all $c > 0$, there are constants α and $\beta > 0$ such that with $p = c/\sqrt{n}$ the random graph $G = G(n, p)$ a.a.s. satisfies the following property. For every proper 3-colouring $E(G) = R \cup B \cup Y$ of the edges of G, there are two colours, say R and B, such that, letting*

$$I = \mathrm{Base}(R) \cap \mathrm{Base}(B), \tag{3.1}$$

we have

(i) $|E(I)| \geqslant \alpha n^2$

and, in fact,

(ii) *the graph I contains at least βn^3 triangles.*

Remark. Note that Lemma 2.2 is *not* a special case of the seemingly stronger Lemma 3.2. In Lemma 2.2 the parameter λ may be arbitrarily small, while in Lemma 3.2 the assumption that the colouring must be proper sets an implicit lower bound on the size of the second largest colour (see (5.13)). In particular, it seems that Lemma 3.2 cannot replace Lemma 2.2 in the proof of Theorem 2.1 in [10].

Nevertheless, the proof of Lemma 3.2, besides some new ideas, uses the ideas from the proof of Lemma 2.2. The relation between the proofs of Lemmas 2.2 and 3.2 is briefly discussed in Section 4.2.

Taking Lemma 3.2 for granted, we are ready to state and prove our results for the two-round game.

Theorem 3.3. *For any ω with $\omega = \omega(n) \to \infty$ as $n \to \infty$, we have*

$$P^*\big(2, cn^{3/2}, \omega\big) = o(1).$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Proof. This follows immediately from Lemma 3.2(i) (with $Y = \emptyset$), since a.a.s. at least one of the ω new edges will fall into I. □

Finally, let us consider the two-round game for $r > 2$ and $N_0 = cn^{3/2}$. If $r \geqslant 4$, then a trivial conservative strategy is to use only two colours for G_0 and then another two colours for G_1. This, together with the threshold result from [18] settles the game at $\Theta(n^{3/2})$, *i.e.*, there exist constants b and $B > 0$ for which we have

$$\lim_{n\to\infty} P^*\big(r, cn^{3/2}, N_1\big) = \begin{cases} 1 & \text{if } N_1 < bn^{3/2}, \\ 0 & \text{if } N_1 > Bn^{3/2}. \end{cases}$$

In the remaining case $r = 3$, we have the following, complete solution.

Theorem 3.4. *We have*

$$\lim_{n\to\infty} P^*\big(3, cn^{3/2}, N_1\big) = \begin{cases} 1 & \text{if } N_1 \ll n, \\ 0 & \text{if } N_1 \gg n. \end{cases}$$

Proof. The '1-statement', that is, the fact that $P^*(3, cn^{3/2}, N_1) \to 1$ as $n \to \infty$, follows by using only two colours for G_0, and then exclusively the third colour for G_1. As long as $N_1 = o(n)$, a.a.s. there will not be any triangles in G_1. The 0-statement follows from the second moment method and Lemma 3.2(ii). □

We remark that the proof above shows that $P^*(3, cn^{3/2}, c'n)$ is bounded away from 0 and 1 for any $c' > 0$.

4. Pairs at distance two in red and blue

Our aim in this section is to prove Lemma 3.2. On the way we shall come close to proving Lemma 2.2 (see Section 4.2). However, we refer the reader to [10] for the complete proof of Lemma 2.2.

Let us introduce a piece of terminology. Let ϱ and d be positive reals. We say that a graph H is (ϱ, τ)-*dense* if, for all $U \subseteq V(H)$ with $|U| \geqslant \varrho |V(H)|$, we have

$$e_H(U) \geqslant \tau \binom{|U|}{2}.$$

We now state a strengthening of Lemma 3.2, which, together with an easy property of (ϱ, τ)-dense graphs, implies Lemma 3.2.

Lemma 4.1. *For all $c > 0$, there is a constant $\tau > 0$ such that, for any $\varrho > 0$, there is a constant $\sigma > 0$ such that for $p = c/\sqrt{n}$ the random graph $G = G(n, p)$ a.a.s. satisfies the following property. For every proper 3-colouring $E(G) = R \cup B \cup Y$ of the edges of G, there are two colours, say R and B, for which, letting*

$$I = \mathrm{Base}(R) \cap \mathrm{Base}(B),$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

there exists a set $U \subseteq V(G)$ *such that*

(i) $|U| \geqslant \sigma n$ *and*

(ii) $I[U]$ *is* (ϱ, τ)*-dense.*

Graphs that are (ϱ, τ)-dense for small ϱ are 'rich in small subgraphs'; in particular, the following result holds [19].

Lemma 4.2. *For all* $\tau > 0$ *there exist* ϱ, n_0, *and* $c_0 > 0$ *such that every* (ϱ, τ)*-dense graph on* $n \geqslant n_0$ *vertices contains at least* $c_0 n^3$ *triangles.*

With Lemmas 4.1 and 4.2 at hand, it is easy to prove Lemma 3.2.

Proof of Lemma 3.2. Given $c > 0$, Lemma 4.1 gives us a constant $\tau > 0$. By Lemma 4.2 we obtain constants ϱ, n_0, and $c_0 > 0$ such that any u-vertex (ϱ, τ)-dense graph with $u \geqslant n_0$ contains at least $c_0 u^3$ triangles. We now feed this constant $\varrho > 0$ to Lemma 4.1, to get a constant $\sigma > 0$.

We claim that the choice of

$$\alpha = \frac{1}{3}\tau\sigma^2 \tag{4.1}$$

and

$$\beta = c_0\sigma^3, \tag{4.2}$$

will do in Lemma 3.2.

To prove our claim, suppose we have a proper 3-colouring of the edges of $G = G(n, p)$. By Lemma 4.1, a.a.s. there are two colours, say R and B, and a set $U \subseteq V(G)$, $|U| \geqslant \sigma n$, for which $I[U] = (\mathrm{Base}(R) \cap \mathrm{Base}(B))[U]$ is (ϱ, τ)-dense. Then

$$|E(I)| \geqslant |E(I[U])| \geqslant \tau\binom{|U|}{2} \geqslant \frac{1}{3}\tau|U|^2 \geqslant \left(\frac{1}{3}\tau\sigma^2\right)n^2 = \alpha n^2.$$

Moreover, by the choice of $\varrho > 0$ and c_0, we deduce from Lemma 4.2 that the number of triangles in the (ϱ, τ)-dense graph $I[U]$ is at least

$$c_0|U|^3 \geqslant c_0\sigma^3 n^3 = \beta n^3.$$

Lemma 3.2 is proved. □

Let us now briefly comment on the organization of this section. The proof of Lemma 4.1 fills the remainder of this section, and will require considerable work. This proof is based on some technical, auxiliary results presented in Section 4.1 and proved in Section 5. The actual proof of Lemma 4.1 is given in Section 4.2.

4.1. Towards the proof of Lemma 4.1

In this section, we state three lemmas and prove two claims that will be crucial in the proof of Lemma 4.1.

Throughout this section and the following one we assume that

$$p = \frac{c}{\sqrt{n}},$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press




Figure 2. The amalgam given by Lemma 4.3

where $0 < c \leqslant 1/5$ is an arbitrary constant that we fix once and for all. We also define once and for all the constants

$$\sigma_0 = (1/200)c^2, \quad \lambda_0 = (1/30)c^2 \quad \text{and} \quad \tau_0 = \frac{1}{800}\lambda_0^2 c^2 \sigma_0. \tag{4.3}$$

Let H be a graph. A pair of disjoint sets U, $V \subseteq V(H)$ will be called $(p, \lambda; H, \varepsilon)$-*lower semi-regular*, or $(p, \lambda; H, \varepsilon)$-*semi-regular* for short, if, for all $U' \subseteq U$, $V' \subseteq V$, with $|U'| \geqslant \varepsilon|U|$ and $|V'| \geqslant \varepsilon|V|$, we have

$$\frac{e_H(U', V')}{p|U'||V'|} \geqslant \lambda.$$

Note that if (U, V) is $(p, \lambda; H, \varepsilon)$-semi-regular then

$$e_H(U, V) \geqslant \lambda p|U||V|. \tag{4.4}$$

Note also that then (U, V) is $(p, \lambda; H, \varepsilon')$-semi-regular for any $\varepsilon' \geqslant \varepsilon$, and (U^*, V) is $(p, \lambda; H, \varepsilon/\varrho)$-semi-regular for any $U^* \subseteq U$ with $|U^*| \geqslant \varrho|U|$.

Our first auxiliary lemma, Lemma 4.3, concerns the existence of a certain substructure in a properly edge-coloured $G(n, p)$. A tripartite graph J on vertex sets Z, U, and W with no edge between Z and W will be called an *amalgam* (of two bipartite graphs). The amalgam guaranteed by Lemma 4.3 in a properly edge-coloured $G(n, p)$ satisfies certain properties with respect to its size, colouring and structure (see Figure 2).

Lemma 4.3. *For all $\varepsilon > 0$, there is $\sigma > 0$ for which the following assertion holds a.a.s. for $G = G(n, p)$. Suppose*

$$E(G) = R \cup B \cup Y \tag{4.5}$$

is a proper colouring of the edges of G. Then there exist two colours, say R and B, and a subgraph J of G which is an amalgam with vertex sets Z, U, and W, as follows:

(i) $|Z| = |W| = \sigma_0 n$ *and* $|U| \geqslant \sigma n$,
(ii) *we have* $E_J(Z, U) \subseteq R$ *and* $E_J(U, W) \subseteq B$,
(iii) *the pairs* (Z, U) *and* (U, W) *are* $(p, \lambda_0; J, \varepsilon)$-*semi-regular*.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Basically, our second lemma concerns the base of the subgraph $J_0 = J[Z, U]$ of an amalgam J as in Lemma 4.3 (as well as, by symmetry, the base of $J_1 = J[U, W]$).

Lemma 4.4. *For all $\varrho > 0$, there is an $\varepsilon > 0$ such that a.a.s. $G = G(n,p)$ satisfies the following property. For all bipartite subgraphs J_0 of G with bipartition (Z, U) and such that*

(i) *$z = |Z| = \sigma_0 n$ and $u = |U| \geqslant n/\log n$, and*
(ii) *(Z, U) is $(p, \lambda_0; J_0, \varepsilon)$-semi-regular,*

the graph $\mathrm{Base}(J_0)[U]$ *is (ϱ, τ_0)-dense.*

Lemma 4.4 follows immediately from the next claim. Let J_0^* be a bipartite graph with bipartition (Z, U^*). Recall that we have the constants σ_0, λ_0 and τ_0, defined in (4.3). Given $\varepsilon' > 0$, we shall say that J_0^* is *ε'-rare* if

(i*) $z = |Z| = \sigma_0 n$ and $u^* = |U^*| \geqslant n/(\log n)^2$,
(ii*) (Z, U^*) is $(p, \lambda_0; J_0^*, \varepsilon')$-semi-regular, and
(iii*) $|\mathrm{Base}(J_0^*)[U^*]| < \tau_0 \binom{u^*}{2}$.

Claim 4.5. *If $\varepsilon' > 0$ is small enough, then a.a.s. $G = G(n,p)$ contains no ε'-rare bipartite subgraph.*

To deduce Lemma 4.4 from Claim 4.5 take $\varepsilon = \varepsilon' \varrho$. In turn, Claim 4.5 follows easily from a special case of Lemma 4.7 below.

Proof of Claim 4.5. Put $F = K_{u^*}$, $\tau' = 1$, and $\alpha = \lambda_0/2e$ into Lemma 4.7 obtaining the desired value of ε'. We will apply the first moment method. By Lemma 4.7, the expected number of ε'-rare bipartite subgraphs of $G(n,p)$ is at most

$$\sum_{u^*} \sum_{T} \binom{n}{z} \binom{n}{u^*} \alpha^T \binom{u^* z}{T} p^T \leqslant \sum_{u^*} \sum_{T} n^{u^*} \left(\frac{\alpha e}{\lambda_0} \right)^T < n^n 2^{-\Omega(n^{3/2}/(\log n)^2)} = o(1).$$

The summation above extends over all integers $n/(\log n)^2 \leqslant u^* \leqslant n$ and $\lambda_0 pzu^* \leqslant T \leqslant \binom{n}{2}$. Note that $\lambda_0 pzu^* \geqslant \lambda_0 \sigma_0 c n^{3/2}/(\log n)^2$. □

As an immediate consequence of Lemmas 4.3 and 4.4 we conclude that, for all $\varrho > 0$, there exists $\sigma > 0$ such that a.a.s. $G = G(n,p)$ contains an amalgam J that satisfies (i)–(iii) of Lemma 4.3 and such that $\mathrm{Base}(J_0)[U]$ is (ϱ, τ_0)-dense. Suppose that $I[U] = (\mathrm{Base}(R) \cap \mathrm{Base}(B))[U]$ is *not* (ϱ, τ)-dense for some τ. Then, because

$$I[U] = (\mathrm{Base}(R) \cap \mathrm{Base}(B))[U] \supseteq (\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U],$$

the graph $(\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U]$ is not (ϱ, τ)-dense either. This means that there exists a set $U^* \subseteq U$, with $u^* = |U^*| \geqslant \varrho |U|$, such that

$$|(\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U^*]| < \tau \binom{u^*}{2}.$$

The subset U^* generates a subamalgam H of J, which, as our next result shows, is very unlikely to appear in $G(n,p)$.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Now come the details. Let Z, U^*, and W be three pairwise disjoint sets of vertices. Given ε' and n, let us call an amalgam H with tripartition (Z, U^*, W) ε'-*rare* if

(i) we have

$$|Z| = |W| = \sigma_0 n, \tag{4.6}$$

$$u^* = |U^*| \geqslant n/(\log n)^2, \tag{4.7}$$

$$\Delta(H) \leqslant 2c\sqrt{n}, \tag{4.8}$$

(ii) the pairs (Z, U^*) and (U^*, W) are $(p, \lambda_0; H, \varepsilon')$-semi-regular,
(iii) letting $H_0 = H[Z, U^*]$ and $H_1 = H[U^*, W]$,

$$|\mathrm{Base}(H_0)[U^*]| \geqslant \tau_0 \binom{|U^*|}{2},$$

and

(iv) we have

$$|(\mathrm{Base}(H_0) \cap \mathrm{Base}(H_1))[U^*]| < \tau_0^3 \binom{|U^*|}{2}. \tag{4.9}$$

Note the symmetry of the roles of H_0 and H_1 in the above definition. We will make use of it later in the proof.

Claim 4.6. *If $\varepsilon' > 0$ is small enough, then a.a.s. $G = G(n, p)$ contains no ε'-rare amalgam.*

Claim 4.6 follows from Lemma 4.7, a delicate result indicating that some bipartite graphs are rare. In order to state Lemma 4.7, we need to introduce more definitions and notation.

Let U^* and W be two disjoint sets of vertices. Suppose $F \subseteq [U^*]^2$ is a graph with vertex set U^*. Suppose now that we have a constant $\varepsilon' > 0$ and integers n and T. Suppose further that $w = |W| = \sigma_0 n$, $u^* = |U^*| \geqslant n/(\log n)^2$, and $|F| = \tau' \binom{u^*}{2}$. We call a bipartite graph H with bipartition (U^*, W) (F, ε')-*rare* if

(i) $|E| = T$ and $\Delta(H) \leqslant 2c\sqrt{n}$,
(ii) (U^*, W) is $(p, \lambda_0; H, \varepsilon')$-semi-regular, and
(iii) we have

$$|F \cap \mathrm{Base}(H)[U^*]| < \tau_0 \tau' |F|. \tag{4.10}$$

Finally, let

$$\mathscr{B}_1 = \mathscr{B}_1(U^*, W; F; \tau', \varepsilon'; n, T) \tag{4.11}$$

be the family of all (F, ε')-rare, bipartite graphs H. We are now ready to state our third auxiliary lemma, the formulation of which is based on the notation and definitions introduced above.

Lemma 4.7. *Let any constants $\tau' > 0$ and $0 < \alpha \leqslant 1$ be given. Then there exist constants $\varepsilon' > 0$ and n_0 such that, with the notation as above, for all $n \geqslant n_0$, all F as above, and*

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

all $0 \leqslant T \leqslant u^*w$, *we have*

$$|\mathscr{B}_1(U^*, W; F; \tau', \varepsilon'; n, T)| \leqslant \alpha^T \binom{u^*w}{T}.$$

Lemma 4.7 is a statement concerning the uniform space of the *bipartite graphs* with vertex classes U^* and W, and T edges. Indeed, this lemma asserts that a random element of this space is extremely unlikely to be a member of the family $\mathscr{B}_1$ defined in (4.11), as the probability that this happens can be made subexponentially small.

The proof of Lemma 4.7 is postponed to Section 5.3.

Proof of Claim 4.6. We first show that, for a suitable ε', ε'-rare amalgams really are rare among all amalgams of that size. We feed $\tau' = \tau_0$ and

$$\alpha = \frac{1}{2}\left(\frac{\lambda_0}{e}\right)^2 \tag{4.12}$$

into Lemma 4.7, and obtain a constant $\varepsilon' > 0$. Let

$$\mathscr{B}_2 = \mathscr{B}_2(Z, U^*, W; \varepsilon'; n, T, T') \tag{4.13}$$

be the family of ε'-rare amalgams H on $Z \cup U^* \cup W$, with T edges with endpoints in W and T' edges with endpoints in Z.

Fact 4.8. *For all* T *and* T', *we have*

$$|\mathscr{B}_2(Z, U^*, W; \varepsilon'; n, T, T')| \leqslant \alpha^T \binom{zu^*}{T'} \binom{wu^*}{T}. \tag{4.14}$$

Proof. For any graph $H \in \mathscr{B}_2$, we let $H_0 = H[Z, U^*]$ and $H_1 = H[U^*, W]$. To show (4.14), we may count the graphs in $\mathscr{B}_2$ by estimating the number of H_0 crudely by

$$\binom{zu^*}{T'}, \tag{4.15}$$

and, for each fixed H_0, estimating the number of H_1 that we may 'put together with H_0' to form a graph H in $\mathscr{B}_2$.

Let us now fix a bipartite graph H_0 with bipartition (Z, U^*) satisfying conditions (i)–(iii) in the definition of ε'-rare amalgams. Moreover, let a subgraph

$$F \subseteq \mathrm{Base}(H_0)[U^*]$$

with $|F| = \tau_0 \binom{|U^*|}{2}$ be fixed. If for some H_1 with bipartition (U^*, W) the amalgam $H = H_0 \cup H_1 \in \mathscr{B}_2$, then

$$H_1 \in \mathscr{B}_1 = \mathscr{B}_1(U^*, W; F; \tau', \varepsilon'; n, T), \tag{4.16}$$

that is, H_1 is (F, ε')-rare. Indeed, conditions (i) and (ii) in the definition of $\mathscr{B}_1$ clearly hold for H_1. To check condition (iii), we observe that, in view of (4.9), we have

$$|F \cap \mathrm{Base}(H_1)[U^*]| \leqslant |(\mathrm{Base}(H_0) \cap \mathrm{Base}(H_1))[U^*]| < \tau_0^3 \binom{|U^*|}{2} = \tau_0^2 |F|.$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Therefore (4.10) holds (with $\tau' = \tau_0$), and so does condition (iii) in the definition of $\mathscr{B}_1$. We conclude that (4.16) is true.

To verify (4.14), it now suffices to put together the bounds in (4.15) and in Lemma 4.7 (with $\tau' = \tau_0$). □

We now use Fact 4.8 to show that the random graph $G(n,p)$ is quite unlikely to contain ε'-rare amalgams H.

Note that if H is an ε'-rare amalgam with $T + T'$ edges, then

$$T,\ T' \geqslant \lambda_0 p|Z||U^*| = \lambda_0 p|W||U^*| = \lambda_0\sigma_0 pnu^*, \tag{4.17}$$

and hence, as (4.7) holds, we have

$$T,\ T' = \Omega(n^{3/2}/(\log n)^2) \gg n\log n. \tag{4.18}$$

Without loss of generality we may assume that $T \geqslant T'$. We are now ready to show that a.a.s. no ε'-rare amalgam occurs as a subgraph in $G = G(n,p)$. As in the proof of Claim 4.5, we shall use the first moment method. The expected number of such subgraphs is

$$\sum |\mathscr{B}_2(Z, U^*, W; \varepsilon'; n, T, T')|\, p^{T+T'}, \tag{4.19}$$

where the sum is over all choices of Z, U^*, $W \subseteq V(G)$ and over all choices of T and T'. We use (4.12), (4.14), (4.17), and (4.18) to estimate (4.19). Below, we write $\sum_{u^*}$ for the sum over $n/(\log n)^2 \leqslant u^* \leqslant n$ and $\sum_{T,T'}$ for the sum over $T,\ T' \geqslant \lambda_0\sigma_0 pnu^*$ (see (4.17)). The quantity in (4.19) is at most

$$\begin{aligned}
&\sum_{u^*}\sum_{T,T'} \binom{n}{u^*}\binom{n}{w}\binom{n}{z}\alpha^T\binom{wu^*}{T'}\binom{zu^*}{T}p^{T'+T} \\
&\leqslant \sum_{u^*}\sum_{T,T'} n^{3n}\alpha^T\left(\frac{e}{\lambda_0}\right)^{2T} \leqslant n^{4n}\sum_{T}\left(\frac{1}{2}\right)^T \\
&\leqslant n^{4n}(n^2)2^{-\Omega(n^{3/2}/(\log n)^2)} = o(1).
\end{aligned}$$

Claim 4.6 follows by Markov's inequality. □

4.2. Proof of Lemma 4.1

We shall use Lemmas 4.3 and 4.4 and Claim 4.6 to prove Lemma 4.1.

Let $c > 0$ be given. A moment's thought shows that the property asserted a.a.s. for $G = G(n,p)$ is increasing, and therefore we may suppose that $c \leqslant 1/5$ (for details concerning this argument, see [12, Lemma 1.10], for instance). We set

$$\tau = \tau_0^3, \tag{4.20}$$

and claim that this choice for τ will do.

To prove our claim above, let an arbitrary $0 < \varrho \leqslant 1$ be given. We have to prove the existence of a suitable constant $\sigma > 0$. To this end, we first invoke Lemma 4.4 with this ϱ, to obtain a constant $\varepsilon(\text{L4.4}) > 0$. Moreover, Claim 4.6 gives us a constant

$$\varepsilon'(\text{C4.6}) > 0. \tag{4.21}$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

We now let

$$\varepsilon' = \min\{\varepsilon(\mathrm{L4.4})/\varrho, \varepsilon'(\mathrm{C4.6})\}, \tag{4.22}$$

and, finally, invoke Lemma 4.3 with $\varepsilon = \varepsilon'\varrho$ to obtain

$$\sigma = \sigma(\mathrm{L4.3}) > 0. \tag{4.23}$$

We claim that σ chosen above will do in Lemma 4.1. In the remainder of this proof we verify this claim.

It follows easily from the Chernoff bound that a.a.s.

$$\Delta(G(n,p)) \leqslant 2c\sqrt{n}. \tag{4.24}$$

Let $G = G(n,p)$ satisfy the conclusions of Lemmas 4.3 and 4.4 and Claim 4.6 with the above parameters, as well as $\Delta(G) \leqslant 2c\sqrt{n}$. (Note that all these properties hold a.a.s. for $G(n,p)$.)

Suppose that we have a colouring $E(G) = R \cup B \cup Y$ with no monochromatic triangles. We shall show that G must then contain the required set U.

Since G satisfies the conclusion of Lemma 4.3, there is an amalgam $J \subseteq G$ as in the statement of that lemma. We follow the notation of Lemma 4.3, so that, *e.g.*, $V(J) = Z \cup U \cup W$ (see Figure 2 as well).

As we have $|U| \geqslant \sigma n$, condition (i) in Lemma 4.1 holds. Hence, the proof of Lemma 4.1 will be finished if we show the following fact.

Fact 4.9. *The graph $I[U]$ is (ϱ,τ)-dense.*

Proof. Our proof strategy has already been unveiled just after the proof of Claim 4.5. Here we give the details.

Let $J_0 = J[Z,U]$ and $J_1 = J[U,W]$. Since G satisfies the conclusion of Lemma 4.4, the graph $\mathrm{Base}(J_0)[U]$ is (ϱ,τ_0)-dense.

We now claim that $(\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U]$ is (ϱ,τ)-dense. Note that, then, $I[U]$ must be (ϱ,τ)-dense, because

$$I[U] = (\mathrm{Base}(R) \cap \mathrm{Base}(B))[U] \supseteq (\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U].$$

Suppose for a contradiction that $(\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U]$ is not (ϱ,τ)-dense, and let $U^* \subseteq U$ be a witness to this fact, that is, suppose that

$$|U^*| \geqslant \varrho|U| \tag{4.25}$$

and

$$|(\mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U^*]| < \tau\binom{|U^*|}{2}. \tag{4.26}$$

We let H be the subgraph of J induced by $Z \cup U^* \cup W$. Also, let $H_0 = H[Z,U^*]$ and $H_1 = H[U^*,W]$, and let $T' = |E(H_0)|$ and $T = |E(H_1)|$. We claim that H is an ε'-rare amalgam. Condition (i) of the definition of an ε'-rare amalgam clearly holds (note that $\Delta(H) \leqslant \Delta(G)$).

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Since (Z, U) and (U, W) are $(p, \lambda_0; J, \varepsilon)$-semi-regular, by the choice of ε' (see (4.22)) and (4.25), we have that (Z, U^*) and (U^*, W) are $(p, \lambda_0; H, \varepsilon')$-semi-regular, so condition (ii) holds too.

We already know that $\mathrm{Base}(J_0)[U]$ is (ϱ, τ_0)-dense, so that

$$|\mathrm{Base}(H_0)[U^*]| = |\mathrm{Base}(J_0)[U^*]| \geqslant \tau_0 \binom{|U^*|}{2}, \tag{4.27}$$

which verifies condition (iii). Finally note that (4.26) yields (iv), since

$$\mathrm{Base}(H_0) \cap \mathrm{Base}(H_1))[U^*] = \mathrm{Base}(J_0) \cap \mathrm{Base}(J_1))[U^*].$$

We have reached a contradiction, for we supposed, in Claim 4.6, that G contains no ε'-rare amalgam. This contradiction completes the proof of Fact 4.9, and consequently the proof of Lemma 4.1. □

Comments on the proof of Lemma 2.2. We observed in the beginning of Section 4 that we would come close to proving Lemma 2.2. We may now expand on this remark a little (we shall be sketchy).

Suppose $p = c/\sqrt{n}$, where $c > 0$ is an arbitrary constant, and consider $G = G(n, p)$. Furthermore, suppose $F \subseteq E(G)$ has cardinality at least $\lambda_0 |E(G)|$ for some fixed $\lambda_0 > 0$. We wish to show that $\mathrm{Base}(F)$ contains an^3 triangles, for some constant $a > 0$.

Recall that Lemma 4.3 asserts the existence of a certain substructure in a $G(n, p)$ (an 'amalgam' of two bipartite graphs) whose edges have been properly 3-coloured (see Figure 2). In the case in which we have a subset of edges F of $G = G(n, p)$ as above, one may prove the existence, within F, of, say, the 'upper half' of the amalgam in Figure 2. In fact, using the notation introduced immediately after the statement of Lemma 4.3, one may prove the existence of the graph J_0. (The proof is roughly speaking 'contained' in the proof of Lemma 4.3.)

With a suitable graph J_0 at hand, one may apply Lemma 4.4, and deduce that $\mathrm{Base}(F)$ contains a (ϱ, τ)-dense, induced subgraph with at least σn vertices, where ϱ, τ, and σ are certain positive constants. It now follows from Lemma 4.2 that $\mathrm{Base}(F)$ must contain an^3 triangles, for some constant a, as required in Lemma 2.2.

5. Proofs of the auxiliary lemmas

Out of the three auxiliary lemmas stated in Section 4.1, only two remain to be proved, namely, Lemmas 4.3 and 4.7. The proofs of both results involve regular and semi-regular pairs, and hence we shall start with a section devoted to these concepts.

5.1. Semi-regularity

The notion of a $(p, \lambda; H, \varepsilon)$-semi-regular pair has already been defined on page 527. When dealing with semi-regular pairs, the following simple observation is often useful.

Lemma 5.1. *If (U, W) is $(p, \lambda; H, \varepsilon)$-semi-regular and $U_0 \subseteq U$ satisfies $|U_0| \geqslant \varepsilon |U|$, then more than $(1 - \varepsilon)|W|$ vertices of W have each at least $\lambda p |U_0|$ neighbours in U_0.*

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

The following powerful generalization of Lemma 5.1, is, in fact, its corollary. Recall that for $V \subseteq V(H)$, $N_H(V) = \bigcup_{v \in V} N_H(v)$ is the union of the neighbourhoods in H of all vertices $v \in V$.

Corollary 5.2. *Let (U, W) be a $(p, \lambda; H, \varepsilon)$-semi-regular pair, let $d < |W|$ be an integer, and fix $U_0 \subseteq U$ with $|U_0| > (\varepsilon/\mu)|U|$, where $\mu = \exp\{-\lambda p d/2\}$. Then the number of d-element subsets W_0 of W such that*

$$|U_0 \setminus N_H(W_0)| > \mu |U_0| \tag{5.1}$$

is less than

$$\varepsilon^{d/2} \binom{d}{d/2} |W|^d / d!.$$

Proof. Let $W_0 \subseteq W$ be a d-element subset of W with a fixed ordering $w_1, \ldots, w_d$ of its elements. We define a sequence

$$U_0 \supseteq \cdots \supseteq U_d = U_0 \setminus \bigcup_{w \in W_0} N(H; w) = U_0 \setminus N_H(W_0) \tag{5.2}$$

of descending subsets of W, putting

$$U_j = U_{j-1} \setminus N(H; w_j) = U_0 \setminus \bigcup_{1 \leqslant k \leqslant j} N(H; w_k), \tag{5.3}$$

for all $1 \leqslant j \leqslant d$. Let us call the ratio

$$|U_j|/|U_{j-1}| \tag{5.4}$$

the *shrinking factor at j* $(1 \leqslant j \leqslant d)$. Observe that this factor depends only on the sequence $w_1, \ldots, w_j$. Observe that for a set W_0 satisfying (5.1), at least $d/2$ shrinking factors are greater than $1 - \lambda p$. Indeed, otherwise we would have

$$|U_d| \leqslant (1 - \lambda p)^{d/2} |U_0| \leqslant \mu |U_0|.$$

Note also that for each $1 \leqslant j \leqslant d$, we have

$$U_{j-1} \geqslant \mu |U_0| > \varepsilon |U|$$

and hence, by Lemma 5.1 there are fewer than $\varepsilon |W|$ vertices $w \in W$ with less than $\lambda p |U_{j-1}|$ neighbours in U_{j-1}.

Let $\mathscr{W}^\times \subseteq W^d = W \times \cdots \times W$ be the collection of ordered d-tuples $(w_1, \ldots, w_d)$ that have at least $d/2$ indices j with shrinking factor greater than $1 - \lambda p$. It follows that

$$|\mathscr{W}^\times| \leqslant \binom{d}{d/2} \varepsilon^{d/2} |W|^d. \tag{5.5}$$

Now let $\mathscr{W} \subseteq [W]^d$ be the set of d-element subsets W_0 of W that satisfy (5.1). Clearly, if $W_0 \in \mathscr{W}$, then any ordered d-tuple $(w_1, \ldots, w_d)$ with $W_0 = \{w_1, \ldots, w_d\}$ belongs to $\mathscr{W}^\times$ (that is, any ordering of a member of $\mathscr{W}$ is in $\mathscr{W}^\times$). This and (5.5) give that

$$d! |\mathscr{W}| \leqslant |\mathscr{W}^\times| \leqslant \binom{d}{d/2} \varepsilon^{d/2} |W|^d,$$

which completes the proof. □

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

The reader may have noted that the concepts of (ϱ, τ)-denseness and ε-semi-regularity are related (the latter is a 'bipartite version' of the former). Furthermore, the reader acquainted with the celebrated regularity lemma of Szemerédi [21] has certainly observed that the notion of ε-semi-regularity is a weaker form of ε-regularity. As it turns out, it is more convenient for us to work with this weaker notion.

However, to be able to assert the existence of suitable semi-regular pairs in our graphs, we shall invoke the regularity lemma, which we now recall. Since we are dealing with sparse graphs (that is, graphs with a subquadratic number of edges), we shall make use of an appropriate variant. We need to introduce some definitions.

A graph H is called $(p; b, \beta)$-*bounded* if for all pairs of disjoint subsets $U, V \subseteq V(H)$ with $|U|, |V| > \beta|V(H)|$ we have $e_H(U, V) \leqslant bp|U||V|$. A pair $U, V \subseteq V(H)$, $U \cap V = \emptyset$, is called $(p; H, \varepsilon)$-*regular* if for all $U' \subseteq U$ and $V' \subseteq V$, with $|U'| \geqslant \varepsilon|U|$ and $|V'| \geqslant \varepsilon|V|$, we have

$$\left| \frac{e_H(U', V')}{p|U'||V'|} - \frac{e_H(U, V)}{p|U||V|} \right| \leqslant \varepsilon.$$

Observe that, clearly, if a pair is $(p; H, \varepsilon)$-regular, then it is also $(p, \lambda; H, \varepsilon)$-semi-regular for $\lambda = e_H(U, V)/p|U||V| - \varepsilon$.

Let $|V(H)| = n$. A partition $V(H) = V_1 \cup \cdots \cup V_k$ is $(p; H, \varepsilon, k)$-*regular* if all but at most $\varepsilon\binom{k}{2}$ pairs (V_i, V_j) are $(p; H, \varepsilon)$-regular, and, for all $i = 1, \ldots, k$, we have $\lfloor n/k \rfloor \leqslant |V_i| \leqslant \lceil n/k \rceil$. The following sparse version of Szemerédi's regularity lemma will be important in what follows (for general discussions on this lemma, see, *e.g.*, [12, 13, 14]).

Lemma 5.3. *For all $\varepsilon > 0$, $b \geqslant 1$, and all integers m and $r \geqslant 1$, there exist $\beta > 0$ and $K \geqslant m$ for which the following holds. For all p and all r-tuples of $(p; b, \beta)$-bounded graphs $(H_1, \ldots, H_r)$, with all the H_i on the same vertex set V with $|V| \geqslant m$, there exists a partition $V = V_1 \cup \cdots \cup V_k$ of V with $m \leqslant k \leqslant K$ that is $(p; H_i, \varepsilon, k)$-regular for all $1 \leqslant i \leqslant r$.*

5.2. Proof of Lemma 4.3

We start with two basic facts.

Fact 5.4. *Let G be a graph on $k \geqslant 1$ vertices and with at least $\gamma\binom{k}{2}$ edges, where $0 < \gamma \leqslant 1$. Then the number of vertices of degree at least $\gamma k/2$ in G is at least $\gamma k/2$.* □

Fact 5.5. *Suppose $k \geqslant 1$ and*

$$E(K_k) = R \cup B \cup Y \cup S \tag{5.6}$$

is a colouring of the edges of the complete graph K_k with colours red, blue, yellow, and sienna, where

$$|R|,\ |B| \geqslant \gamma\binom{k}{2}$$

and

$$|S| \leqslant \frac{1}{8}\gamma^2\binom{k}{2}. \tag{5.7}$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Then there is a vertex x in K_k and distinct colours $C_1, C_2 \in \{R, B, Y\}$ such that x is incident to at least $\gamma k/8$ edges of colour C_1 and at least $\gamma k/8$ edges of colour C_2.

Proof. For a colour C and vertex x, let $d(C, x)$ be the number of edges of colour C incident to x in colouring (5.6). For $C = R$ and B, let

$$U_C = \left\{ u \in V(K_k) : d(C, x) \geqslant \frac{1}{2}\gamma k \right\}. \tag{5.8}$$

By Fact 5.4, we have $|U_R|, |U_B| \geqslant \gamma k/2$. Clearly, we are done if $U_R \cap U_B \neq \emptyset$. Hence we suppose that $U_R \cap U_B = \emptyset$. Consider the $|U_R||U_B|$ edges in $E(U_R, U_B) = E_{K_k}(U_R, U_B)$. There is a colour $C \in \{R, B, Y\}$ for which we have

$$|E(U_R, U_B) \cap C| \geqslant \frac{1}{3}(|U_R||U_B| - |S|).$$

Without loss of generality, we may assume that $C \neq R$. By averaging, we may deduce that there is a vertex $x \in U_R$ for which we have

$$d(C, x) \geqslant \frac{|E(U_R, U_B) \cap C|}{|U_R|} \geqslant \frac{1}{3}\left(|U_B| - \frac{|S|}{|U_R|}\right) \geqslant \frac{1}{3}\left(\frac{1}{2}\gamma k - \frac{(1/16)\gamma^2 k^2}{\gamma k/2}\right) = \frac{1}{8}\gamma k,$$

and we are again done: x is incident to at least $\gamma k/2 \geqslant \gamma k/8$ edges of colour R and to at least $\gamma k/8$ edges of colour $C \neq R$. □

We are now ready to prove Lemma 4.3.

Proof of Lemma 4.3. Let $\varepsilon > 0$ be given. Since the smaller ε is, the stronger is the conclusion in Lemma 4.3, we may suppose that

$$\varepsilon \leqslant \frac{1}{30}. \tag{5.9}$$

Put

$$\delta = \min\left\{\varepsilon^2, \frac{1}{10^2}\lambda_0, \frac{1}{2 \times 10^4}c^4\right\} \quad \text{and} \quad m = \frac{16}{3\lambda_0^2}. \tag{5.10}$$

Let

$$\beta = \beta(\delta, m) \quad \text{and} \quad K = K(\delta, m) \tag{5.11}$$

be the constants given by Lemma 5.3, for the constants $\varepsilon = \delta$, $b = 2$, m, and $r = 3$ as above. We now let

$$\sigma = \frac{1}{K(\delta, m)}, \tag{5.12}$$

and claim that this choice for σ will do. We now proceed to prove this claim. Thus, let the graph G and the edge-colouring $E(G) = R \cup B \cup Y$ be as in the statement of Lemma 4.3.

The expected number of triangles in G is $(1/6 + o(1))(np)^3 = (1/6 + o(1))c^3 n^{3/2}$. Also, the expected number of edges $e = \{a, b\}$ together with two further vertices x and y ($x \neq y$)

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

with $x, y \in N(G;a) \cap N(G;b)$ (that is, the number of diamonds) is

$$\binom{n}{2}\binom{n-2}{2}p^5 \sim \left(\frac{1}{4}+o(1)\right)c^5 n^{3/2}.$$

Standard variance calculations, combined with Chebyshev's inequality, show that both quantities above are concentrated around their means, *i.e.*, a.a.s. the number of triangles is $(1/6+o(1))c^3n^{3/2}$, and the number of diamonds is a.a.s. $(1/4+o(1))c^5n^{3/2}$. Therefore, the number of triangles in G that are 'solitary', *i.e.*, that share no edge with any other triangle, is a.a.s.

$$(1+o(1))\left(\frac{1}{6}c^3n^{3/2}-\frac{1}{2}c^5n^{3/2}\right) \geqslant \frac{1}{7}c^3n^{3/2},$$

where we used that $c \leqslant 1/5$. We may hence suppose that G contains at least $c^3n^{3/2}/7$ solitary triangles.

By adjusting the notation, we may suppose that

$$|R| \geqslant |B| \geqslant |Y|.$$

Since we are supposing that the colouring (4.5) contains no monochromatic triangle, we may deduce that $|B \cup Y| \geqslant c^3n^{3/2}/7$; indeed, each solitary triangle must contain either a blue edge or a yellow edge, for otherwise it would be entirely red. In particular, we have

$$|R|,\ |B| \geqslant \frac{1}{14}c^3n^{3/2}. \tag{5.13}$$

By Chernoff's inequality, we may also suppose that, for each pair $U, W \subseteq V(G)$ with $U \cap W = \emptyset$ and $|U|, |W| \geqslant n/\log n$, we have

$$\frac{1}{2}|U||W|p \leqslant e_G(U,W) = |E_G(U,W)| \leqslant 2|U||W|p, \tag{5.14}$$

and

$$e_G(U) = |E(G[U])| \leqslant |U|^2 p. \tag{5.15}$$

Note that it follows from (5.14) that G is a.a.s. $(p;2,\beta)$-bounded for all $\beta > 0$.

Let us now apply Lemma 5.3 to obtain a partition $V(G) = V_1 \cup \cdots \cup V_k$ of $V(G)$ with $m \leqslant k \leqslant K$ that is $(p;F,\delta,k)$-regular for $F = R$, $F = B$, and $F = Y$. We now state and prove a claim that will be used in what follows.

Claim 5.6. *Let $1/2 > \lambda > \frac{2}{25}\lambda_0$. Suppose $F \subseteq E(G)$ is a subset of edges of $G = G(n,p)$, with $|F| \geqslant \lambda cn^{3/2}$. Let $V_1, \ldots, V_k$ be a $(p;F,\delta,k)$-regular partition of $V(G)$, where $m \leqslant k \leqslant K$. Suppose for simplicity that $|V_1| = \cdots = |V_k|$. Let Φ be the set of pairs $\{i,j\}$ $(1 \leqslant i < j \leqslant k)$ for which (V_i, V_j) is $(p;F,\delta)$-regular and*

$$e_F(V_i,V_j) \geqslant \frac{1}{2}\lambda p\left(\frac{n}{k}\right)^2. \tag{5.16}$$

Then

$$|\Phi| \geqslant \frac{5}{8}\lambda\binom{k}{2}. \tag{5.17}$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Proof. Let $\ell = x\binom{k}{2}$ be the number of pairs (V_i, V_j) satisfying (5.16), and recall that we are assuming that $|V_i| = n/k$ $(1 \leqslant i \leqslant k)$. Using that G satisfies (5.14) and (5.15) for all 'large' U and $W \subseteq V(G)$, we deduce that

$$\lambda c n^{3/2} \leqslant |F| \leqslant 2\ell \left(\frac{n}{k}\right)^2 p + \left(\binom{k}{2} - \ell\right) \frac{\lambda p}{2} \left(\frac{n}{k}\right)^2 + k\left(\frac{n}{k}\right)^2 p,$$

and so

$$\lambda \leqslant x + \frac{1}{4}(1-x)\lambda + \frac{1}{k}.$$

Solving the last inequality for x, we obtain $x \geqslant 3\lambda/4$, because $1/k \leqslant 1/m = 3\lambda^2/16$. Hence, at least

$$\frac{3}{4}\lambda\binom{k}{2} - \delta\binom{k}{2} \geqslant \frac{5}{8}\lambda\binom{k}{2}$$

pairs (V_i, V_j) are $(p; F, \delta)$-regular, each satisfying (5.16); that is, the set Φ satisfies (5.17), as required. □

For $F = R$, B, and Y, let Φ_F be the set of pairs $\{i, j\}$ $(1 \leqslant i < j \leqslant k)$ such that (V_i, V_j) is $(p; F, \delta)$-regular and

$$e_F(V_i, V_j) \geqslant \frac{1}{2}\left(\frac{1}{14}c^2\right) p \left(\frac{n}{k}\right)^2 = \frac{1}{28} c^2 p \left(\frac{n}{k}\right)^2. \tag{5.18}$$

We apply Claim 5.6 with $F = R$ and B and $\lambda_0 = c^2/14$ (see (5.13)). That claim tells us that

$$|\Phi_R|,\ |\Phi_B| \geqslant \frac{1}{25} c^2 \binom{k}{2}. \tag{5.19}$$

Since

$$3 \times \frac{1}{28} c^2 p \left(\frac{n}{k}\right)^2 \leqslant \frac{1}{2} p \left(\frac{n}{k}\right)^2,$$

the fact that (5.14) holds for all 'large' U and $W \subseteq V(G)$ implies that if a pair $\{i, j\}$ $(1 \leqslant i < j \leqslant k)$ is not a member of $\Phi_R \cup \Phi_B \cup \Phi_Y$, then (V_i, V_j) is *not* $(p; F, \delta)$-regular for some $F \in \{R, B, Y\}$. Put

$$\Phi_S = [k]^2 \setminus (\Phi_R \cup \Phi_B \cup \Phi_Y).$$

The observations above, the fact that $V_1, \ldots, V_k$ is a $(p; F, \delta, k)$-regular partition of $V(G)$ for $F \in \{R, B, Y\}$, and our choice of δ (see (5.10)) imply that

$$|\Phi_S| \leqslant 3\delta\binom{k}{2} \leqslant \frac{3}{2 \times 10^4} c^4 \binom{k}{2} < \frac{1}{8}\left(\frac{1}{25}c^2\right)^2 \binom{k}{2}. \tag{5.20}$$

We now apply Fact 5.5 to the following colouring of the edges of the complete graph K_k on $[k] = \{1, \ldots, k\}$:

$$E(K_k) = \Phi_R \cup \Phi_B \cup \Phi_Y \cup \Phi_S.$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

In view of (5.19) and (5.20), we may apply Fact 5.5 with $\gamma = c^2/25$. We then obtain $i_0 \in [k] = V(K_k)$ and two disjoint sets $\Gamma'_F \subseteq [k]$ ($F = R$ and B, say) with

$$|\Gamma'_F| \geqslant \frac{1}{8}\left(\frac{1}{25}c^2\right)k = \frac{1}{200}c^2k, \tag{5.21}$$

such that, for all $j \in \Gamma'_F$, we have that

$$(V_{i_0}, V_j) \text{ is } (p; F, \delta)\text{-regular} \tag{5.22}$$

and

$$e_F(V_{i_0}, V_j) \geqslant \frac{1}{28}c^2 p\left(\frac{n}{k}\right)^2. \tag{5.23}$$

We let $\Gamma_F \subseteq \Gamma'_F$ ($F = R$ and B) be an arbitrary subset of Γ'_F with $|\Gamma_F| = (c^2/200)k$. Moreover, let

$$U = V_{i_0}, \quad Z = \bigcup_{j\in\Gamma_R} V_j, \quad \text{and} \quad W = \bigcup_{j\in\Gamma_B} V_j,$$

and let

$$J = R[Z, U] \cup B[U, W]. \tag{5.24}$$

Claim 5.7. *We claim that J, Z, U, and W satisfy* (i), (ii) *and* (iii) *of Lemma* 4.3.

Proof. We start by noticing that

$$|Z| = |W| = \frac{1}{200}c^2k \times \frac{n}{k} = \frac{1}{200}c^2n = \sigma_0 n.$$

Moreover,

$$|U| = \frac{n}{k} \geqslant \frac{n}{k}(\delta, m) = \sigma n.$$

Property (ii) asserted in Lemma 4.3 is clear from the definition of J (see (5.24)). We are now left with proving (iii), that is, that (Z, U) and (U, W) are $(p, \lambda_0; J, \varepsilon)$-semi-regular. We deal with (Z, U), as the other case is analogous.

Let $U' \subseteq U$ and $Z' \subseteq Z$ be such that $|U'| \geqslant \varepsilon|U|$ and $|Z'| \geqslant \varepsilon|Z|$. We have to show that

$$\frac{e_J(U', Z')}{p|U'||Z'|} \geqslant \frac{1}{30}c^2 = \lambda_0. \tag{5.25}$$

Put $Z'_j = Z' \cap V_j$ ($j \in \Gamma_R$). Clearly, we have $e_J(U', Z') = \sum_{j\in\Gamma_R} e_J(U', Z'_j)$. Therefore, we have

$$\begin{aligned}\frac{e_J(U', Z')}{p|U'||Z'|} &= \sum_{j\in\Gamma_R} \frac{e_J(U', Z'_j)}{p|U'||Z'_j|} \times \frac{|U'||Z'_j|}{|U'||Z'|} \\ &= \left(\sum_{j\in\Gamma'_R} + \sum_{j\in\Gamma''_R}\right) \frac{e_J(U', Z'_j)}{p|U'||Z'_j|} \times \frac{|Z'_j|}{|Z'|},\end{aligned} \tag{5.26}$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

where $\Gamma'_R = \{j \in \Gamma_R : |Z'_j|/|V_j| < \delta\}$, and $\Gamma''_R = \Gamma_R \setminus \Gamma'_R$. We now claim that

$$\sum_{j \in \Gamma''_R} \frac{|Z'_j|}{|Z'|} \geqslant 1 - \varepsilon. \tag{5.27}$$

Indeed,

$$\sum_{j \in \Gamma'_R} |Z'_j| \leqslant \delta \sum_{j \in \Gamma'_R} |V_j| \leqslant \delta |Z|.$$

Moreover, $|Z'| \geqslant \varepsilon |Z|$, and hence

$$\sum_{j \in \Gamma'_R} |Z'_j| \leqslant \delta \left(\frac{1}{\varepsilon} |Z'| \right) \leqslant \varepsilon |Z'|, \tag{5.28}$$

as $\delta \leqslant \varepsilon^2$ (see (5.10)). It follows from (5.28) that (5.27) does indeed hold.

We now go back to (5.26). Clearly, the right-hand side of (5.26) is, by (5.22), (5.23), and (5.27), at least

$$\sum_{j \in \Gamma''_R} \frac{e_J(U', Z'_j)}{p|U'||Z'_j|} \times \frac{|Z'_j|}{|Z'|} \geqslant \left(\frac{1}{28} c^2 - \delta \right) \sum_{j \in \Gamma''_R} \frac{|Z'_j|}{|Z'|} \geqslant \left(\frac{1}{28} c^2 - \delta \right)(1 - \varepsilon) \geqslant \frac{1}{30} c^2 = \lambda_0.$$

We have thus shown that (5.25) does indeed hold, and so (Z, U) is $(p, \lambda_0; J, \varepsilon)$-semi-regular. This concludes the proof of Claim 5.7. □

Claim 5.7 completes the proof of Lemma 4.3. □

5.3. Proof of Lemma 4.7

The basic underlying idea in this proof comes from [15, Lemma 11] (*cf.* [12, Lemma 8.30]).

We start by letting arbitrary positive constants τ', and α be given. To prove Lemma 4.7, we need to define an appropriate constant $\varepsilon' > 0$. Put

$$\varepsilon' = \min \left\{ \frac{1}{16} (\alpha/3)^{96/\lambda_0 \tau' \sigma_0}, \frac{1}{4} \tau' \exp\{-\lambda_0 c^2/3\} \right\}. \tag{5.29}$$

We claim that this choice for ε' will do.

Recall that we wish to estimate the number of bipartite graphs H with bipartition (U^*, W), where $|W| = w = \sigma_0 n$ and $|U^*| = u^* \geqslant n/(\log n)^2$, belonging to the family

$$\mathscr{B}_1 = \mathscr{B}_1(U^*, W; F; \tau', \varepsilon'; n, T),$$

where

$$F \subseteq [U^*]^2, \qquad |F| = \tau' \binom{u^*}{2}. \tag{5.30}$$

In order to obtain this estimate, let us make the following observation about the graphs $H \in \mathscr{B}_1$.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Lemma 5.8. *Suppose $H \in \mathscr{B}_1$. Then H is a union of two edge-disjoint subgraphs H' and H'' with $|E(H')| = |E(H'')| = T/2$, and the pair (U^*, W) is both $(p, \lambda_0/3; H', \varepsilon')$-semi-regular and $(p, \lambda_0/3; H'', \varepsilon')$-semi-regular.*

Proof. It suffices to use the 'probabilistic method', taking a random partition. The result follows from standard exponential estimates for the tail of the hypergeometric distribution. We omit the details. □

For each graph $H \in \mathscr{B}_1$, we fix a decomposition $H = H' \cup H''$. By Fact 5.4, there exists a set $U^{**} \subseteq U^*$ such that

$$u^{**} = |U^{**}| \geqslant \tau' u^*/2 \tag{5.31}$$

and for all $v \in U^{**}$ we have $\deg_F(v) \geqslant \tau' u^*/2$. We now need to fix some notation. Let

$$U^{**} = \{v_i : 1 \leqslant i \leqslant u^{**} = |U^{**}|\},$$

and let

$$F_i = N(F; v_i), \qquad W^{(i)} = N(H''; v_i) \tag{5.32}$$

be the neighbourhoods of v_i in the graph F and H'', respectively. Because of our choice of U^{**}, we have

$$|F_i| = \deg_F(v_i) \geqslant \frac{1}{2}\tau' u^* \tag{5.33}$$

for all $1 \leqslant i \leqslant u^{**}$.

Let $d_i = |W^{(i)}|$. Then, by the $(p, \lambda_0/3; H'', \varepsilon')$-semi-regularity of (U^*, W), it follows from Lemma 5.1 that at least $(1 - \varepsilon')u^*$ vertices of U^* have at least $(\lambda_0/3)p|W|$ neighbours in W. Hence, for at least $u^{**} - \varepsilon' u^* > u^{**}/2$ indices i, we have $d_i \geqslant (\lambda_0/3)p|W|$ (note that $\varepsilon' < \tau'/4$). Thus, we may suppose that

$$\frac{1}{3}\lambda_0 p|W| \leqslant d_i \leqslant 2c\sqrt{n} \quad \text{for all} \quad 1 \leqslant i \leqslant u^{**}/2 \tag{5.34}$$

(recall that $\Delta(H) \leqslant 2c\sqrt{n}$). It will be convenient to set

$$W^{(i)} = \{w_1^{(i)}, \ldots, w_{d_i}^{(i)}\} \tag{5.35}$$

for all $1 \leqslant i \leqslant u^{**}/2$.

Let

$$\mu_0 = \exp\{-\lambda_0^2 c^2 \sigma_0/18\}. \tag{5.36}$$

We shall say that v_i $(1 \leqslant i \leqslant u^{**}/2)$ is *bad* if

$$\left|F_i \backslash N\left(H'; W^{(i)}\right)\right| \geqslant \mu_0 |F_i|.$$

We claim that more than $u^{**}/4$ vertices are bad. Suppose for a contradiction that the number of vertices v_i $(1 \leqslant i \leqslant u^{**}/2)$ that are *not* bad is at least $u^{**}/4$. Let v_i be one of them. Clearly, the number of edges of $F \cap \mathrm{Base}(H)[U^*]$ adjacent to v_i is at least

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

$|F_i \cap N(H'; W^{(i)})|$ (this is the set of H'-neighbours of the H''-neighbours of v_i which are also F-neighbours of v_i). Since v_i is not bad, we have

$$\left|F_i \cap N\left(H'; W^{(i)}\right)\right| \geqslant (1-\mu_0)|F_i| \geqslant \frac{1}{50}\lambda_0^2 c^2 \sigma_0 |F_i|, \tag{5.37}$$

where we used that $x = (1/18)\lambda_0^2 c^2 \sigma_0 \leqslant 1/2$, and hence $1 - e^{-x} \geqslant x/2$, say. However, this means that v_i contributes to $F \cap \mathrm{Base}(H)[U^*]$ at least as many pairs as in (5.37) (with some pairs counted twice, when all v_i are considered). Consequently, by (4.3), (5.30), (5.31) and (5.33),

$$|F \cap \mathrm{Base}(H)[U^*]| \geqslant \frac{1}{4}u^{**} \times \frac{1}{2}\left(\frac{1}{50}\lambda_0^2 c^2 \frac{1}{2}\tau' u^*\right) \geqslant (\tau')^2 \binom{u^*}{2} = \tau'|F|, \tag{5.38}$$

which is in contradiction to the fact that $H \in \mathscr{B}_1$ (see (4.10)).

Completion of the proof of Lemma 4.7. Suppose $H \in \mathscr{B}_1$ is given. In view of Lemma 5.8, we may decompose H as a union $H' \cup H''$, so that $|E(H')| = |E(H'')| = T/2$ and (U^*, W) is both $(p, \lambda_0/3; H', \varepsilon')$-semi-regular and $(p, \lambda_0/3; H'', \varepsilon')$-semi-regular.

We estimate the cardinality of

$$\mathscr{B}_1 = \mathscr{B}_1(U^*, W; F; \sigma_0, \lambda_0, \tau', \varepsilon'; n, T)$$

as follows. We 'generate' the members $H \in \mathscr{B}_1$ in two rounds: first H', then H''. We do not require that H' and H'' above be disjoint. Since we are after an upper bound for $|\mathscr{B}_1|$, this is justified.

The number of choices of H' is bounded crudely by $\binom{wu^*}{T/2}$. With fixed H', for every H'' such that $H' \cup H'' \in \mathscr{B}_1$, there exist $u^{**}/4$ bad vertices $v_i \in U^*$ with $d_i = |N(H'; v_i)| \geqslant (\lambda_0/3)pw$. For each one of them we apply Corollary 5.2 to H' with $d = d_i$, $U_0 = F_i$, $W_0 = W^{(i)}$, $\varepsilon = \varepsilon'$, $\lambda = \lambda_0/3$ and $\mu = \exp\{-\lambda_0 pd/6\}$.

We have $|F_i \backslash N(H'; W^{(i)})| > \mu_0|F_i| \geqslant \mu|F_i|$, and thus (5.1) holds. We also have

$$|F_i| \geqslant \tau' u^*/2 > \varepsilon' \exp\{\lambda_0 c^2/3\} u^* \geqslant (\varepsilon'/\mu)u^*.$$

Hence, by Corollary 5.2 the number of choices of the neighbourhood $N(H''; v_i)$ is at most

$$(\varepsilon')^{d/2}\binom{d}{d/2} w^d/d! < (\varepsilon')^{d/2} 2^d \binom{w}{d}. \tag{5.39}$$

Thus, an upper bound on the number of choices of H'' may be obtained by bounding first the number of choices of the $u^{**}/4$ bad vertices by 2^{u^*}, and then summing the products

$$(2\sqrt{\varepsilon'})^D \prod \left\{\binom{w}{d(v)} : v \in U^*\right\} \tag{5.40}$$

over all degree sequences $\mathbf{d} = \{d(v) : v \in U^*\}$, where $D = (1/12)\lambda_0 p u^{**} w$ is a lower bound on the sum of the degrees of all bad vertices (*cf.* (5.34)). Indeed, with a fixed set of bad vertices U^{***}, and fixed degree sequence $\mathbf{d}$, by (5.39) there are at most

$$\prod_{v \in U^{***}} (\varepsilon')^{d(v)/2} 2^{d(v)} \binom{w}{d(v)} \times \prod_{v \in U^* \backslash U^{***}} \binom{w}{d(v)}$$

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

choices of the neighbourhoods which together create H''. To estimate the quantity in (5.40), we observe that, because $\Delta(H) \leqslant 2c\sqrt{n}$, we have $T \leqslant 2c\sqrt{n}|U^*| = 2cu^*\sqrt{n}$. Moreover, we recall that $u^{**} \geqslant \tau' u^*/2$, $p = c/\sqrt{n}$, and $w = \sigma_0 n$, and hence we have

$$\frac{D}{T} \geqslant \frac{(1/12)\lambda_0 p u^{**} w}{2c\sqrt{n}u^*} \geqslant \frac{1}{48}\lambda_0 \tau' \sigma_0.$$

Therefore, given H', there are at most

$$2^{u^*}(2\sqrt{\varepsilon'})^{(\lambda_0\tau'\sigma_0/48)T} \prod \left\{ \binom{w}{d(v)} : v \in U^* \right\} \tag{5.41}$$

choices for H''. We now sum (5.41) over all choices of $\mathbf{d} = \{d(v): v \in U^*\}$. Recalling that $\mathbf{d}$ is a partition of $T/2$, and that

$$T \geqslant \lambda_0 p u^* w = \lambda_0 c \sigma_0 u^* \sqrt{n} \gg u^* \tag{5.42}$$

(*cf.* (4.17) and (4.18)), we see that this sum is at most

$$(4\sqrt{\varepsilon'})^{(\lambda_0\tau'\sigma_0/48)T} \binom{wu^*}{T/2}. \tag{5.43}$$

To obtain the desired bound on $\mathscr{B}_1$ we have to multiply the above by $\binom{wu^*}{T/2}$, the number of choices for H'. As, crudely,

$$\binom{wu^*}{T/2}^2 < 3^T \binom{wu^*}{T},$$

Lemma 4.7 is proved by our choice of ε'. □

6. Concluding remarks

We close by mentioning a few remarks and open problems.

The online game for more colours. We hope to address the online game when the Painter has $r > 2$ colours in the near future.

General graphs. It would be most interesting to generalize our results for general graphs H, that is, for avoidance games in which the Painter has to avoid monochromatic copies of a given graph H. However, our methods are deeply based on the simple structure of the triangles and hence a completely new approach may be needed.

Our results above suggest that, in the two-round game for arbitrary graphs H, the first interesting case is the one in which $r = 2$ and N_0 is around the threshold for the corresponding Ramsey property, that is, $N_0 = \Theta(n^{2-1/d_2(H)})$, where

$$d_2(H) = \max\{(|E(F)| - 1)/(|V(F)| - 2): F \subseteq H,\ |V(F)| > 2\}$$

(see [19]).

Deterministic games. In the deterministic version of our online game, say for $r = 2$ and $H = K_k$, the edges are generated not randomly, but by an adversary. Hence, this is a two-person game with the payoff to the Painter equal to the number of coloured edges until she is forced to form a monochromatic clique K_k.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

Define the *online Ramsey number* $\bar{R}(k)$ as the value of this game. The question is whether it is true that $\bar{R}(k) = o(R(k)^2)$ as $k \to \infty$, where $R(k)$ is the classical Ramsey number. Some preliminary results in this direction have been proved by Kurek and Ruciński [16]. In particular, $\bar{R}(3) = 8$ and $\bar{R}(k,l) = o(R(k,l)^2)$ as $k \to \infty$ while $l \geqslant 3$ is fixed, where $R(k,l)$ and $\bar{R}(k,l)$ are the corresponding off-diagonal numbers.

Finally, we mention that other games inspired by Ramsey's theorem have been investigated in several forms. The reader is referred to, for instance, Beck [2, 3, 4, 5], Beck and Csirmaz [1], Erdős and Selfridge [7], and Seress [20].

Acknowledgements

We thank Svante Janson for his interest and suggestions. We also thank the anonymous referee for careful reading and many useful suggestions. Part of this work was done when some of the authors were at the meeting *Combinatorics, Probability and Computing*, at the Mathematisches Forschungsinstitut Oberwolfach.

References

[1] Beck, J. and Csirmaz, L. (1982) Variations on a game. *J. Combin. Theory Ser. A* **33** 297–315.
[2] Beck, J. (1981) van der Waerden and Ramsey type games. *Combinatorica* **1** 103–116.
[3] Beck, J. (1985) Random graphs and positional games on the complete graph. In *Random Graphs '83* (Poznań, 1983), North-Holland, Amsterdam, pp. 7–13.
[4] Beck, J. (1994) Deterministic graph games and a probabilistic intuition. *Combin. Probab. Comput.* **3** 13–26.
[5] Beck, J. (1996) Foundations of positional games. In *Proceedings of the Seventh International Conference on Random Structures and Algorithms* (Atlanta, GA, 1995), Vol. 9, pp. 15–47.
[6] Bollobás, B. (1985) *Random Graphs*, Academic Press, London.
[7] Erdős, P. and Selfridge, J. L. (1973) On a combinatorial game. *J. Combin. Theory Ser. A* **14** 298–301.
[8] Frankl, P. and Rödl, V. (1986) Large triangle-free subgraphs in graphs without K_4. *Graphs Combin.* **2** 135–144.
[9] Friedgut, E. (1999) Sharp thresholds of graph properties, and the k-sat problem. *J. Amer. Math. Soc.* **12** 1017–1054.
[10] Friedgut, E., Rödl, V., Ruciński, A. and Tetali, P. (2002) A sharp threshold for random graphs with a monochromatic triangle in every edge coloring. Submitted.
[11] Janson, S., Łuczak, T. and Ruciński, A. (1990) An exponential bound for the probability of nonexistence of a specified subgraph in a random graph. In *Random graphs '87* (Poznań, 1987), Wiley, Chichester, pp. 73–87.
[12] Janson, S., Łuczak, T. and Ruciński, A. (2000) *Random Graphs*, Wiley-Interscience, New York.
[13] Kohayakawa, Y. (1997) Szemerédi's regularity lemma for sparse graphs. In *Foundations of Computational Mathematics* (Rio de Janeiro, 1997), Springer, Berlin, pp. 216–230.
[14] Kohayakawa, Y. and Rödl, V. (2003) Szemerédi's regularity lemma and quasi-randomness. In *Recent Advances in Algorithms and Combinatorics*, Springer, pp. 289–351.
[15] Kohayakawa, Y., Łuczak, T. and Rödl, V. (1996) Arithmetic progressions of length three in subsets of a random set. *Acta Arith.* **75** 133–163.
[16] Kurek, A. (2002) The sparsest Ramsey graphs are complete graphs. PhD thesis 1998.
[17] Łuczak, T., Ruciński, A. and Voigt, B. (1992) Ramsey properties of random graphs. *J. Combin. Theory Ser. B* **56** 55–68.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press

[18] Rödl, V. and Ruciński, A. (1994) Random graphs with monochromatic triangles in every edge coloring. *Random Struct. Alg.* **5** 253–270.
[19] Rödl, V. and Ruciński, A. (1995) Threshold functions for Ramsey properties. *J. Amer. Math. Soc.* **8** 917–942.
[20] Seress, Á. (1992) On Hajnal's triangle-free game. *Graphs Combin.* **8** 75–79.
[21] Szemerédi, E. (1978) Regular partitions of graphs. In *Problèmes Combinatoires et Théorie des Graphes* (Colloq. Internat. CNRS, Univ. Orsay, Orsay, 1976), CNRS, Paris, pp. 399–401.

https://doi.org/10.1017/S0963548303005881 Published online by Cambridge University Press