RAIRO-Theor. Inf. Appl. 45 (2011) 301–309
DOI: 10.1051/ita/2011118

Available online at:
www.rairo-ita.org

THE FIBONACCI AUTOMORPHISM OF FREE BURNSIDE GROUPS

Ashot S. Pahlevanyan[1]

Abstract. We prove that the Fibonacci morphism is an automorphism of infinite order of free Burnside groups for all odd $n \geq 665$ and even $n = 16k \geq 8000$.

Mathematics Subject Classification. 20F28, 20E36, 20F50, 20M05.

1. Introduction

The question of study of automorphisms of free Burnside groups was stated by Ol'shanskii in the Kourovka Notebook [7]. The first results were obtained by Cherepanov in [4,5] and by Atabekyan in [2,3]. In paper [4] it was proved that the Fibonacci morphism is an automorphism of infinite order of free Burnside groups for all odd $n > 10^{10}$ and even $n = 16k \geq 8000$.

This paper shows that the bound of odd n can be decreased from $n > 10^{10}$ to $n \geq 665$.

Consider an automorphism $\varphi : F_2 \to F_2$ of the absolutely free group F_2 of rank two with free generators $\{a, b\}$, given on generators by formulae

$$\varphi : a \mapsto b, \quad \varphi : b \mapsto ab.$$

This automorphism is called after Fibonacci since the lengths of words $\varphi^k(a)$ are equal to corresponding members of the numerical Fibonacci sequence. If we consider the sequence of mirror copies of words $\varphi^k(a)$, we obtain the iterations of the following morphism

$$h : a \mapsto b, \quad \varphi : b \mapsto ba.$$

Keywords and phrases. Free periodic groups, Burnside groups, group automorphisms, Fibonacci morphism, Fibonacci sequence, Fibonacci word, golden ratio.

[1] Faculty of Mathematics and Mechanics, Yerevan State University, A. Manoogian st. 1, Yerevan-0025, Armenia. apahlevanyan@ysu.am

Article published by EDP Sciences

© EDP Sciences 2011

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

This morphism is also called after Fibonacci. All the statements of this paper that we prove for the first morphism hold for the second morphism either.

Automorphism φ naturally induces an automorphism of free the Burnside group $B(2,n)$, which we denote by the same letter φ. Let us remember that a free Burnside group $B(2,n)$ is the quotient $F_2/{F_2}^n$, where ${F_2}^n$ is the subgroup generated by all possible n-powers of elements of F_2. Obviously the group $B(2,n)$ has a presentation $B(2,n) = \langle a_1, a_2 | A^n = 1, \ \textit{for all words}\, A = A(a_1, a_2)\rangle$.

Theorem 1.1. *For arbitrary odd $n \geq 665$ and arbitrary even $n = 16k \geq 8000$ the Fibonacci automorphism φ has infinite order in the group $Aut(B(2,n))$.*

Theorem 1.1 strengthens the similar result of paper [4], decreasing the bound of odd n to $n \geq 665$.

To prove Theorem 1.1 we prove the following result that is individually interesting.

Proposition 1.2. *For any natural k no forth power of a non-empty word occurs in a cyclic word $\varphi^k(a)$.*

As usual, by a cyclic word we mean a word written on a circle without fixing its start. Proposition 1.2 strengthens one of the results of paper [6] by Karhumäki, where a similar statement is proved without the assumption that the word $\varphi^k(a)$ is cyclic. Our proof of Proposition 1.2 does not depend on paper [6] by Karhumäki.

Proposition 1.2 also strengthens the Lemma 1.3 of paper [4] by Cherepanov, according to which no 24th power of a non-empty word occurs in a cyclic Fibonacci sequence. Bearing on paper [6] by Karhumäki in [9] Mignosi and Pirillo proved the following interesting result:

Proposition 1.3. *The Fibonacci infinite word contains no fractional power with an exponent grater than $2 + ((\sqrt{5}+1)/2)$ and, for any real number $\varepsilon > 0$, it contains a fractional power with an exponent grater than $2 + ((\sqrt{5}+1)/2) - \varepsilon$.*

Theorem 1.1 implies

Corollary 1.4. *For arbitrary odd $n \geq 665$ the quotient group*

$$Aut(B(2,n))/Inn(B(2,n))$$

is infinite.

Proof. According to the famous theorem of S. I. Adian (see Thm. VI.3.4 of [1]) the center of $B(m,n)$ is trivial for $n \geq 665$ and $m > 1$. Therefore $Inn(B(2,n))$ is isomorphic to $B(2,n)$. Since any inner automorphism of the group $B(2,n)$ has a finite order, from Theorem 1.1 it follows that for any natural number l each automorphism φ^l is not inner. Hence the quotient $Aut(B(2,n))/Inn(B(2,n))$ is infinite. □

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

2. The Proof of Proposition 1.2

We adhere to the following notions and notations of monograph [1].

Definition 2.1. The word A is called *primitive* if it cannot be presented in a form D^r for $r > 1$.

We say that the word E occurs in a word X, if there exist words R and Q such that $X = REQ$ holds. If the word R (word Q) is empty, then E is a prefix (suffix) of X. If X is a word over the alphabet that does not contain the letter $*$ and $X = REQ$, the word $R * E * Q$ is called an occurrence of word E in a word X. E is called a base of the occurrence $R * E * Q$. For a given word X we denote by $\overline{X}$ the cyclic word generated by X, that is the word X written on a circle without fixing its start. For a given word X by $\partial(X)$ we denote the length of X, that is the number of its letters over the alphabet $\{a, b\}$. For the equality by definition of two words or two occurrences in a same word we use the symbol $\rightleftharpoons$.

Consider an automorphism $\varphi : a \mapsto b, \varphi : b \mapsto ab$ of the group $B(2, n)$. Let us first write out a few images

$$\varphi^k(a) : a \mapsto b \mapsto ab \mapsto bab \mapsto \underbrace{ab}\,\underbrace{bab} \mapsto \underbrace{bab}\,\underbrace{abbab} \mapsto \underbrace{abbab}\,\underbrace{babbabb}.$$

Denote

$$X_0 \rightleftharpoons \varphi^0(a) = a,\ X_k \rightleftharpoons \varphi^k(a),\ k = 1, 2, \ldots$$

Since $X_{k+1} = X_{k-1} \cdot X_k$, the lengths of words of the sequence X_k, $k = 1, 2, \ldots$ form a Fibonacci sequence. Let us denote

$$A \rightleftharpoons X_k, B \rightleftharpoons X_{k-1}, C \rightleftharpoons X_{k-2}, D \rightleftharpoons X_{k-3},$$

$$E \rightleftharpoons X_{k-4}, F \rightleftharpoons X_{k-5}, G \rightleftharpoons X_{k-6}, H \rightleftharpoons X_{k-7}.$$

Then $A = CDC, B = DC, C = ED$ and $X_{k+1} = BA = DCCDC$.

Let us recall the following statements from [1], that we often refer to.

Lemma 2.2 (see Lem. I.2.2 in [1])**.** *If $AB = BA$, then there exists a word D, such that $A = D^t, B = D^r$, for some $t, r \geq 0$.*

Lemma 2.3 (see Lem. I.2.9 in [1])**.** *Suppose $A^t A_1 = B^r B_1$, where $\partial(A^t A_1) \geq \partial(AB)$, A_1 is a suffix of A, B_1 - a prefix of B. If A is a primitive word, then for some k, $B = A^k$ holds.*

Lemma 2.4 (see Lem. IV.2.16 in [1])**.** *If no elementary α-power of rank 1 occurs in a word X, then $X \overset{\alpha}{\sim} Y \Rightarrow X = Y$.*

Definition 2.5 (see Def. I.4.34 in [1])**.** Suppose $\mathcal{A} = \bigcup_{i=1}^{\infty} \mathcal{A}_i$, where

$$X \in \mathcal{A} \Leftrightarrow X \in \mathcal{R}_{\alpha-1} \& Norm(\alpha, X, 9) = \varnothing.$$

Elements of the set $\mathcal{A}$ are called *absolutely reduced*.

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

From Propositions 1.2 and 1.3 follows

Corollary 2.6. *For any natural k the cyclic word $\varphi^k(a)$ contains no fractional power $2+((\sqrt{5}+1)/2)$ of a non-empty word.*

Proof. Let us prove it by induction on k. The base of induction is obvious. Suppose the statement is true for all natural $l \leq k$ and prove it for $k+1$. Since all cyclic shifts of the word $X_{k+1} = DCCDC$ occur in a Fibonacci word $X_{k+3} = DCCDCCDCDCCDC$ that contains no $2+((\sqrt{5}+1)/2)$ fractional power according to proposition 1.3, the word $\overline{X_{k+1}}$ contains no fractional $2+((\sqrt{5}+1)/2)$ power of a non-empty word either. □

Lemma 2.7. *None of the words $\overline{X_k}$ is a proper power, that is $\overline{X_k} \neq Z^t, t \geq 2$, for $k = 1, 2, \ldots$*

Proof. Since the cyclic shift of a proper power is itself a proper power, it is enough to prove that X_k is not a proper power. We prove this by induction on k. For $k \leq 5$ the proof is obvious. Suppose that the lemma is proved for all numbers $l \leq k$, and prove it for $k+1$. We can assume that the word Z is primitive. Let $X_{k+1} = DCCDC = Z^t, t \geq 2$. Then $CDCDC = (CD)^2C = {Z_1}^t$ for some cyclic shift Z_1 of Z. Since $t \geq 2$, $\partial(Z_1) + \partial(CD) < \partial(CDCDC)$ and by Lemma 2.3 we obtain $\overline{B} = CD = {Z_1}^p$. Therefore $B = Z^p$ and $A = Z^q$, $p, q \geq 1$, $p \neq q$. This contradicts the inductive assumption. □

Lemma 2.8. *If $\overline{X_k}$ contains a power Z^t then $t < 4$.*

Proof. The proof is by induction on k. For $k \leq 5$ the proof is obvious since the word $\overline{X_5} = \overline{bababbab}$ does not contain Z^4. For $k = 6$ we have the word $\overline{X_6} = \overline{abbabbababbab}$ that does not contain a subword Z^4 with $\partial(Z) \leq 3$. For $k = 7$ no word Z^4 with $\partial(Z) \leq 3$ occurs in $\overline{X_7} = \overline{bababbababbabbababbab}$. The word $\overline{X_7}$ does not contain Z^4 with $\partial(Z) = 4, 5$ either. Let $k \geq 7$. Suppose the statement is proved for all $l \leq k$ and prove it for $k+1$. Let Z^4 occur in $\overline{X_{k+1}} = \overline{DCCDC}$. Z^4 does not occur in a word $\overline{A} = \overline{CDC}$ by inductive assumption. Since D is a suffix of C, any subword of word $\overline{DCCDC}$ of length three over the alphabet $\{C, D\}$ occurs in $\overline{CDC}$. Therefore Z^4 does not occur in a subword of word $\overline{DCCDC}$ of length three over the alphabet $\{C, D\}$.

Let us prove that Z^4 does not occur in a subword of $\overline{DCCDC}$ of length four over the alphabet $\{C, D\}$ either. Let us write out subwords of length four of word $\overline{DCCDC}$ over the alphabet $\{C, D\}$:

$$DCCD, CCDC, CDCD, DCDC, CDCC.$$

Since the word $DCDC$ is a suffix of $CCDC$, it is enough to consider the case of the occurrence of Z^4 in words $DCCD, CCDC, CDCC, CDCD$.

I. Suppose Z^4 occurs in $DCCD$. Then it contains the base CC of the occurrence $D * CC * D$. We have obvious inequalities $2\partial(C) < 4\partial(Z) < 2\partial(D) + 2\partial(C)$. Therefore $\partial(Z) < \partial(C)$ and $\partial(Z) + \partial(C) < 2\partial(C)$. By Lemma 2.3 we obtain

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

$C = Z^p, p \geq 1$. According to Lemma 2.7 we have $p \not\geq 2$, and using $\partial(D) < \partial(C) = \partial(Z)$ we obtain $p \neq 1$.

II. Suppose Z^4 occurs in $CDCC = EDDEDED$. Then it contains the base $DED = DC$ of the occurrence $ED * DED * ED$. We have $\partial(Z) < \partial(C)$. Let us consider the following cases:

(1) If Z^4 contains the suffix E of the base of the occurrence $ED * DEDE * D$, then from $\partial(Z) < \partial(D) + \partial(E) = \partial(C)$ follows $(DE)^2 = {Z_1}^{t_1} Z_1', t_1 \geq 2$ for some cyclic shift Z_1 of Z. According to Lemma 2.3 we obtain $DE = {Z_1}^p, p \geq 1$. Hence $ED = C = {Z_2}^p$. By Lemma 2.7 we have $p \not\geq 2$ and by $\partial(Z) < \partial(C)$ the inequality $p \neq 1$ holds.

Thus, we can assume that Z^4 is contained in the base $EDDEDE$ of the occurrence $*EDDEDE * D$ in a word $CDCC$ and, at the same time, contains the base of the occurrence $ED * DED * ED$.

(2) If Z^4 contains the base $DDED$ of the occurrence $E * DDED * ED$, then from equality $DDED = FEDED = FCC$, where F is a suffix of C, and inequality $\partial(Z) < \partial(C)$ follows that ${Z_1}' {Z_1}^{t_1} = FC^2, t_1 \geq 2$ for some cyclic shift Z_1 of Z. Therefore $C = ED = {Z_2}^p$, for some cyclic shift Z_2 of Z. Since $\partial(CDCC) < 4\partial(C)$, the case $p = 1$ is impossible, and $p > 1$. This contradicts the Lemma 2.7.

(3) Thus, one can assume that Z^4 occurs in the base $DDEDE$ of occurrence $E * DDEDE * D$ and at the same time does not contain the prefix D and suffix E of that base. Then Z^4 occurs in $\overline{A} = \overline{EDDED}$. This contradicts the inductive assumption.

III. Now let Z^4 occur in $CDCD = EDDEDD$. Then it contains the base $DED = DC$ of the occurrence $ED * DED * D$. First note that $Z^t \neq CDCD$ holds, since in the contrary case $\partial(Z) < \partial(CD)$, hence $CD = Z^p, p \geq 2$. Thus, $B = DC = {Z_1}^p$ for some cyclic shift Z_1 of Z. This is a contradiction to inductive assumption. According to the case above the word Z^4 does not occur in the base $DDED$ of the occurrence $E * DDED * D$ since that base is equal to the base of the occurrence $E * DDED * C$ in an already considered word $CDCC$. Let us consider the following cases:

(1) If Z^4 contains the base of the occurrence $EF * EFEEFE * FE$ in a word $CDCD$ then $EFEEFE = (C)^2 = {Z_1}^{t_1} Z_1'$ and in view of $\partial(Z_1) < \partial(C)$ we have $C = {Z_1}^p$. This contradicts the Lemma 2.7 for $p \geq 2$ and the inequality $\partial(D) < \partial(C)$ for $p = 1$.

(2) Thus Z^4 occurs in the base $FEEFEFE$ of the occurrence $EFE*$ $FEEFEFE*$ in a word $CDCD$ and contains the base of the occurrence $EFE * FEEFE * FE$. Then $\partial(Z) < \partial(E) + \partial(F)$ holds. If Z^4 contains the base of the occurrence $EFE * FEEFEF * E$ then the word $E(EF)^2 = GFEFEF$ is periodic with period EF and $E(EF)^2 = {Z_1}' {Z_1}^{t_1}$. According to Lemma 2.3 we have $EF = {Z_1}^p$ and $FE = {Z_2}^p = D$. In view of Lemma 2.7 $p \not\geq 2$ holds and by $\partial(Z) < \partial(E) + \partial(F)$ we have $p \neq 1$.

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

(3) It remains to consider the case when Z^4 strictly occurs in the base of the occurrence $EF * EFEEFEF * E$ and at the same time does not contain the prefix E and suffix F of that base. Then $\partial(Z) < \partial(FE)$ and $EEFE = G(FE)^2 = {Z_1}'{Z_1}^{t_1}$ where $t_1 \geq 2$. According to Lemma 2.3 we have $D = FE = {Z_2}^p$ for some cyclic shift Z_2 of Z, in spite of Lemma 2.7.

IV. Now suppose Z^t occurs in a word $CCDC$. Then it contains the base $EDD = CD$ of the occurrence $ED * EDD * ED$ in a word $CCDC$. Since $\partial(D) < \partial(C)$ we have $\partial(Z) < \partial(C)$. Therefore ${Z_1}'{Z_1}^{t_1} = EDD = CD$ for some cyclic shift Z_1 of Z, where $t_1 \geq 2$, E is a suffix of D and $\partial(Z_1) < \partial(D) + \partial(E)$. According to Lemma 2.3 we obtain that $D = {Z_1}^p, p \geq 1$. By Lemma 2.7 we have $p = 1$ and $Z_1 = D$. The base of the occurrence $ED * EDD * ED$ in a word $CCDC$ is non-continuable to the right relative to period $Z_1 = D$ since the first letters of words D and E are different by definition of words X_k. Then Z^t occurs in a word $\overline{A} = \overline{CCD} = \overline{EDEDD}$ that contradicts the inductive assumption.

Thus we have proved that no word of form Z^4 occurs in a subword of length four of cyclic word $\overline{DCCDC}$ over the alphabet $\{C, D\}$. Let us now prove that it does not occur in a cyclic word $\overline{DCCDC}$ either. Assuming the contrary we obtain that Z^t contains one of the words CCD, CDC, DCC and DCD. In view of obvious inequalities $2\partial(C) < 2\partial(D) + \partial(C)$ and $\partial(C) + \partial(D) < 2\partial(C) + \partial(D)$ we obtain that more than half of the word Z^4 occurs in one of the following words CCD, CDC, DCC, DCD.

(1) Let Z^4 contain the base of the occurrence

$$D * CCD * C = D * EHGGFEFEFE * C$$

in a word $DCCDC$. Since $\partial(D) + \partial(EH) + \partial(C) \leq \partial(GGFEFEFE)$, we have $GG(FE)^3 = {Z_1}'{Z_1}^{t_1}$ for some cyclic shift Z_1 of Z, where $t_1 \geq 2$. Using the Lemma 2.3 we obtain that $D = FE = {Z_2}^p$ for some cyclic shift Z_1 of Z. By Lemma 2.7 we have $p = 1$ and $D = Z_2$. Since the first letters of C and D are different, the word $D * EHGGFEFEFE * C$ is non-continuable to the right relative to period $D = Z_2$. Then D^4 is a suffix of the occurrence $*DCCD*C = *DEFGFEFEFE*C$. Therefore FG is a suffix of the word $D = FE = FGF$, hence $FG = GF$. By Lemma 2.2 we get $E = T^p$ for some word T and $p \geq 2$. This contradicts the Lemma 2.7.

(2) Suppose Z^4 contains the base of the occurrence $C * DCC * D$ in a cyclic word $CDCCD$. Since D is a suffix of C, we have $DCC = {Z_1}'{Z_1}^{t_1}$ for some cyclic shift Z_1 of Z, and, obviously, $t_1 \geq 2$ holds. Then, according to Lemma 2.3, the word $C = {Z_2}^p$ is a proper power. By Lemma 2.7 we have $p = 1$ and $C = Z_2$. But since C is non-continuable to the right and to the left side one cannot count the word C^4 because of the inequality $\partial(D) < \partial(C)$, we obtain a contradiction.

(3) Let Z^4 contain the base of the occurrence

$$C * CDC * D = EFE * GHGF(EFE)^2 * FE$$

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

or

$$D*CDC*C = FE*GHGF(EFE)^2*EFE$$

in a cyclic word $CCDCD$ or $DCDCC$ respectively. One has $E = GF = GHG$. In view of the obvious inequality

$$\partial(F(EFE)^2) > \partial(EFE) + \partial(FE) + \partial(E)$$

by Lemma 2.3 and Lemma 2.7 we obtain $C = EFE = Z_2$. The suffix C of the base of the occurrence $C*CDC*D$ is not continuable to the right relative to period C because the first letters of words C and D are different. In view of the inequality $\partial(D) < \partial(C)$ the base of the occurrence $*CCDC*D$ does not end with word C^4. Now let Z^4 contain the base of the occurrence $D*CDC*C$. Consider the maximal power of the word C that occurs in a word $DCDCC$, where one C of that power coincides with the suffix C of the base of the occurrence $D*CDC*C$. We can continue the occurrence $D*CDC*C$ to the right relative to period $C = Z_2$. Now let us count from right to left the maximal power of C that occurs in a word $DCDCC$. We have the equalities

$$DCDCC = FEEFEFEEFEEFE = FEEFCCC.$$

It is obvious that the equality $EEF = EFE = C$ has to hold, and therefore $EF = FE$. Then $D = FE = T^p, p \geq 2$ that contradicts the Lemma 2.7.

(4) Finally suppose Z^4 contains the base of the occurrence

$$C*DCD*C = EF*EDCD*C$$

in a cyclic word $CDCDC$. Let us repeat the reasoning of case one. Having changed only the occurrence $D*CCD*C$ by the occurrence $EF*EDCD*C = EF*CCD*C$ in a word $CDCDC$, we can assume, that Z^4 does not contain the base of the occurrence $EF*CCD*C$. Therefore Z^4 occurs in a base of the occurrence $EF*EDCDC*$ and at the same time contains the base of the occurrence

$$EFE*DCD*C = EFE*GFEDD*C = EFE*GDDD*ED.$$

We have the inequality $2\partial(D) + \partial(E) \leq \partial(GD^3)$. Then $GD^3 = {Z_1}'{Z_1}^{t_1}$ for some cyclic shift Z_1 of Z, and $t_1 \geq 2$. Hence $D = {Z_2}^p$. By Lemma 2.7 we have $p = 1$ and $D = Z_2$. Since the first letters of words C and D are different, the occurrence $EFE*GFEDD*C$ is not continuable to the right relative to period D. If D^4 is a suffix of the base of occurrence $*CDCD*C = *CFGFEDD*C$ then from right to left we read D^2, $D = FE$, and the equality $FG = GF = E$ must hold. Therefore $E = T^p, p \geq 2$ that contradicts the Lemma 2.7. Thus we proved that

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

no word of form $Z^t, t \geq 4$ can occur in a word $\overline{X_{k+1}} = \overline{DCCDC}$. The Lemma is proved completely. □

It remains to note that the Lemma 2.8 is a reformulation of Proposition 1.2.

3. Proof of the main result

Now turn to the proof of Theorem 1.1.

Proof. Suppose that φ has a finite order in $Aut(B(2,n))$, that is $\varphi^k = id$. Then, particularly $\varphi^k(a) = a$, that is the word $X_k = a$. Therefore $a^{-1}X_k$ is equal to the empty word in $B(2,n)$. Consider two possible cases:

(1) If X_k starts with the letter b, then $a^{-1}X_k$ is not reducible and obviously contains no forth power of a non-empty word by Proposition 1.2 since all letters in X_k are positive.
(2) If X_k starts with the letter a, then we reduce $a^{-1}a$ and the result $X_k{}'$ contains no fourth power by Lemma 1.2. Hence, by definition 2.5 the irreducible word $a^{-1}X_k$ is absolutely reduced for odd $n \geq 665$ and according to Lemma 2.4 it cannot be equal to the empty word in $B(2,n)$. This proves the Theorem 1.1 for odd n.

To prove the Theorem 1.1 for even $n = 16k \geq 8000$ (as in [4]) we use Theorem 2(i) of [8] according to which if a non-empty freely non-reducible word X is equal to one in $B(m,n)$, then X contains a non-empty subword of the form $A^{(n/2)-1240}$. Again by Proposition 1.2 the word $a^{-1}X_k$ contains no subword of the form $A^{n/2-1240}$, hence it cannot be equal to the empty word in $B(2,n)$. □

Appendix

The author thanks the Referee for suggesting the following much shorter proof of Proposition 1.2 using some well-known properties of Fibonacci words.

Proof. Let h denote the Fibonacci morphism given by

$$h : a \mapsto b, \quad h : b \mapsto ba.$$

It is well-known that the reversal of $h(b)$ is a conjugate of $h(b)$ for all $k \geq 0$. Thus the square of the reversal of $h^{n-1}(b)$ is a factor of the cube $h^{n-1}(b)^3$, which is a factor of the Fibonacci infinite word $\lim_{k\to\infty} h^k(b)$ for all $n \geq 3$. Therefore the square of $h^{n-1}(b)$ does not contain 4th powers (see [6], Thm. 2). The claim now follows from the fact that the word $\varphi^n(a)$ equals the reversal of $h^{n-1}(b)$, which can be proved by induction. □

Acknowledgements. The author is grateful to the anonymous referees for detailed review of the paper and helpful suggestions.

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press

References

[1] S.I. Adian, *The Burnside Problem and Identities in Groups.* Nauka (1975); English translation, Springer-Verlag (1979) 1–336.
[2] V.S. Atabekyan, Non-φ-admissible normal subgroups of free Burnside groups. *J. Contemp. Math. Anal.* **45** (2010) 112–122.
[3] V.S. Atabekyan, Normal automorphisms of free Burnside groups. *Izv. RAN. Ser. Math.* **75** (2011) 3–18.
[4] E.A. Cherepanov, Free semigroup in the group of automorphisms of the free Burnside group. *Comm. Algebra* **33** (2005) 539–547.
[5] E.A. Cherepanov, Normal automorphisms of free Burnside groups of large odd exponents. *Int. J. Algebra Comput.* **16** (2006) 839–847.
[6] J. Karhumäki, On cube-free ω-words generated by binary morphisms. *Disc. Appl. Math.* **5** (1983) 279–297.
[7] Kourovka Notebook, *Unsolved Problems in Group Theory.* Novosibirsk (2006).
[8] I.G. Lysënok, Infinite Burnside groups of even exponent. *Izv. Math.* **60** (1996) 453–654.
[9] F. Mignosi and G. Pirillo, Repetitions in the Fibonacci infinite word. *Informatique Théorique et Applications* **26** (1992) 199–204.

Communicated by J. Karhumäki.
Received April 12, 2010. Accepted March 1st, 2011.

https://doi.org/10.1051/ita/2011118 Published online by Cambridge University Press