Combinatorics, Probability and Computing (2004) 13, 31–35. © 2004 Cambridge University Press
DOI: 10.1017/S0963548303005820 Printed in the United Kingdom

On Generalized Random Railways

HANS GARMO[1], SVANTE JANSON[1] and MICHAŁ KAROŃSKI[2]

[1]Department of Mathematics, Uppsala University, PO Box 480, S-751 06 Uppsala, Sweden
(e-mail: hans.garmo@math.uu.se, svante.janson@math.uu.se)
http://www.math.uu.se/~svante

[2]Department of Discrete Mathematics, Adam Mickiewicz University, Poznań, Poland
(e-mail: karonski@amu.edu.pl)

and

Department of Mathematics and Computer Science, Emory University, Atlanta, USA
(e-mail: michal@mathcs.emory.edu)
http://main.amu.edu.pl/~karonski

Received 27 February 2002; revised 21 January 2003

We consider a random generalized railway defined as a random 3-regular multigraph where some vertices are regarded as switches that only allow traffic between certain pairs of attached edges. It is shown that the probability that the generalized railway is functioning is linear in the proportion of switches. Thus there is no threshold phenomenon for this property.

1. Introduction

A *railway* is a 3-regular multigraph where each vertex is regarded as a switch where one of the attached edges is regarded as a main track that branches into the two others. Thus, a train that approaches a vertex along the main track (for that vertex) can continue on any of the two other tracks, but a train coming on one of the branching tracks has to leave the vertex on the main track.

A *random railway* is a random 3-regular multigraph where each vertex is regarded as a switch in a randomly chosen orientation. Random railways were introduced and studied by Garmo [6], where the reader can also find examples and a historical background.

Railways can be regarded as a special case of multigraphs with restrictions on the connections at each vertex; we may at each vertex of a multigraph impose any set of rules allowing passage from one edge only to certain others. In other words, instead of the usual line graph of a multigraph G (or its directed version), we consider a certain subgraph of it. Other such (multi)graphs with restrictions have been considered by Andersson [1] and Janson and Wormald [9], but will not be considered here.

https://doi.org/10.1017/S0963548303005820 Published online by Cambridge University Press

A railway is *functioning* if a train starting on an arbitrary edge in an arbitrary direction can reach any other edge while obeying the rule above at each vertex.

Remark. A stronger requirement is that a train starting on an arbitrary edge in an arbitrary direction can reach any other edge in any other direction. Such railways are said to be *two-way functioning* [6]. In other words, a railway is two-way functioning if the corresponding restricted directed edge graph is strongly connected. As is shown in [6], the probability that a random railway is functioning but not two-way functioning decreases exponentially. The same applies to the generalization studied here; thus the results below hold without changes if 'functioning' is replaced by 'two-way functioning'.

Garmo [6] showed that the probability that a random railway with n vertices is functioning converges to $1/3$ as $n \to \infty$. (Since the number of vertices in a 3-regular multigraph is even, we assume throughout the paper that n is even without further comment.) On the other hand, if we consider a random 3-regular multigraph without restrictions, it is a.a.s. functioning, *i.e.*, functioning with probability tending to 1 as $n \to \infty$. Indeed, it is easy to see that a 3-regular multigraph is functioning if and only if it is connected, which holds a.a.s.: see, *e.g.*, [2]. The purpose of the present paper is to study the transition between these cases by considering intermediate cases where we mix switches and ordinary vertices.

We define the generalized random railway $R_{n,p}$ to be a random 3-regular multigraph on n vertices, where each vertex with probability p is replaced by a switch in a random orientation, all random choices being independent. The remaining vertices, which are not made into switches, allow traffic from any edge to any other edge. Alternatively, we could start with a random railway and randomly, with probability $1-p$, replace switches by 'superswitches' allowing traffic from any edge to any other. Hence, $R_{n,1}$ is the random railway considered by Garmo, while $R_{n,0}$ is just a random 3-regular multigraph. The results just given for these extreme cases connect in the following way.

Theorem. *Let $0 \leqslant p \leqslant 1$ be fixed. Then, as $n \to \infty$,*

$$\mathbb{P}(R_{n,p} \text{ is functioning}) \to 1 - 2p/3.$$

We thus have a smooth linear transition as p varies between 0 and 1. This is in contrast to the behaviour of many properties of random graphs (or random subsets in general), where there is a sharp threshold: see, *e.g.*, [4] and [3] for early examples and results and [8] for a survey. In particular, Friedgut [5] has shown that a property of random subsets, or random graphs, has a sharp threshold unless the property only depends on small subsets, or can be approximated in a certain sense by such a 'local' property. In view of this, the result above is somewhat surprising, since 'functioning' is a global property. There are two explanations for this behaviour. First, the random replacement of vertices by switches yields a random subset as in Friedgut's theorem, but there are other random factors too, *viz.* the underlying random 3-regular multigraph and the orientation of the switches, which may blur the threshold. Secondly, and more importantly, as is shown in [6] (see the proof

https://doi.org/10.1017/S0963548303005820 Published online by Cambridge University Press




Figure 1. A paddle wheel

below), the global property of 'functioning' is essentially (*i.e.*, approximately) determined by the local property 'there are no paddle wheels'. Friedgut [5] has given simple artificial examples where a global property is essentially determined by a local property. This paper exhibits an example, although in a more complicated situation, showing that the same can happen for natural global properties, which stresses the need for the rather complicated arguments used in various applications of Friedgut's theorem to specific properties.

Remarks. **(1)** We clearly obtain the same asymptotical result if we consider random 3-regular multigraphs with fixed numbers of switches and ordinary vertices (superswitches).

(2) Another interesting problem is whether $R_{n,p}$ is Hamiltonian, *i.e.*, has an allowed cycle through all vertices. For $p = 0$, the probability of this tends to $1/e$. Indeed, Robinson and Wormald [10] showed that a random 3-regular simple graph is a.a.s. Hamiltonian; their proof extends to showing that a random 3-regular multigraph is a.a.s. Hamiltonian if and only if it lacks loops: see [8]. For $p \geqslant 3(1 - \sqrt{3}/2) \doteq 0.402$, the first moment method (a simple calculation of the expected number of Hamilton cycles) shows that the probability tends to 0. For smaller $p > 0$, we do not know. (We conjecture a sharp threshold.)

2. Proof of the theorem

We use the construction of $R_{n,p}$ where we start with a random railway R with n vertices, and with probability $1 - p$ replace switches by superswitches (ordinary vertices). Clearly, if the railway R is functioning then so is $R_{n,p}$, since we only have added more connections.

As is shown in [6], the main obstruction to R being functioning is the presence of *paddle wheels*, where a k-paddle wheel ($k \geqslant 1$) is a set of k switches connected in a cycle such that the main track of each is connected to one of the branching tracks of the next switch: see Figure 1. It is evident that the presence of a paddle wheel implies that the railway is malfunctioning, since a train entering the paddle wheel is trapped there. Conversely (see [6]),

$$\mathbb{P}(R \text{ is malfunctioning but has no paddle wheels}) \to 0$$

because the probability of the existence of some other trapping configuration in a random railway tends to 0. In other words, a random railway is, a.a.s., functioning if and only if it contains no paddle wheels. Furthermore (see again [6]), the number of paddle wheels is asymptotically Poisson-distributed: more precisely, if X_k is the number of k-paddle wheels

https://doi.org/10.1017/S0963548303005820 Published online by Cambridge University Press




Figure 2. A paddle wheel with some trees leading to it

in R, then

$$X_k \xrightarrow{\mathrm{d}} \mathrm{Po}\left(\frac{1}{k}\left(\frac{2}{3}\right)^k\right), \tag{2.1}$$

jointly for all $k \geqslant 1$. Moreover, we have the uniform bound

$$\mathbb{E}X_k \leqslant \left(\frac{4}{5}\right)^k, \tag{2.2}$$

which enables us to sum (2.1) over all k by a truncation argument [6]. Hence, for the total number of paddle wheels in R,

$$\sum_k X_k \xrightarrow{\mathrm{d}} \mathrm{Po}\left(\sum_{k=1}^{\infty} \frac{1}{k}\left(\frac{2}{3}\right)^k\right) = \mathrm{Po}\big(\ln 3\big).$$

In order to extend these results to $R_{n,p}$, we use further the structure result of [7], which shows that, a.a.s., if the random railway R has $m \geqslant 0$ paddle wheels $W_1, \ldots, W_m$, then the edge set of R can be partitioned in a large set E_0 and sets $E_1, \ldots, E_m$, where E_0 is functioning in the sense that a train starting on any edge in E_0 in any direction can reach any other edge in E_0, while E_i, $i = 1, \ldots, m$, contains the edges such that a train starting there in a certain direction gets trapped by W_i (if it starts on an edge in E_i in the other direction it can always reach E_0). If W_i is a k-paddle wheel, then the set E_i contains the k edges in W_i, the k edges leading in to it, and possibly some further edges forming small trees attached to W_i; see Figure 2. Moreover, any edge may be reached from some, and thus from every, edge in E_0.

Now suppose that R is of this type and that we change it into R', say, by converting some switches into ordinary vertices, thus adding another possible connection through some vertices. It is obvious that if some paddle wheel remains, then a train entering the paddle wheel is trapped and R' is not functioning. On the other hand, it is equally obvious that if no paddle wheel remains, which means that at least one switch in each paddle wheel is replaced by an ordinary vertex, then a train reaching W_i may, through the new connection, find its way back to E_0. Consequently, R' is then functioning.

https://doi.org/10.1017/S0963548303005820 Published online by Cambridge University Press

In particular, doing the random substitution giving $R_{n,p}$, we see that $R_{n,p}$ is a.a.s. functioning if and only if it contains no paddle wheels, and thus

$$\mathbb{P}(R_{n,p} \text{ has no paddle wheels}) - \mathbb{P}(R_{n,p} \text{ is functioning}) \to 0. \tag{2.3}$$

Finally, a k-paddle wheel in R has probability p^k to survive to $R_{n,p}$, so if $X_{k,p}$ is the number of k-paddle wheels in $R_{n,p}$, then it follows immediately from (2.1) that

$$X_{k,p} \xrightarrow{\mathrm{d}} \mathrm{Po}\left(\frac{1}{k}\left(\frac{2}{3}\right)^k p^k\right),$$

jointly for all $k \geqslant 1$. The bound (2.2) again enables us to sum this and obtain

$$\sum_k X_{k,p} \xrightarrow{\mathrm{d}} \mathrm{Po}\left(\sum_{k=1}^{\infty} \frac{1}{k}\left(\frac{2p}{3}\right)^k\right) = \mathrm{Po}(-\ln(1-2p/3)).$$

Consequently,

$$\begin{aligned}\mathbb{P}(R_{n,p} \text{ has no paddle wheels}) &= \mathbb{P}\left(\sum_x X_{k,p} = 0\right) \\ &\to \exp(\ln(1-2p/3)) = 1 - 2p/3,\end{aligned}$$

and the theorem follows by (2.3). □

Acknowledgement

We thank Pontus Andersson for pointing out an error in a preliminary version of this paper.

References

[1] Andersson, P. Hamilton cycles in generalized random railways. In preparation.
[2] Bollobás, B. (2001) *Random Graphs*, 2nd edn, Cambridge University Press.
[3] Bollobás, B. and Thomason, A. (1987) Threshold functions. *Combinatorica* **7** 35–38.
[4] Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. *Publ. Math. Inst. Hungar. Acad. Sci.* **5** 17–61.
[5] Friedgut, E. (1999) Sharp thresholds of graph properties, and the k-sat problem. *J. Amer. Math. Soc.* **12** 1017–1054.
[6] Garmo, H. (1996) Random railways as 3-regular graphs. *Random Struct. Alg.* **9** 113–136.
[7] Garmo, H. (1999) Asymptotic properties of the connectivity number of random railways. *Adv. Appl. Probab.* **31** 720–741.
[8] Janson, S., Łuczak, T. and Ruciński, A. (2000) *Random Graphs*, Wiley, New York.
[9] Janson, S. and Wormald, N. C. Rainbow Hamilton cycles in random regular graphs. In preparation.
[10] Robinson, R. W. and Wormald, N. C. (1992) Almost all cubic graphs are Hamiltonian. *Random Struct. Alg.* **3** 117–125.

https://doi.org/10.1017/S0963548303005820 Published online by Cambridge University Press