The Canadian Entomologist

Index for Volume 141 (2009) / Index pour le volume 141 (2009)



Entomological Society of Canada /
Société d'entomologie du Canada

© 2010 Entomological Society of Canada

https://doi.org/10.4039/entcontin142 Published online by Cambridge University Press

Note – The annual index usually is printed in the last issue of each calendar year. For the tardy appearance of the Volume 141 (2009) index, the Editor-in-Chief apologises to all but especially to those dedicated readers who were eagerly awaiting its publication in the November/December issue.

Nota – L'index annuel parait généralement dans le dernier numéro de l'année civile. Le directeur de la publication tient à présenter ses excuses pour la parution tardive de l'index du volume 141, 2009 à tous et, plus particulièrement, aux fidèles lecteurs qui lisent avec grand intérêt cet index.

Ashekian C. See Sinclair *et al*., 397
Bauce É. See Berthiaume *et al*., 309
Bauce É. See Carleton *et al*., 614
Behan-Pelletier VM, Eamer B. *Ceratozetes* and *Ceratozetoides* (Acari: Oribatida: Ceratozetidae) of North America, 246
Belhabib R *et al*. Host selection and reproductive performance of *Phloeosinus bicolor* (Coleoptera: Curculionidae: Scolytinae) in indigenous and exotic *Cupressus* in Tunisia, 595
Ben Jamaa ML. See Belhabib *et al*., 595
Benttinen J, Preisser E. Avian kleptoparasitism of the digger wasp *Sphex pensylvanicus*, 604.
Berthiaume R. See Carleton *et al*., 614
Berthiaume R *et al*. Daily oviposition pattern of *Lambdina fiscellaria* (Lepidoptera: Geometridae) under laboratory conditions, 309
Björklund N. Non-destructive tree-trunk funnel trap for capturing *Hylobius warreni* (Coleoptera: Curculionidae) ascending stems of trees, 422
Bleiker KP *et al*. Transport of fungal symbionts by mountain pine beetles, 503
Bogdanović AM *et al*. Sexual dimorphism in *Ephedrus persicae* (Hymenoptera: Braconidae: Aphidiinae): intraspecific variation in size and shape, 550
Boone CK *et al*. Assemblage of Hymenoptera arriving at logs colonized by *Ips pini* (Coleoptera: Curculionidae: Scolytinae) and its microbial symbionts in western Montana, 172
Borden JH. See Poland *et al*., 158
Boucher S. *Phytomyza petoei* (Diptera: Agromyzidae), a leaf miner on cultivated mint, newly recorded in the Nearctic Region, 126
Brochu K, Wheeler TA. Systematics and ecology of the Nearctic species of *Neophyllomyza* (Diptera: Milichiidae), 103
Brodo, F. See Starý and Brodo, 1
Buddle CM. See Schroeder *et al*., 515
Cappuccino N. See Norman *et al*., 609
Carleton D *et al*. Seasonal parasitism of hemlock looper (Lepidoptera: Geometridae) eggs in eastern Canada, 614
Carroll AL. See Smith *et al*., 56
Carter ME *et al*. Analysis of genetic diversity in an invasive population of Asian long-horned beetles in Ontario, Canada, 582
Ciaramitaro TM. See Poland *et al*., 158
Cox GH. See Heard and Cox, 406
de Groot P. See Lyons *et al.*, 40
Delisle J. See Carleton *et al*., 614
Digweed SC *et al*. Current status of invasive alien birch-leafmining sawflies (Hymenoptera: Tenthredinidae) in Canada, with keys to species, 201
Duguma D *et al*. Seasonal dynamics of *Urophora quadrifasciata* on spotted knapweed in the Arkansas Ozarks, 70
Dzurisin JDK. See Prior KM *et al*., 161
Eamer B. See Behan-Pelletier and Eamer, 246
Elmhirst J. See Sinclair *et al*., 397
Evenden ML. Biology of *Caloptilia fraxinella* (Lepidoptera: Gracillariidae) on ornamental green ash, *Fraxinus pennsylvanica* (Oleaceae), 31
Evenden ML. See Wins-Purdy *et al*., 86
Foottit RG *et al*. Identification, distribution, and molecular characterization of the apple aphids *Aphis pomi* and *Aphis spiraecola* (Hemiptera: Aphididae: Aphidinae), 478
Foottit RG. See Pike *et al*., 561
Forbes MR. See Norman *et al*., 609
Friesen CP. See Westwood and Friesen, 80

© 2010 Entomological Society of Canada

https://doi.org/10.4039/entcontin142 Published online by Cambridge University Press

Gavloski J. See Wise *et al*., 619
Gibbs JJ. New species in the *Lasioglossum petrellum* species group identified through an integrative taxonomic approach, 371
Gibson GAP, Reigada C. The bizarre male of *Spalangia dozieri* (Hymenoptera: Pteromalidae): adaptations for male phoresy or the result of sexual selection? 112
Gotoh T. See Gries *et al*., 145
Graf G. See Pike *et al*., 561
Grant GG *et al*. Pheromone identification of *Dioryctria abietivorella* (Lepidoptera: Pyralidae) from an eastern North American population: geographic variation in pheromone response, 129
Grebennikov VV. See Jendek and Grebennikov, 236
Gries G. See Gries *et al*., 53
Gries G. See Gries *et al*., 145
Gries R *et al*. *Lymantria dispar* sex pheromone is a behavioral antagonist to pheromonal attraction of male *Lymantria mathura*, 53
Gries R *et al*. Spacing of traps baited with species-specific *Lymantria* pheromones to prevent interference by antagonistic components, 145
Grogan T. See Sinclair *et al*., 397
Hajek AE. See Long *et al*., 153
Hammon R. See Pike *et al*., 561
Harrison RG. See Carter *et al*., 582
Heard SB, Cox GH. Plant module size and attack by the goldenrod spindle-gall moth, 406
Hébert C. See Carleton *et al*., 614
Hébert C. See Berthiaume *et al*., 309
Hellmann JJ. See Prior *et al*., 161
Hueppelsheuser T. See Sinclair *et al*., 397
Ivanović A. See Bogdanović *et al*., 550
Jendek E, Grebennikov VV. *Agrilus sulcicollis* (Coleoptera: Buprestidae), a new alien species in North America, 236
Jones GC. See Lyons *et al*., 40
Judd GJR. See Wins-Purdy *et al*., 86
Kabera A. See Sadeghi *et al*., 95
Kavallieratos NG. See Bogdanović *et al*., 550
Krauth SJ. See Boone *et al*., 172
Kring TJ. See Duguma *et al*., 70
Lamb RJ *et al*. Measuring the performance of aphids: fecundity *versus* biomass, 401
Lamontagne L. See Berthiaume *et al*., 309
Langor D. See Majka *et al*., 317
Langor DW. See Digweed *et al*., 201
Lauzon CR. See Bleiker *et al*., 503
Laws AN. Density-dependent reductions in grasshopper fecundity in response to nematode parasitism, 415
Lee J-W, Lee S-M. Two new species in the genus *Jezarotes* (Hymenoptera: Ichneumonidae: Acaenitinae) from Korea, 496
Lee S-M. See Lee and Lee, 496
Li Z *et al*. Ultrastructure of antennal sensilla of female *Ceratosolen solmsi marchali* (Hymenoptera: Chalcidoidea: Agaonidae: Agaoninae), 463
Lieutier F. See Belhabib *et al*., 595
Lindgren BS. See Smith *et al*., 56
Long SJ *et al*. *Sirex* species (Hymenoptera: Siricidae) and their parasitoids in *Pinus sylvestris* in eastern North America, 153
Lowery DT. See Foottit *et al*., 478
Lowery DT. See Olivier *et al*., 425
Lushai G. See Foottit *et al*., 478
Lyons DB *et al*. Host selection by *Agrilus planipennis* (Coleoptera: Buprestidae): inferences from sticky-band trapping, 40
Mackay PA. See Lamb *et al*.. 401
MacQuarrie CJK. See Digweed *et al*., 201
Majka CG *et al*. Latridiidae (Coleoptera) of Atlantic Canada: new records, keys to identification, new synonyms, distribution, and zoogeography, 317
Mann J. See Sinclair *et al*., 397
Maw HEL. See Foottit *et al*., 478
Maw HEL. See Pike *et al*., 561

© 2010 Entomological Society of Canada

https://doi.org/10.4039/entcontin142 Published online by Cambridge University Press

Michiels K. See Sadeghi *et al*., 95
Migui SM. See Lamb *et al*.. 401
Millar JG. See Grant *et al*., 129
Miller DG. See Pike *et al*., 561
Morneau L. See Digweed *et al*., 201
Nakamura K. See Gries *et al*., 53
Norman K *et al*. Parasitism of a successful weed biological control agent, *Neogalerucella calmariensis*, 609
Nouira S. See Belhabib *et al*., 595
Nystrom KL. See Digweed *et al*., 201
Olivier CY *et al*. Phytoplasma diseases and their relationships with insect and plant hosts in Canadian horticultural and field crops, 425
Pelini SL. See Prior *et al*., 161
Peng Y. See Li *et al*., 463
Picard I. See Berthiaume *et al*., 309
Pike KS *et al*. New species of *Braggia* (Hemiptera: Aphididae) on buckwheat in western North America, 561
Pinault L *et al*. Interaction of foliage and larval age influences preference and performance of a geometrid caterpillar, 136
Poland TM *et al*. 1-Octen-3-ol is repellent to *Ips pini* (Coleoptera: Curculionidae: Scolytinae) in the midwestern United States, 158
Potter SE. See Bleiker *et al*., 503
Preisser E. See Benttinen and Preisser, 604
Prior KM *et al*. Biology of larvae and adults of *Erynnis propertius* at the northern edge of its rang, 161
Pureswaran DS. See Poland *et al*., 158
Quiring D. See Carleton *et al*., 614
Quiring D. See Pinault *et al*., 136
Raffa KF. See Boone *et al*., 172
Reigada C. See Gibson and Reigada, 112
Royer L. See Carleton *et al*., 614
Rücker WH. See Majka *et al*., 317
Sadeghi A *et al*. Acute and chronic insecticidal activity of a new mannose-binding lectin from *Allium porrum* against *Acyrthosiphon pisum via* an artificial diet, 95
Saint-Germain M. See Schroeder *et al*., 515
Schaefer PW. See Gries *et al*., 53
Schaefer PW. See Gries *et al*., 145
Scharbach R. See Lyons *et al*., 40
Schroeder B *et al*. Activity of flying beetles (Coleoptera) at two heights in canopy gaps and intact forests in a hardwood forest in Quebec, 515
Sinclair BJ *et al*. *Dasineura plicatrix* (Diptera: Cecidomyiidae): a recent introduction into North America, 397
Six DL. See Bleiker *et al*., 503
Six DL. See Boone *et al*., 172
Smagghe G. See Sadeghi *et al*., 95
Smirle MJ. See Foottit *et al*., 478
Smith GD *et al*. Life history of a secondary bark beetle, *Pseudips mexicanus* (Coleoptera: Curculionidae: Scolytinae), in lodgepole pine in British Columbia, 56
Smith MT. See Carter *et al*., 582
Spence JR. See Digweed *et al*., 201
Sperling FAH. See Sperling and Sperling, 521
Sperling JLH, Sperling FAH. Lyme borreliosis in Canada: biological diversity and diagnostic complexity from an entomological perspective, 521
Starý J, Brodo, F. Arctic species of the subgenus *Symplecta sensu stricto* (Diptera: Limoniidae), 1
Starý P. See Bogdanović *et al*., 550
Starý P. See Pike *et al*., 561
Stobbs LW. See Olivier *et al*., 425
Takács S. See Gries *et al*., 145
Thurston G. See Pinault *et al*., 136
Tomanović Ž. See Bogdanović *et al*., 550
Trudel R. See Grant *et al*., 129
Turgeon JJ. See Carter *et al*., 582

© 2010 Entomological Society of Canada

https://doi.org/10.4039/entcontin142 Published online by Cambridge University Press

Turnock WJ. See Wise *et al*., 619
Van Damme EJM. See Sadeghi *et al*., 95
Westwood AR, Friesen CP. Occurrence and habitat of the endangered white flower moth, *Schinia bimatris* (Lepidoptera: Noctuidae), in Manitoba, 80
Wheeler TA. See Brochu and Wheeler, 103
Whitehouse C. See Wins-Purdy *et al*., 86
Wiedenmann RN. See Duguma *et al*., 70
Wins-Purdy AH *et al*. Effect of horticultural oil on oviposition behaviour and egg survival in the oblique banded leafroller (Lepidoptera: Tortricidae), 86
Williams DJM. See Digweed *et al*., 201
Williams DW. See Long *et al*., 153
Wise IL *et al*. Sequential decision plan for controlling *Mamestra configurata* in spring canola, 619
Yang D. See Li *et al*., 463
Yang P. See Li *et al*., 463
Žikić V. See Bogdanović *et al*., 550

© 2010 Entomological Society of Canada

https://doi.org/10.4039/entcontin142 Published online by Cambridge University Press