Ergod. Th. & Dynam. Sys. (2014), **34**, 1914–1929 © *Cambridge University Press, 2013*
doi:10.1017/etds.2013.34

Robust minimality of iterated function systems with two generators

ALE JAN HOMBURG†‡ and MEYSAM NASSIRI§

† *KdV Institute for Mathematics, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
‡ *Department of Mathematics, VU University Amsterdam, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands*
(e-mail:a.j.homburg@uva.nl)
§ *School of Mathematics, Institute for Research in Fundamental Sciences (IPM), PO Box 19395-5746, Tehran, Iran*
(e-mail:nassiri@ipm.ir)

(*Received* 8 *January* 2013 *and accepted in revised form* 8 *April* 2013)

Abstract. We prove that every compact manifold without boundary admits a pair of diffeomorphisms that generates C^1 robustly minimal dynamics. We apply the results to the construction of blenders and robustly transitive skew product diffeomorphisms.

1. *Introduction*

In this paper we study robust minimality of iterated function systems. By an iterated function system (IFS) one means the action of the semigroup generated by a family of diffeomorphisms. The study of IFSs, apart from being important in itself, has a remarkable role in understanding certain (single) dynamical systems. For instance, one can embed an IFS into a skew product over the full shift with a sufficient number of symbols, so that it shows up in many dynamical systems exhibiting some form of hyperbolicity.

Let $\mathcal{F}$, $\mathcal{G}$ be two families of diffeomorphisms on a compact manifold M. Denote

$$\mathcal{F} \circ \mathcal{G} = \{f \circ g \mid f \in \mathcal{F}, g \in \mathcal{G}\}$$

and, for $k \in \mathbb{N}$,

$$\mathcal{F}^k := \mathcal{F}^{k-1} \circ \mathcal{F}, \quad \mathcal{F}^0 := \{\mathrm{Id}\}.$$

Write $\langle \mathcal{F} \rangle^+$ for the semigroup generated by $\mathcal{F}$, that is, $\langle \mathcal{F} \rangle^+ = \bigcup_{n=0}^{\infty} \mathcal{F}^k$. The action of the semigroup $\langle \mathcal{F} \rangle^+$ is called the *iterated function system associated with* $\mathcal{F}$ and we denote it by IFS($\mathcal{F}$). For $x \in M$, we write the orbits of the action of this semigroup as

$$\langle \mathcal{F} \rangle^+(x) = \{f(x) : f \in \langle \mathcal{F} \rangle^+\}.$$

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

A sequence $\{x_n : n \in \mathbb{N}\}$ is called a branch of an orbit of IFS($\mathcal{F}$) if for each $n \in \mathbb{N}$ there is $f_n \in \mathcal{F}$ such that $x_{n+1} = f_n(x_n)$. We say that IFS($\mathcal{F}$) is minimal if every orbit has a branch which is dense in M. We say that a property P holds C^r robustly for IFS($\mathcal{F}$) if it holds for IFS($\tilde{\mathcal{F}}$) for every family $\tilde{\mathcal{F}}$ whose elements are C^r perturbations of elements of $\mathcal{F}$.

Here we prove the following theorem.

THEOREM A. *Every boundaryless compact manifold admits a pair of diffeomorphisms that generates a C^1 robustly minimal IFS.*

The number of generators in Theorem A is optimal. Indeed, a single diffeomorphism cannot be C^1 robustly minimal. Recall that a diffeomorphism is minimal if every orbit is dense, but Pugh's closing lemma yields the existence of periodic points for C^1 generic diffeomorphisms. There are several examples of robustly minimal IFSs: in dimension one [**5, 8**]; in any dimension but local and with many generators [**13**]; on every boundaryless compact manifold and with many generators [**6**] and where at least three generators are required [**7**].

The study of minimal IFSs has important consequences for the study and construction of robustly transitive diffeomorphisms. Recall that a diffeomorphism is transitive if it has a dense orbit, and it is (C^1) robustly transitive if every nearby diffeomorphism (in the C^1 topology) is also transitive. Some parts of the proof of Theorem A are closely related to the construction of blenders. The notion of blender introduced by Bonatti and Díaz [**3**] is the main tool to construct robustly transitive diffeomorphisms (cf. [**15**]). The results of this paper permit us to give a straightforward and general construction of blenders and robustly transitive dynamics in arbitrary dimension along the lines of [**14**]. It can be summarized as follows (cf. Theorem 5.6):

> *Let F be a diffeomorphism with invariant compact partially hyperbolic set Γ such that $F|_\Gamma$ is conjugate (appropriately) to an iterated function system* IFS($\mathcal{F}$). *If* IFS($\mathcal{F}$) *is* strongly robustly minimal*, then F is* robustly transitive *on* Γ.

The strong robust minimality of an IFS roughly means that one may change the perturbed family of maps at each iteration and still obtain dense orbits for all points (cf. Definition 4.1). The main result in this direction is Theorem 4.2, which implies the following improvement of Theorem A.

THEOREM B. *Every boundaryless compact manifold admits a pair of diffeomorphisms that generates a C^1 strongly robustly minimal IFS.*

We provide two constructions that prove Theorem A, in §§2 and 3. These constructions yield different types of examples in the applications to robust transitivity. In §4 we study the strong robust minimality of IFSs and then we complete the proof of Theorem B. The connection to blenders and an application to the construction of robustly transitive skew product diffeomorphisms are presented in §5.

2. *Minimal iterated function systems*

In this section we prove Theorem A.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

2.1. *Local robust minimality.*

Definition 2.1. Let $\mathcal{F}$ be a family of maps on the metric space X, and $Y \subset X$. We say that IFS($\mathcal{F}$) is *minimal on* Y if for every $x \in Y$, the orbit $\langle \mathcal{F} \rangle^+(x)$ has a branch which is dense in Y.

Recall that a map ϕ on a metric space (X, d) is a contraction if and only if there is a constant $0 < \kappa < 1$ such that $d(\phi(x), \phi(y)) \leq \kappa d(x, y)$, for all $x, y \in X$.

The following proposition is a modification of [**14**, Proposition 2.3] (see also [**13**]). We give a short proof in §4.

PROPOSITION 2.2. *Let U and V be two open disks in $\mathbb{R}^n$ containing 0, and $\phi : U \to V$ be a diffeomorphism with $\phi(0) = 0$. If $D\phi_0$ is a contraction, then there exists $k \in \mathbb{N}$ depending only on $D\phi_0$ such that for any small $\varepsilon > 0$ there exist $r > 0$ and vectors $c_1, \ldots, c_k \in B_\varepsilon(0)$ such that*

$$B_r(0) \subset \overline{\mathcal{O}^+_{\mathcal{F}}(0)},$$

where $\mathcal{F} = \{\phi + c_1, \ldots, \phi + c_k\}$ and $\mathcal{O}^+_{\mathcal{F}}(0)$ is a branch of the orbit $\langle \mathcal{F} \rangle^+(0)$. Moreover, IFS($\mathcal{F}$) *is robustly minimal on $B_r(0)$.*

2.2. *Proof of Theorem A.*

Definition 2.3. (Weak hyperbolic point) Let p be a hyperbolic periodic point of a diffeomorphism g of period k. We say that p is *δ-weak hyperbolic* if

$$1 - \delta < \mathrm{m}(D_p g^k|_{E^s_p}) < \|D_p g^k|_{E^s_p}\| < 1 < \mathrm{m}(D_p g^k|_{E^u_p}) < \|D_p g^k|_{E^u_p}\| < \frac{1}{1-\delta}.$$

Here E^s_p and E^u_p are the stable and unstable subspaces at p.

Proof of Theorem A. Let M be a compact manifold of dimension $m \geq 2$. We refer to [**5**] or [**8**] for iterated functions systems in dimension one. The proof has three steps.

Step 1. It is a standard consequence of Morse theory that one can take a Morse–Smale diffeomorphism $f_\circ$ on M with a unique attracting fixed point o. We assume that the norm of $Df_\circ^{-1}(o)$ is sufficiently close to one (i.e. o is a δ-weak hyperbolic attractor with sufficiently small δ).

We then modify slightly the dynamics of $f_\circ$ in a neighborhood U of o in order to obtain a new Morse–Smale diffeomorphism f such that it has a periodic attracting orbit $\{f^i(p)\}$ in U with sufficiently large period n and with very weak attraction. The period of the attracting orbit and the rate of its weak attraction depend on the dimension of the manifold.

To do the modification we may assume that the open set U is diffeomorphic to the product $\mathbb{D}^2 \times \mathbb{D}^{m-2}$. If $m = 2$, then we replace the dynamics on U by the local diffeomorphism f on $\mathbb{D}^2$ as in Figure 1.

If $m > 2$ then we define h_1 as the two-dimensional map on $\mathbb{D}^2$ defined for the case $m = 2$, and a contraction h_2 with a unique fixed point o_2 on $\mathbb{D}^{m-2}$ so that o_2 is a δ-weak hyperbolic attractor, with a small $\delta > 0$. Then we modify f on U to be the product $h_1 \times h_2$.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press




FIGURE 1. Dynamics of f if $\dim(M) = 2$ (left) and if $\dim(M) > 2$ (right).

Step 2. (Local minimality) Let $\mathcal{V}$ be a small neighborhood of f such that it only contains Morse–Smale diffeomorphisms. We find a diffeomorphism $g \in \mathcal{V}$ arbitrarily close to f in such a way that its dynamics close to the periodic orbit $\mathcal{O}_f(p) = \{f^i(p)\}$ is slightly different from f and is chosen so that the IFS of $\{f, g\}$ is robustly minimal in a neighborhood of the periodic orbit $\{f^i(p)\}_i$. Here we use the results of §2.1.

Let $V_i := f^i(V_0)$ be sufficiently small neighborhoods of $f^i(p)$, for $i = 0, \ldots, n-1$. By the assumptions, $1 - \epsilon < \mathrm{m}(Df^n(p)) < \|Df^n(p)\| < 1$ for some small $\epsilon > 0$. So we may apply Proposition 2.2 for the map f^n restricted to V_0, which gives very small vectors $c_1, \ldots, c_k$ (where $k < n$ if δ is small enough) on the local coordinates on V_0 so that the IFS of $\{f^n, T_{c_1} \circ f^n, \ldots, T_{c_k} \circ f^n\}$ is robustly minimal on a neighborhood D_0 of p. Here T_c is defined in local coordinates as the translation by the vector c.

Let $g_\circ$ be a diffeomorphism on M such that on V_{n-i} it is equal to $f^{1-i} \circ T_{c_i} \circ f^i$. Observe that $g_\circ$ is well defined as the V_i are disjoint and the c_i are sufficiently small. Moreover, we may assume that $g_\circ$ is arbitrarily close to f by taking c_i small enough. In particular, $g_\circ \in \mathcal{V}$, so it is Morse–Smale. Further, $g_\circ^n$ on V_0 is equal to $T_{c_1} \circ T_{c_n} \circ f^n$ and $f^{i-1} \circ g_\circ \circ f^{n-i}$ on V_0 is equal to $T_{c_i} \circ f^n$. Let g be any diffeomorphism sufficiently close to $g_\circ$ in the C^1 topology. Then the IFS of $\{f, g\}$ is robustly minimal on D_0. We call such a local set with robustly minimal dynamics a *blending region* of the IFS.

Observe that $\hat{g} := g \circ f^{-2}$ is very close to f^{-1} and so it is Morse–Smale, provided that $g_\circ$ is close enough to f. Moreover, the semigroup generated by $\{f, g \circ f^{-2}\}$ contains the semigroup generated by $\{f, g\}$.

From now on, we denote $g \circ f^{-2}$ by $\hat{g}$ for a given g. Thus, for any g close to $g_\circ$, IFS($\{f, \hat{g}\}$) is robustly minimal on D_0. We may also assume that for any g close to $g_\circ$, $\hat{g}$ has a periodic repeller $p' \in D_0$ such that $W^u(p')$ contains D_0.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

Step 3. (Globalization) We show that there exists a diffeomorphism g close to $g_\circ$ such that both positive and negative orbits of the blending region D_0 of IFS($\{f, \hat{g}\}$) cover the entire manifold M. If so, then, for any point $x \in M$ and any open set $O \subset M$, there exist diffeomorphisms $h_1, h_2 \in \langle\{f, \hat{g}\}\rangle^+$ such that $h_1(x) \in D_0$ and $h_2^{-1}(O) \cap D_0 \neq \emptyset$. Since D_0 is a blending region of IFS($\{f, \hat{g}\}$), it follows that there exists $h_3 \in \langle\{f, \hat{g}\}\rangle^+$ such that $h_3(h_1(x)) \in h_2^{-1}(O) \cap D_0$, and so for $h := h_2 \circ h_3 \circ h_1 \in \langle\{f, \hat{g}\}\rangle^+$ we have $h(x) \in O$.

Thus to complete the proof of Theorem A it remains to show the above statement. Recall that for a Morse–Smale diffeomorphism the manifold M is the union of the stable manifolds of finitely many periodic points. Also, the union of basins of attraction of periodic attractors for a Morse–Smale diffeomorphism is an open and dense subset of the manifold: the stable manifold $W^s(p)$ for f and the unstable manifold $W^u(p')$ for $\hat{g}$ are open and dense.

Pick g so that $\hat{g}$ and f have no common periodic points, while $\hat{g}$ maps the periodic points of f into the basin of attraction of $\{f^i(p)\}$. The negative orbits of D_0 then cover M: pick a point $x \in M$, then iterating x by f sufficiently often brings it close to some periodic orbit of f since f is Morse–Smale, a further iterate by $\hat{g}$ maps it into the basin of attraction of $\{f^i(p)\}$, after which an iterate of f brings it into D_0. Similarly for positive orbits of D_0. They cover M if f^{-1} maps the periodic points of $\hat{g}$ into the basin of attraction of the periodic attractor for $\hat{g}^{-1}$.

Since these basins of attraction are open and dense in M, these properties can be achieved for g arbitrarily close to $g_\circ$. This completes the proof of Theorem A. □

3. *Alternative construction*

In this section we describe an alternative argument leading to Theorem A. The local part of the argument, producing a blending region, is through an alternative construction. The global part, the mechanism to iterate into and out of the blending region, will be the same. We start with the construction of minimal IFSs generated by affine maps on Euclidean spaces.

3.1. *Iterated function systems generated by affine maps.*

We will provide an affine contraction S and an affine expansion T (i.e. T^{-1} is a contraction) so that the IFS generated by S and T is minimal on all of $\mathbb{R}^m$. Here we assume $m \geq 2$; compare [**2**] for minimal IFSs on $[0, \infty)$ generated by affine maps.

A set Δ is called invariant for an iterated function system IFS($\mathcal{F}$) if $\mathcal{F}(\Delta) = \Delta$. Consider an iterated function system IFS($\mathcal{F}$) generated by contractions on $\mathbb{R}^n$. Such an IFS admits a unique compact invariant set that contains all limit points of branches. This follows from the observation that $\mathcal{F}$ is a contraction in the Hausdorff metric on the class of compact sets [**11**]. We call such a set the attractor of the IFS.

The constructed affine maps S, T will be such that $S \circ T$ is a contraction and the IFS generated by the two affine contractions S and $S \circ T$ possesses an attractor with non-empty interior. Similar ideas were independently used by Volk [**16**] in his construction of persistent attractors for endomorphisms.

Consider the rotation R in $\mathbb{R}^m$, $m \geq 2$,

$$R(x_1, \ldots, x_m) = (\pm x_m, x_1, \ldots, x_{m-1}),$$

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

where the sign is such that $R \in \mathrm{SO}(m)$: a minus sign for even m and a plus sign for odd m. Define further the translation $H(x_1, \ldots, x_m) = (x_1 + s, x_2, \ldots, x_m)$ and the affine map S on $\mathbb{R}^m$ by

$$S(x_1, \ldots, x_m) = H \circ rR(x_1, \ldots, x_m) = (\pm r x_m + s, r x_1, \ldots, r x_{m-1}),$$

for constants $0 < r < 1$, $s > 0$. Likewise, define an affine map T on $\mathbb{R}^m$ by

$$T(x_1, \ldots, x_m) = \left(-a x_1, a x_2, \ldots, a x_{m-1}, -a x_m - 2\frac{s}{r}\right)$$

with $a > 1$. Similar to S, T is the composition of a map from $\mathrm{SO}(m)$ which is multiplied by a factor, a, and a translation. Note that S is a contraction by $r < 1$, while T is an expansion as $a > 1$. Compute

$$\begin{aligned} S \circ T(x_1, \ldots, x_m) &= H^{-1} \circ arR(x_1, \ldots, x_m) \\ &= (\mp arx_m - s, -arx_1, arx_2, \ldots, arx_{m-1}). \end{aligned}$$

The affine map $S \circ T$ is a contraction for $ar < 1$.

LEMMA 3.1. *There are constants $0 < r < 1$, $s > 0$, $a > 1$ with $ar < 1$ such that the iterated function system* IFS($\mathcal{G}$) *with $\mathcal{G} = \{S, S \circ T\}$ has an attractor Δ with non-empty interior. Moreover, this interior contains the fixed point of T.*

Proof. Define the box $B(1, v_2, \ldots, v_m)$ with corners $(\pm 1, \pm v_2, \ldots, \pm v_m)$. We will find r, s, a and $v_2, \ldots, v_m$ such that

$$S(B) \cup S \circ T(B) \supset B. \tag{1}$$


This is a consequence of the conditions

$$r v_m + s > 1, 0 > -r v_m + s, r > v_2, r v_2 > v_3, \ldots, r v_{m-1} > v_m. \tag{2}$$

In fact, (2) implies

$$ar v_m + s > 1, 0 > -ar v_m + s, ar > v_2, ar v_2 > v_3, \ldots, ar v_{m-1} > v_m, \tag{3}$$

and (2) and (3) together give (1).



https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

The repelling fixed point of T is located at $(0, \ldots, 0, -2s/(r(a+1)))$. It lies in B if

$$2s < v_m r(a+1). \tag{4}$$

Now, (2) and (4) can be satisfied by taking suitable $v_m < \cdots < v_2 < r$ all near 1 and $s = 1 - v_m^2$ near 0 so that (2) holds, and a with $ar < 1$.

Since S and $S \circ T$ are contractions, there is a ball O that is mapped into itself by both S and $S \circ T$, that is, $\mathcal{G}(O) \subset O$. Thus

$$\Delta = \lim_{p \to \infty} \mathcal{G}^p(O)$$

is a non-empty compact set that is invariant for IFS($\mathcal{G}$). Since S and $S \circ T$ are contractions, Δ is the unique compact set that is invariant for $\mathcal{G}$ [**11**]. Because $\mathcal{G}(B) \supset B$, the set Δ contains B. □

COROLLARY 3.2. *The iterated function system* IFS($\{S, T\}$) *is minimal on* $\mathbb{R}^m$.

Proof. Note first that $\mathcal{G}$ is minimal on Δ. Indeed, $\mathcal{G}^n(\Delta)$ consists of 2^n sets with diameter shrinking to zero as $n \to \infty$ and together covering Δ. So, given $x \in \Delta$ and open V intersecting Δ, $\langle \mathcal{G} \rangle^+(x)$ intersects V. This implies that $\mathcal{G}$ is minimal on Δ.

The corollary follows, as in the globalization step in the proof of Theorem A in the previous section, since both backward iterates of Δ under IFS($\{S, T\}$) cover $\mathbb{R}^n$ (true since S is a contraction with attracting fixed point inside Δ) and forward iterates of Δ under IFS($\{S, T\}$) cover $\mathbb{R}^n$ (true since T^{-1} is a contraction with attracting fixed point inside Δ). □

The proof of Lemma 3.1 gives more than its statement as it includes arguments for C^1 robust occurrence of invariant sets with non-empty interior. Denote by $\mathrm{Diff}^1(\mathbb{R}^m)$ the set of diffeomorphisms on $\mathbb{R}^m$, endowed with the C^1 compact-open topology.

COROLLARY 3.3. *There exists a neighborhood* $W \subset \mathrm{Diff}^1(\mathbb{R}^m) \times \mathrm{Diff}^1(\mathbb{R}^m)$ *of* $(S, S \circ T)$ *such that for each* (f_1, f_2) *in this neighborhood,* IFS($\mathcal{F}$) *with* $\mathcal{F} = \{f_1, f_2\}$ *admits an invariant set with non-empty interior.*

In rescaled coordinates $(y_1, \ldots, y_m) = h(x_1, \ldots, x_m) = (\delta x_1, \ldots, \delta x_m)$, $\delta > 0$, the attractor Δ is multiplied by a factor δ. The affine maps computed in the $(y_1, \ldots, y_m)$ coordinates become

$$h \circ S \circ h^{-1}(y_1, \ldots, y_m) = (\pm r y_m + \delta s, r y_1, \ldots, r y_{m-1})$$

and

$$h \circ T \circ h^{-1}(y_1, \ldots, y_m) = \left(-a y_1, a y_2, \ldots, a y_{m-1}, -a y_m - 2\frac{\delta s}{r} \right);$$

the maps are unaltered except for the translation vector which is multiplied by δ. In other words, if S and T are affine maps as above so that $\mathcal{G}(\mathbb{R}^m; S, T)$ has an attractor Δ, then replacing s in their expressions by δs yields an IFS with attractor $\delta\Delta$.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

3.2. *Second proof of Theorem A.* The proof of Theorem A in §2.2 consists of a local part, the construction of a blending region in steps 1 and 2, and a global part in step 3 involving a mechanism to move from the blending region to other parts of the manifold, and back, by iterating. The same construction can be followed here, where Lemma 3.1 used in a chart on the manifold provides a blending region. Compare [**6**].

Specifically we obtain two diffeomorphisms g_1, g_2 on M generating a robustly minimal IFS with the following properties:

(1) for $i = 1, 2$, g_i has a unique attracting fixed point p_i and a unique repelling fixed point q_i;

(2) there is a blending region, containing p_1 and q_2, on which g_1 and $g_1 \circ g_2$ are contractions.

The diffeomorphisms g_1, g_2 are not C^1 close to the identity, but the construction can be done so that $\mathrm{m}(Dg_1)$, $|Dg_1|$, $\mathrm{m}(Dg_2)$, $|Dg_2|$ are everywhere close to 1.

4. *Strong robust minimality*

In this section we introduce the notion of strong robust minimality for IFSs.

4.1. *Notation and definitions.* Let X be a metric space, and $\mathcal{F}$ be a family of maps on X.

- For any $Y \subset X$ and $x \in X$, we denote $\mathcal{F}(Y) := \bigcup_{\phi\in\mathcal{F}} \phi(Y)$ and $\mathcal{F}(x) := \mathcal{F}(\{x\})$.
- If each element of $\mathcal{F}$ is invertible, then we denote $\mathcal{F}^{-1} = \{f^{-1} \mid f \in \mathcal{F}\}$.
- Let $\mathbf{F} = (\mathcal{F}_1, \mathcal{F}_2, \dots)$, where $\mathcal{F}_n$ is a family of maps on X, for any $n \in \mathbb{N}$. Then we denote $\mathcal{F}^{(n)} = \mathcal{F}_n \circ \cdots \circ \mathcal{F}_1$, $\mathcal{F}^{(1)} = \mathcal{F}_1$, $\mathcal{F}^{(0)} = \{\mathrm{Id}\}$,

$$\langle\langle \mathbf{F} \rangle\rangle^+ := \bigcup_{n=0}^{\infty} \mathcal{F}^{(n)},$$

  and its orbits by $\langle\langle \mathbf{F} \rangle\rangle^+(x) := \{f(x) : f \in \langle\langle \mathbf{F} \rangle\rangle^+\}$, for $x \in X$.
- We also denote $\mathcal{F}_m^{(n)} := \mathcal{F}_{m+n} \circ \cdots \circ \mathcal{F}_{m+1}$, for $n, m \in \mathbb{N} \cup \{0\}$.
- Let $x \in X$. A sequence $\{x_n : n \in \mathbb{N}\} \subset X$ is called a branch of the orbit $\langle\langle \mathbf{F} \rangle\rangle^+(x)$ if for any $n \in \mathbb{N}$ there is $f_n \in \mathcal{F}^{(n)}$ such that $x_{n+1} = f_n(x_n)$.

Let $\mathcal{F}$ be a family (with k elements) of C^r diffeomorphisms of a manifold M. A neighborhood $\mathcal{U}$ of $\mathcal{F}$ is the set of all families $\mathcal{F}'$ whose elements are C^r perturbations of elements of $\mathcal{F}$.

Definition 4.1. We say that $\mathrm{IFS}(\mathcal{F})$ is C^r *strongly* robustly minimal if there is a neighborhood $\mathcal{U}$ of $\mathcal{F}$ such that for any $x \in M$ and any sequence $\mathbf{F} = (\mathcal{F}_1, \mathcal{F}_2, \dots)$ in $\mathcal{U}$ there is a branch of the orbit $\langle\langle \mathbf{F} \rangle\rangle^+(x)$ which is dense in M.

It is clear that this is stronger than the robust minimality defined in the introduction. We will see how this notion is natural and effective when one applies the robust minimality of IFSs to construct examples of robustly transitive diffeomorphisms.

We say that a map ϕ on a metric space (X, d) is bi-Lipschitz with constants $\lambda > 0$ and κ, if $\lambda d(x, y) \le d(\phi(x), \phi(y)) \le \kappa d(x, y)$ for all $x, y \in X$. If $0 < \kappa < 1$ then ϕ is a contraction. Then, we also say λ is the *contraction lower bound* of ϕ.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

4.2. *Sufficient conditions for strong robustness.*

THEOREM 4.2. *Let M be a boundaryless compact manifold and D be an open subset of M. Let $\mathcal{F}$ be a finite family of diffeomorphisms of M such that:*
(1) *for any $f \in \mathcal{F}$, $f|_{\overline{D}}$ is a contraction;*
(2) $\overline{D} \subset \mathcal{F}(D)$.
Then IFS($\mathcal{F}$) *is strongly robustly minimal on D.*

Let $\mathcal{G}$ be a family of diffeomorphisms of M such that $\mathcal{F} \subset \mathcal{G}$ and $M = \mathcal{G}(D) = \mathcal{G}^{-1}(D)$. Then IFS($\mathcal{G}$) *is strongly robustly minimal (on M).*

Before proving the theorem we prove a basic lemma.

LEMMA 4.3. *Let X be a connected metric space, and $D \subset X$ be a bounded open set. Let $0 < \lambda < \kappa < 1$. Let $\mathbf{F} = (\mathcal{F}_1, \mathcal{F}_2, \dots)$ be such that for any $n \in \mathbb{N}$, $\mathcal{F}_n$ is a family of homeomorphisms of X such that:*
(1) *for any $f \in \mathcal{F}_n$, $f|_D$ is bi-Lipschitz with constants λ and κ;*
(2) $\overline{D} \subset \mathcal{F}_n(D)$;
(3) *Z_n is δ-dense in D, where $Z_n = \{x \in D \mid f(x) = x \text{ for some } f \in \mathcal{F}_n\}$, and $2\delta > 0$ is smaller than the Lebesgue number of a covering $\mathcal{A}$ of $\overline{D}$ with $\mathcal{A} \subseteq \{f(D) \cap \overline{D} \mid f \in \mathcal{F}_n\}$.*
Then for any $x \in D$, $\langle\langle \mathbf{F} \rangle\rangle^+(x)$ has a branch which is dense in D.

Proof. It is enough to prove the following.

CLAIM 4.4. *Let $B \subset D$ be an open ball. Then there is $n \in \mathbb{N}$ such that for any $m \in \mathbb{N} \cup \{0\}$ there is $f \in \mathcal{F}_m^{(n)} = \mathcal{F}_{m+n} \circ \cdots \circ \mathcal{F}_{m+1}$ such that $f(D) \subset B$.*

Proof. Assume that $B = B_r(x)$ for some $x \in D$. Let $k \in \mathbb{N}$ be (the smallest integer) larger than $\ln(\delta/\mathrm{diam}(D))/\ln(\kappa)$.

(a) If $r \geq 2\delta$, let $B' = B_\delta(x)$ be the ball of radius δ and the same center. Then it follows from (3) that for any $i \in \mathbb{N}$, B' contains some $z \in Z_i$. Thus, there is some $f_i \in \mathcal{F}_i$ with an attracting fixed point in B'. By applying this fact at most k times we see that there is $f \in \mathcal{F}_{m+k} \circ \cdots \circ \mathcal{F}_{n+2} \circ \mathcal{F}_{m+1}$ such that $f(D) \subset B$. In particular, the lemma follows for $n = k$.

(b) If $r < 2\delta$, let $k(r) \in \mathbb{N}$ be the smallest integer larger than $\ln(r/(2\delta))/\ln(\lambda)$. Then, it follows from (2) that for any $i \in \mathbb{N}$, there is some $g_i \in \mathcal{F}_i$ such that $B \subset g_i(D)$. Then by (1), $g_i^{-1}(B) \subset D$ and it contains a ball of radius $r\lambda^{-1}$. By applying this fact at most $k(r)$ times we see that there is $g \in \mathcal{F}_{i+k(r)} \circ \cdots \circ \mathcal{F}_{i+1}$ such that $g^{-1}(B \cap g(D))$ contains a ball of radius 2δ. Thus from (a), for any $m \in N \cup \{0\}$, there is $h \in \mathcal{F}_{m+k} \circ \cdots \circ \mathcal{F}_{m+2} \circ \mathcal{F}_{m+1}$ such that $h(D) \subset g^{-1}(B \cap g(D))$. In particular, $(g \circ h)(D) \subset B$. For $i = m + k$, $n = k + k(r)$ and $f = g \circ h$, it follows that $f(D) \subset B$ and $f \in \mathcal{F}_{m+n} \circ \cdots \circ \mathcal{F}_{m+2} \circ \mathcal{F}_{m+1}$, completing the proof of claim. □

Now, by considering a countable topological base for D and applying the claim repeatedly one obtains a branch of $\langle\langle \mathbf{F} \rangle\rangle^+(x)$ which is dense in D. □

Remark 4.5. Conditions (2) and (3) in Lemma 4.3 are equivalent to what are called the covering and the well-distributed properties respectively in [**14**, Definition 2.4].

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

Proof of Theorem 4.2. Let $\delta > 0$ be a number such that 2δ is smaller that the Lebesgue number of the covering $\mathcal{A} = \{f(D) \cap \overline{D} \mid f \in \mathcal{F}\}$ of $\overline{D}$.

Let Y_n be the set of fixed points of elements of $\mathcal{F}^n$. It follows from the main result of [**11**] that there is $n_0 \in \mathbb{N}$ large enough such that for any $n \geq n_0$, Y_n is $(\delta/2)$-dense in D.

It is easy to see that the family $\mathcal{F} \cup \mathcal{F}^{n_0}$ satisfies hypotheses (1)–(3) of Lemma 4.3 for some bi-Lipschitz constants $0 < \lambda^{1/2} < \kappa^2 < 1$. Let $\mathcal{V}$ be a sufficiently small neighborhood of $\mathcal{F}$, and let $\mathbf{F} = (\mathcal{F}_1, \mathcal{F}_2, \dots)$ be a sequence of perturbations of $\mathcal{F}$ in $\mathcal{V}$. Then, for any $m \in \mathbb{N}$, $\mathcal{F}_m^{(n_0)}$ is close to $\mathcal{F}^{n_0}$, and the set of its fixed points (denoted by $Z_m^{(n_0)}$) is δ-dense in D. Consequently, for any $n, m \in \mathbb{N}$, the family $\mathcal{F}_n \cup \mathcal{F}_m^{(n_0)}$ satisfies hypotheses (1)–(3) of Lemma 4.3 for bi-Lipschitz constants λ and κ.

Now let $\{B_i\}$ be a topological base of D, so that each B_i is a ball. Fix $i \in \mathbb{N}$, and let $B = B_i$ be a ball of radius r. By the proof of Claim 4.4 (to apply part (a) we consider the families $\mathcal{F}_i^{(n_0)}$, and to apply part (b) we consider the families $\mathcal{F}_i$), there are non-negative integers k and $k(r)$ such that for any $m \in \mathbb{N}$, there is $f \in \mathcal{F}_m^{(n)}$ such that $f(D) \subset B$, where $n = kn_0 + k(r)$.

Using this statement inductively, one finds a dense branch in D for any orbit $\langle\langle \mathbf{F} \rangle\rangle^+(x)$, $x \in D$. This completes the proof of the first part.

If $\mathcal{G}$ is a family of diffeomorphisms of M such that $\mathcal{F} \subset \mathcal{G}$ and $M = \mathcal{G}(D) = \mathcal{G}^{-1}(D)$, then by compactness of M there are finite subsets $\mathcal{G}_\pm \subset \langle \mathcal{G}^{\pm 1} \rangle^+$ such that each $\mathcal{G}_\pm(D)$ covers the manifold M. This property is robust under small perturbation of $\mathcal{G}$. Thus any point in M has iterations (either backward or forward) in D. This allows the above argument to be repeated to show that any orbit $\langle\langle \mathbf{G} \rangle\rangle^+(x)$, $x \in M$, has a branch which is dense in M. In other words, IFS($\mathcal{G}$) is strongly robustly minimal (on M). □

The following result gives a short proof for Proposition 2.2.

PROPOSITION 4.6. *There exist a family of maps $\mathcal{F}$ as in Proposition 2.2 such that* IFS($\mathcal{F}$) *is strongly robustly minimal on some open ball $B_\delta(0)$.*

Proof. Let $\lambda > 0$ such that $B_\lambda(0) \subset D\phi_0(B_1(0))$. We consider a cover of the closed unit ball of $\mathbb{R}^n$ by k balls of radius λ,

$$\overline{B_1(0)} \subset \bigcup_{i=1}^{k} B_\lambda(b_i) = \bigcup_{i=1}^{k} (B_\lambda(0) + b_i).$$

It follows that for $\delta > 0$ small enough,

$$\overline{B_\delta(0)} \subset \bigcup_{i=1}^{k} (\phi(B_\delta(0)) + \delta b_i).$$

To complete the proof, it is enough to apply the previous theorem for $D = B_\delta(0)$ and the family $\mathcal{F} = \{\phi + c_1, \dots, \phi + c_k\}$, where $c_i = \delta b_i$. □

4.3. *Conclusion of proof of Theorem B.*

Proof of Theorem B. It is easy to see that the pair of diffeomorphism (in both proofs) of Theorem A satisfies the hypotheses of Theorem 4.2 and so it generates a strongly robustly

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

minimal IFS. The same occurs for the one-dimensional case. In fact, on the circle the pair of diffeomorphisms (one is a Morse–Smale diffeomorphism close to identity and the other one is an irrational rotation) in [**8**] also satisfies the hypotheses of Theorem 4.2. □

5. *Skew products and robust transitivity*

Following the ideas developed in [**14**], we discuss relations between robustly minimal IFSs constructed in the previous sections and blenders. We demonstrate how, through these relations, the minimal IFSs provide elementary constructions of robustly transitive skew products. More precisely, we show how the constructions in this paper provide an alternative construction for one of the main results from [**3**] on robustly transitive skew products (see Theorem 5.2 below).

The material in the previous sections is readily translated to skew product systems of diffeomorphisms over shift maps; a blending region will give rise to a symbolic blender. Blenders are certain hyperbolic invariant sets used in the construction of robustly transitive dynamics, as explored in [**3**] (see also [**4, 5**] and the survey [**15**]). The original notion of blenders is in a context of partial hyperbolicity with one-dimensional central directions. The generalization of blenders to higher-dimensional central directions was first considered in [**13, 14**], where symbolic blenders and their geometric models are introduced and applied to robustly transitive dynamics in symplectic settings. A further study of their geometric and dynamical properties is in [**1**].

5.1. *Symbolic blenders.* Let g_1, g_2 be two diffeomorphisms on M generating a robustly minimal IFS as in the previous sections. For reasons of definiteness and clarity we assume the properties listed in §3.2. Write $h_1 = g_1$, $h_2 = g_1 \circ g_2$ and let $\mathcal{H} = \{h_1, h_2\}$. Recall that IFS($\mathcal{H}$) acts minimally on a set Δ. More specifically, there are open sets $E_{\text{in}} \subset \Delta \subset E_{\text{out}}$ on which

$$E_{\text{in}} \subset \mathcal{H}(E_{\text{in}}) \subset \Delta \subset \mathcal{H}(E_{\text{out}}) \subset E_{\text{out}} \tag{5}$$

and h_1 and h_2 are contractions on E_{out}.

With $\Sigma = \{1, 2\}^{\mathbb{Z}}$, let H_0 on $\Sigma \times M$ be defined by

$$H_0(\omega, y) = (\sigma(\omega), h_{\omega_0}(y)),$$

where $(\sigma(\omega))_k = \omega_{k+1}$ is the left shift operator. For the one-sided symbol space $\Sigma_+ = \{1, 2\}^{\mathbb{N}}$, a map $H_{0,+}$ on $\Sigma_+ \times M$ is defined likewise, using the same formula. The symbol spaces Σ_+ and Σ will be endowed with the product topology, and we write $\Sigma = \Sigma_- \times \Sigma_+$.

A natural situation is where Σ occurs, through a topological conjugacy, as an invariant set of a diffeomorphism. This gives sense to notions of (partial) hyperbolicity and of invariant manifolds such as stable manifolds. It will in fact be the context of our applications on robustly transitive skew product systems. Henceforth, we work in the setup of diffeomorphisms $f : N \to N$ on a compact manifold N possessing a maximal hyperbolic invariant set in an open set $U \subset N$ on which f is topologically conjugate to σ on Σ. We consider skew product systems $(x, y) \mapsto (f(x), g_y(x))$ on $N \times M$; write $\mathcal{S}^1(N, M)$ for the set of such diffeomorphisms endowed with the C^1 topology. Note that C^1 small perturbations of f yield a perturbed hyperbolic set in U on which the

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

dynamics remains topologically conjugate to σ acting on Σ. For convenience, as we restrict diffeomorphisms to their hyperbolic sets, we write $\mathcal{S}^1(\Sigma, M)$ for the C^1 skew product maps $H : \Sigma \times M \to \Sigma \times M$,

$$H(\omega, y) = (\sigma(\omega), h_\omega(y)),$$

endowed with the C^1 topology in the sense just described.

Suppose H is partially hyperbolic on $\Sigma \times M$ in the sense that contraction and expansion rates of σ dominate those of $y \mapsto h_\omega(y)$. That is, with $E^s \oplus E^u$ denoting the splitting in stable and unstable directions for σ,

$$|D\sigma(\omega)|_{E^s_\omega}| < \mathrm{m}(Dh_\omega(y)) \le |Dh_\omega(y)| < \mathrm{m}(D\sigma(\omega)|_{E^u_\omega})$$

for each $y \in M$ and $\omega \in \Sigma$. The partially hyperbolic skew product systems in $\mathcal{S}^1(\Sigma, M)$ form an open subset of $\mathcal{S}^1(\Sigma, M)$; we may assume that H_0 is partially hyperbolic. We refer to [**5**] for more on partial hyperbolicity. As a consequence of partial hyperbolicity, there are strong stable and strong unstable manifolds of points in $\Sigma \times M$ that project to the stable and unstable manifolds of σ under the projection to the base space Σ. These strong stable manifolds provide a strong stable lamination $\mathcal{F}^{ss}$ of $\Sigma \times M$. For definiteness, a local strong stable manifold is a compact part of a strong stable manifold that projects to some $\Sigma_- \times \{\omega_+\}$ under the projection to the base space Σ. Likewise, strong unstable manifolds of points in $\Sigma \times M$ provide a strong unstable lamination $\mathcal{F}^{uu}$ of $\Sigma \times M$. Local strong unstable manifolds project to $\{\omega_-\} \times \Sigma_+$ under the projection to the base space Σ. A strong stable or unstable lamination of a set is called minimal if each of its leaves lies dense in the set.

The maximal invariant set B of H in $\Sigma \times E_{\mathrm{out}}$ in the following proposition is called a symbolic blender [**14**]. The local unstable set of B is to be interpreted as the union of local strong unstable manifolds through points in B.

PROPOSITION 5.1. *Any skew product map $H \in \mathcal{S}^1(\Sigma, M)$ sufficiently close to $H_0(\omega, m) = (\sigma(\omega), h_{\omega_0}(m))$ possesses a maximal invariant set $B \subset \Sigma \times E_{\mathrm{out}}$ with the following properties:*

- *H is topologically mixing on B;*
- *the strong unstable lamination of B is minimal;*
- *any local strong stable manifold inside $\Sigma \times E_{\mathrm{in}}$ intersects the local unstable set of B.*

Proof. The image $H^k_{0,+}(\Sigma_+ \times E_{\mathrm{in}})$ contains 2^k strips $\Sigma_+ \times U_i \subset \Sigma_+ \times E_{\mathrm{out}}$ with the diameter of U_i going to 0 as $k \to \infty$ and together covering $\Sigma_+ \times E_{\mathrm{in}}$. Take an open set $U \subset \Sigma_+ \times \Delta$. A high iterate $H^n_{0,+}(U)$ contains a strip $\Sigma_+ \times J$ in $\Sigma_+ \times \Delta$. By the above description of iterates $H^k_{0,+}(\Sigma_+ \times E_{\mathrm{in}})$, we see that further iterates $H^{n+k}_{0,+}(U)$ contain strips in $\Sigma_+ \times \Delta$, that lie increasingly dense in it as $k \to \infty$.

This reasoning also applies to small perturbations H_+ of $H_{0,+}$, where also the fiber maps may depend on all of ω instead of just ω_0 (for one-dimensional central directions this is also pursued in [**10**]). Suppose $H_+(\omega, x) = (\sigma(\omega), h_\omega(x))$ is such that h_ω depends continuously on ω and is uniformly close to h_{ω_0}. We note the following changes in the reasoning. The inclusions (5) get replaced by

$$\Sigma \times E_{\mathrm{in}} \subset H_+(\Sigma_+ \times E_{\mathrm{in}}), \quad H_+(\Sigma_+ \times E_{\mathrm{out}}) \subset \Sigma_+ \times E_{\mathrm{out}}. \tag{6}$$

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

An iterate H_+^n maps $\Sigma_+ \times E_{\text{out}}$ to 2^n strips R_i. The diameter of a strip is the maximal real number r so that each $R_i \cap \{\omega\} \times E_{\text{out}}$ is contained in a ball of radius r. The map H_+ on $\Sigma_+ \times E_{\text{out}}$ acts by contractions in the fibers $\{\omega\} \times E_{\text{out}}$. Hence $H_+^n(\Sigma_+ \times E_{\text{in}})$ are 2^n strips with diameter going to zero as $n \to \infty$ and together covering $\Sigma_+ \times E_{\text{in}}$. Iterates under H_+ of $\Sigma_+ \times E_{\text{in}}$ or $\Sigma_+ \times E_{\text{out}}$ therefore converge to an invariant set that contains $\Sigma_+ \times E_{\text{in}}$. As above, for any open set U, $H_+^{n+k}(U)$ contains strips lying increasingly dense in the invariant set of H_+ as k increases.

We proceed with skew products over the shift operator on two-sided symbol spaces. For H C^1-close to H_0, there is a strong stable foliation close to the affine foliation, that is, with leaves close to $\Sigma_- \times \{\omega_+\} \times \{m\}$. Write $\pi^+ : \Sigma \times E_{\text{out}} \to \Sigma_+ \times E_{\text{out}}$ for the continuous projection along local strong stable manifolds. The existence of a strong stable foliation that is close to the affine foliation means that π^+ is C^0-close to the identity.

We can copy the previous reasoning. Observe that H^n maps a curve $\{\omega_-\} \times \Sigma_+ \times \{m\}$ to 2^n curves that are each a graph of a map $\Sigma_+ \to \Sigma_- \times E_{\text{out}}$. Likewise, H^n maps $\Sigma \times E_{\text{out}}$ to 2^n strips R_i. Since π^+ is close to the identity, (6) gets replaced by

$$\Sigma \times E_{\text{in}} \subset \pi^+ H(\Sigma_+ \times E_{\text{in}}), \quad \pi^+ H(\Sigma_+ \times E_{\text{out}}) \subset \Sigma_+ \times E_{\text{out}}.$$

Invariance of the strong stable foliation gives that these inclusions also hold for iterates of H. Further, by uniform C^1-closeness of h_ω to h_{ω_0},

$$\lim_{i \to \infty} \operatorname{diam}(R_i) = 0.$$

The diameter of a strip, as before, is the maximal real number r such that each $R_i \cap \{\omega\} \times E_{\text{out}}$ is contained in a ball of radius r. This proves the proposition. □

5.2. *Robustly transitive skew product diffeomorphisms.* As an application of symbolic blenders from Proposition 5.1 we show how it gives constructions of robustly transitive diffeomorphisms, obtaining a main result in [**3**] in a straightforward manner.

Let N, M be compact manifolds and let $f : N \mapsto N$ be a diffeomorphism with a compact hyperbolic locally maximal invariant set Λ_f, on which f is topologically mixing. Below we will use the fact that unstable manifolds for f of points in Λ_f lie dense in Λ_f; see, for example, [**12**, §18.3]. Recall that a C^1 small perturbation $\tilde{f}$ of f possesses a hyperbolic attractor $\Lambda_{\tilde{f}}$ near Λ_f. Moreover, $\tilde{f}$ restricted to $\Lambda_{\tilde{f}}$ is topologically conjugate to f restricted to Λ_f. Let $R_1, \ldots, R_k$ be a Markov partition for Λ_f, through which $f|_{\Lambda_f}$ is conjugate by φ to the full shift $\sigma : \Sigma \to \Sigma$ with k symbols.

THEOREM 5.2. [**3**] *There is a diffeomorphism $F : N \times M \to N \times M$, $F(x, y) = (f(x), g(x, y))$ with f as above, that is topologically mixing on $\Lambda_f \times M$. Moreover, F is robustly topologically mixing in $\mathcal{S}^1(N \times M)$; that is, there is an open neighborhood U of F in $\mathcal{S}^1(N \times M)$ such that each $\tilde{F}(x, y) = (\tilde{f}(x), \tilde{g}(x, y))$ from U is topologically mixing on $\Lambda_{\tilde{f}} \times M$.*

Proof. Let $\Omega_f \subset \Lambda_f$ be a Smale horseshoe for an iterate f^k of f, that is, an invariant set for f^k on which f^k is topologically conjugate to the shift σ on the symbol space $\Sigma = \{1, 2\}^{\mathbb{Z}}$ (see, for example, [**12**, Theorem 6.5.5]). For simplicity we assume $k = 1$ for now. With

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

the Smale horseshoe comes a Markov partition of two sets U_1, U_2 covering Ω_f, for which Ω_f is the maximal invariant set in $U_1 \cup U_2$.

Consider a skew product system $(x, y) \mapsto (f(x), g(x, y))$ with the following properties:

(1) through the conjugation of $f|_{\Omega_f}$ with σ, the skew product system on $\Omega_f \times M$ is given by
$$G(\omega, y) = (\sigma(\omega), g_{\omega_0}(y)),$$
where g_1, g_2 are as in §3.2;

(2) for $(x, y) \in U_i \times M$, $i = 1, 2$, $g(x, y)$ does not depend on x.

Such a skew product system exists as g_1 and g_2 are isotopic to the identity.

We now consider the diffeomorphism F restricted to $\Omega_f \times M$. The repelling fixed point q_2 of g_2 in Δ gives a fixed point $(2^\infty, q_2)$ for G that is repelling in $\{2^\infty\} \times M$. This corresponds to a fixed point Q_2 for F that is repelling within its central fiber. Similarly, F has a fixed point P_1 coming from the fixed point $(1^\infty, p_1)$ for G; this fixed point is attracting within its central fiber. Note that $W^u(Q_2)$ is dense in $\Lambda_f \times M$, since unstable manifolds for f are dense in Λ_f. We claim that
$$W^u(Q_2) \subset \overline{W^{uu}(P_1)},$$
establishing that $W^{uu}(P_1)$ lies dense in $\Lambda_f \times M$. Namely, take a point $q \in W^u(Q_2)$ and a neighborhood V of it, iterate backwards and note that $H^{-m}(V)$ intersects $W^{uu}(P_1)$ by the existence of a symbolic blender of F on $\Omega_f \times M$ (compare [**3**, Lemma 1.9]). Density of $W^{uu}(P_1)$ implies that H is topologically mixing on $\Lambda_f \times M$, since iterates of an open set intersect the local stable manifold of P_1 and thus accumulate onto $W^{uu}(P_1)$. This construction is robust under perturbations of F (see also Proposition 5.1), proving that F is robustly topologically mixing. □

5.3. *Sufficient conditions for robust transitivity.* Let $f : N \to N$ be a diffeomorphism with a hyperbolic invariant set Λ_f on which f is topologically conjugate to the shift $\sigma : \Sigma \to \Sigma$ with k symbols. First we state an application of the well-known results of [**9**].

LEMMA 5.3. (Normal hyperbolicity) *Let $F : N \times M \to N \times M$ be a skew product diffeomorphism such that $F(x, y) = (f(x), g(x, y))$ with f as above, and such that $g(x, \cdot)$ is uniformly dominated by $f|_{\Lambda_f}$. Then for any G in a C^1 neighborhood of F in* Diff$(N \times M)$*:*

- *there is a unique continuation Γ_G of $\Gamma_F := \Lambda_f \times M$;*
- *$G|_{\Gamma_G}$ is conjugate by ϕ_G to the C^1 skew product H_G on $\Sigma \times M$,*
$$(\omega, y) \mapsto (\sigma(\omega), h_{G,\omega}(y));$$
- *for any $\omega \in \Sigma$, $\phi_G^{-1}(\{\omega\} \times M)$ is diffeomorphic to M and it is C^1 close to $\phi_F^{-1}(\{\omega\} \times M) = \{\varphi^{-1}(\omega)\} \times M$.*

Definition 5.4. If F satisfies the previous lemma, then we say that F is *robustly transitive on* Γ_F if for any G in some neighborhood of F, Γ_G is a transitive set for G. Analogously, one may define robustly topologically mixing.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

On the other hand, one can prove the following lemma (see [**14**] for a proof).

LEMMA 5.5. *Let* $H : \Sigma \times M \to \Sigma \times M$, $H(\omega, y) = (\sigma(\omega), g_{\omega_0}(y))$, *where* $\mathcal{G} = \{g_i\}$ *is a family of diffeomorphisms such that* IFS($\mathcal{G}$) *is minimal. Then* H *is transitive; and the strong unstable lamination is minimal.*

As a corollary one obtains the following criterion for robust transitivity.

THEOREM 5.6. (Sufficient condition for robust transitivity) *Let* F *be as in Lemma 5.3 and suppose that for* $i = 1, \ldots, k$,

$$F|_{R_i \times M} = f|_{R_i} \times g_i$$

with $\bigcup_{i=1}^{k} R_i = \Lambda_f$ *and such that* IFS($\{g_1, \ldots, g_k\}$) *is strongly robustly minimal. Then* F *is robustly transitive on* Γ_F, *and the strong unstable lamination is robustly minimal in* Γ_F.

We remark that Theorem 4.2 provides a sufficient condition to guarantee the main hypothesis of this theorem—the IFS being strongly robustly minimal. Therefore, Theorems 4.2 and 5.6 together provide a general framework to prove robust transitivity, reformulating to the role of blenders.

Acknowledgements. The second author is grateful to Enrique Pujals for many insightful discussions. He was supported in part by a grant from IPM (No. 88370121).

REFERENCES

[1] P. G. Barrientos, Y. Ki and A. Raibekas. Symbolic blender horseshoe and applications, in preparation, 2012.

[2] V. Bergelson, M. Misiurewicz and S. Senti. Affine actions of a free semigroup on the real line. *Ergod. Th. & Dynam. Sys.* **26** (2006), 1285–1305.

[3] C. Bonatti and L. J. Díaz. Persistent non-hyperbolic transitive diffeomorphisms. *Ann. of Math.* (2) **143** (1996), 357–396.

[4] C. Bonatti and L. J. Díaz. Abundance of C^1-robust homoclinic tangencies. *Trans. Amer. Math. Soc.* **364** (2012), 5111–5148.

[5] C. Bonatti, L. J. Díaz and M. Viana. *Dynamics Beyond Uniform Hyperbolicity (Encyclopaedia of Mathematical Sciences, 102)*. Springer, Berlin, 2005.

[6] F. H. Ghane, A. J. Homburg and A. Sarizadeh. C^1 robustly minimal iterated function systems. *Stoch. Dyn.* **10** (2010), 155–160.

[7] F. H. Ghane and M. Saleh. C^1-robustly minimal IFS with three generators. *Internat. J. Bifur. Chaos Appl. Sci. Engrg.* **20** (2010), 3373–3377.

[8] A. S. Gorodetskiĭ and Yu. S. Il'yashenko. Certain properties of skew products over a horseshoe and a solenoid. *Dynamical Systems, Automata, and Infinite Groups (Proceedings of the Steklov Institute of Mathematics, 231)*. Ed. R. I. Grigorchuk. MAIK Nauka/Interperiodica, Moscow, 2000, pp. 90–112.

[9] M. Hirsch, C. Pugh and M. Shub. *Invariant Manifolds (Lecture Notes in Mathematics, 583)*. Springer, Berlin, 1977.

[10] A. J. Homburg. Circle diffeomorphisms forced by expanding circle maps. *Ergod. Th. & Dynam. Sys.* **32** (2012), 2011–2024.

[11] J. E. Hutchinson. Fractals and self-similarity. *Indiana Univ. Math. J.* **30** (1981), 713–747.

[12] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems. With a Supplement by Anatole Katok and Leonardo Mendoza (Encyclopedia of Mathematics and Its Applications, 54)*. Cambridge University Press, Cambridge, 1997.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press

[13] M. Nassiri. Robustly transitive sets in nearly integrable Hamiltonian systems. *PhD Thesis*, IMPA, 2006.
[14] M. Nassiri and E. Pujals. Robust transitivity in Hamiltonian dynamics. *Ann. Sci. Éc. Norm. Supér.* **45** (2012), 191–239.
[15] E. R. Pujals and M. Sambarino. Homoclinic bifurcations, dominated splitting, and robust transitivity. *Handbook of Dynamical Systems*. Vol. 1B. Elsevier, Amsterdam, 2006, pp. 327–378.
[16] D. Volk. Persistent massive attractors of smooth maps. *Ergod. Th. & Dynam. Sys.* to appear.

https://doi.org/10.1017/etds.2013.34 Published online by Cambridge University Press