Animal (2008), **2:2**, pp 292–297 © The Animal Consortium 2008
doi: 10.1017/S1751731107001450

animal

Tail docking in pigs: acute physiological and behavioural responses

M. A. Sutherland†, P. J. Bryer, N. Krebs and J. J. McGlone

Department of Animal and Food Sciences, Pork Industry Institute, Texas Tech University, Lubbock, TX 79409, USA

(Received 20 July 2007; Accepted 1 November 2007)

*Tail docking of piglets is a routine procedure on farms to control tail-biting behaviour; however, docking can cause an acute stress response. The objectives of this research were to determine the stress responses to tail docking in piglets and to compare two methods of tail docking; cautery iron (CAUT) and the more commonly used blunt trauma cutters (BT). At approximately 6 days of age, piglets were tail docked using CAUT (*n *= 20), BT (*n *= 20) or sham tail docked with their tails remaining intact (CON;* n *= 40). Blood samples were taken prior to tail docking and at 30, 60 and 90 min after tail docking to evaluate the effect of tail docking on white blood cell (WBC) measures and cortisol concentrations. The above experiment was repeated to observe behaviour without the periodic blood sampling, so as not to confound the effects of blood sampling on piglet behaviour. Piglet behaviour was recorded in the farrowing crate using 1 min scan-samples via live observations for 60 min prior to and 90 min after tail docking. Total WBC counts were reduced (*P *> 0.05) among BT and CAUT compared with CON piglets 30 min after tail docking. Cortisol concentrations were higher (*P *< 0.01) among BT compared with CON and CAUT piglets 60 min after tail docking. Cautery and BT-docked piglets spent more (*P *< 0.05) time posterior scooting compared with CON piglets between 0 and 15 min, and 31 and 45 min after tail docking. Piglets tail docked using CAUT and BT tended to spend more (*P *< 0.07) time sitting than CON piglets between 0 and 15 min post tail docking. Elevated blood cortisol can be reduced by the use of the CAUT rather than the BT method of tail docking. Although the tail docking-induced rise in cortisol was prevented by using CAUT, the behavioural response to BT and CAUT docking methods was similar.*

Keywords: animal welfare, behaviour, cortisol, pigs, tail docking

Introduction

Tail biting in pigs is a behaviour that is a serious animal welfare problem. Tail biting affects the welfare of the pigs being bitten directly, also the procedure of tail docking as a common means of preventing tail biting is an animal welfare issue as it causes acute trauma and pain. Currently, the cause of tail-biting outbreaks among pigs is unknown, but the cause appears to be multi-factorial. The onset of tail-biting behaviour has been associated with many factors including physical (i.e. floor type), environmental, nutritional and feeding management, over-crowding, gender, genetics, length of tail and lack of substrates (Fraser, 1987; Fraser and Rushen, 1987; McGlone *et al.*, 1990; McGlone and Nicholson, 1992; Guy *et al.*, 2002; Jankevicius and Widowski, 2003 and 2004; Walker and Bilkei, 2006). Currently, tail docking is routinely used on farms in the USA as a solution to this problem.

† E-mail: mhairi.sutherland@ttu.edu

Commonly used tail-docking techniques include surgical tail docking (the tail is cut off using a sharp knife), heated docking iron (the tail is severed using a cautery iron (CAUT)) and rubber ring tail docking (a constrictive rubber ring is applied to the tail). Physiological and behavioural indices of stress have been used to measure the acute stress response to tail docking in pigs (Noonan *et al.*, 1994; Prunier *et al.*, 2005), lambs (Mellor and Holmes, 1988; Graham *et al.*, 1997 and 2002; Kent *et al.*, 1998) and dairy cows (Petrie *et al.*, 1996; Eicher *et al.*, 2000; Schreiner and Ruegg, 2002). Physiological indices of stress shown to increase in response to tail docking include cortisol concentrations (Mellor and Holmes, 1988; Petrie *et al.*, 1996; Graham *et al.*, 1997; Kent *et al.*, 1998), haematological values (Schreiner and Ruegg, 2002) and haptoglobin concentrations (Eicher *et al.*, 2000). Behaviours shown to change in response to tail docking include eating (Eicher *et al.*, 2000), restlessness (Graham *et al.*, 1997; Kent *et al.*, 1998), foot stamping (Graham *et al.*, 1997; Kent *et al.*, 1998), head turning (Graham *et al.*, 1997; Kent *et al.*, 1998), total active

https://doi.org/10.1017/S1751731107001450 Published online by Cambridge University Press

behaviour (Graham *et al.*, 1997 and 2002; Kent *et al.*, 1998), time spent in abnormal postures (Graham *et al.*, 1997 and 2002; Kent *et al.*, 1998), vocalisation (Noonan *et al.*, 1994), tail wagging (Noonan *et al.*, 1994) and tail jamming (Noonan *et al.*, 1994).

There is limited literature on the acute stress response of pigs to tail docking. Prunier *et al.* (2005) showed that adrenocorticotropin hormone (ACTH), cortisol and lactate did not differ between pigs tail docked using a CAUT- and control-handled pigs, at 1 day of age. However, tail jamming (clamping of tail stump between the hind limbs without side-to-side movement) and wagging, and grunting were shown to be greater in tail-docked pigs compared with control-handled pigs for up to 1 min after tail docking (Noonan *et al.*, 1994). Currently, there is limited literature comparing the stress response caused by different methods of tail docking in pigs, but the acute stress response caused by different methods of tail docking in lambs has been thoroughly evaluated in the literature (Graham *et al.*, 1997; Kent *et al.*, 1998). Graham *et al.* (1997) compared the stress response to different methods of tail docking in lambs and found that tail docking using the rubber ring method resulted in a marked increase in cortisol concentrations and active behaviours, whereas tail docking using a heated docking iron produced levels of behaviour and cortisol similar to that of control-handled lambs.

Analgesics and anaesthetics are not routinely used to relieve the pain associated with tail docking on commercial pig farms in the USA or at research or teaching institutions (FASS, 1999). Therefore, it would be beneficial to assess different methods of tail docking to determine methods that may reduce the stress caused by this routine procedure. Pain and distress can be assessed in animals using a range of physiological and behavioural indices and the assessment of pain or distress is improved by using several indices (Molony and Kent, 1997). Therefore, the objective of this study was to determine whether different methods of tail docking (conventional blunt trauma cutting (BT) *v.* CAUT) would have an effect on the stress caused by tail docking as measured by behaviour and physiological measures of stress that have previously been shown to change in response to tail docking.

Material and methods

Pigs used in this study were PIC USA genetics using the Camborough-22 sow line. All animals were fed a diet to meet or exceed National Research Council nutrient requirements. Water was provided *ad libitum*. All animal procedures were approved by the Texas Tech University Animal Care and Use Committee.

Experiment 1: physiological response to tail docking

Eight weight-matched and healthy piglets from 10 sows were allocated to one of two treatment groups: docked ($n = 40$) and non-docked (CON, $n = 40$). Within each litter, two gilts and two barrows were allocated to the CON treatment group. Of the four pigs (two gilts and two barrows) allocated to the tail-docking treatment, one gilt and one barrow were allocated to one of two docking treatments: BT or CAUT. The same number of gilts and barrows were allocated to each treatment.

Within the first 3 days after farrowing, piglets were routinely castrated, ear notched (for pig ID) and given an iron shot (100 mg). Piglets were given 3 days to recover from this initial processing experience. At 6 days of age (± 2 days), piglets were tail docked or left intact, depending on which treatment group the pig was in. Piglets were removed from the sow individually and taken to an adjoining room separated by a closed door, so as not to disturb the remaining sows and piglets in the farrowing room. Piglets were held by one handler with the tail facing outward. The second handler marked a length of 2 cm on the pigs' tail starting at the base on the pigs' tail and then either sham cut the pigs' tail or cut the tail using BT or CAUT. Sham cutting involved placing two fingers, one on either side of the tail, and making a cutting motion on the tail. Tail docking was performed using one of two methods: (1) conventional BT cutting with disinfected stainless-steel cutting pliers or (2) cutting with a commercial cutting CAUT (Meador TNSC; Meador Swine Health Developers, Gretna, NE, USA). Regardless of the tail-docking method (BT or CAUT), tails were docked to a tail length of 2 cm. Once all the piglets from one litter were tail docked or sham handled, they were returned to their home pen all together. The tail-docking treatment was randomised over time.

Prior to (baseline), and 30, 60 and 90 min after tail docking, pigs were held in a supine position and 5 ml blood was obtained by anterior vena cava puncture (catching and blood sampling took $\sim$1 min). Blood was collected into vacutainers containing sodium heparin. Whole blood was analysed to determine white cell counts, differential leucocyte counts and platelet counts using a cell counter (Cell-Dyn® 3700, Abbott laboratories, Abbott Park, IL, USA) and the neutrophil-to-lymphocyte (N : L) ratio was calculated by dividing the percentage of neutrophils by the percentage of lymphocytes. Blood samples were centrifuged and plasma collected for analysis of cortisol using an enzyme immunoassay kit (Assay designs, Ann Arbor, MI, USA). Intra- and inter-assay CV were 4.7% and 6.0%, respectively.

Experiment 2: behavioural response to tail docking

Eight weight-matched and healthy piglets from 10 sows were allocated to one of two treatment groups: docked ($n = 40$) and non-docked (CON, $n = 40$). Within each litter, two gilts and two barrows were allocated to the CON treatment groups. Of the four pigs (two gilts and two barrows) allocated to the tail-docking treatment, one gilt and one barrow were allocated to one of two docking treatments: BT or CAUT. The same number of gilts and barrows were allocated to each treatment.

Within the first 3 days after farrowing, piglets were routinely castrated, ear notched (for pig ID) and given an iron shot (100 mg). Piglets were given 3 days to recover

https://doi.org/10.1017/S1751731107001450 Published online by Cambridge University Press

Table 1 *Description of behaviours*

Behaviour	Description
Walking[†]	Relatively low-speed locomotion in which propulsive force derives from the action of legs
Sitting[†]	Resting on the caudal part of the body
Standing[†]	Assuming or maintaining an upright position on extended legs
Lying, without contact	Maintaining a recumbent position and not in contact with other piglets or the sow
Lying, with contact	Maintaining a recumbent position while contacting another piglet/s or the sow
Massaging[†]	Rhythmic and sustained mechanical manipulation of the mammary of the sow by the piglets prior to and after nursing
Nursing[†]	The act of releasing milk to suckling young
Aggressive interactions[‡]	Attacks between two or more piglets including bites and pushes, primarily of the ears and neck
Scooting	Caudal part of the body being dragged across the ground
Total active	All behaviours combined, with the exception of the lying behaviour
Total resting	Lying without contact plus lying with contact

[†]Hurnik *et al.* (1995).
[‡]McGlone (1985).

from this initial processing experience. At 6 days of age (±2 days), piglets were tail docked or left intact, depending on which treatment group the pig was in. One and a half hours prior to tail docking, experimental piglets were individually marked with a heavy-duty marking pen (Super Mark Pen, Fearing International Ltd, Northampton, UK) using a series of lines in the cross sectional plan to differentiate among individual pigs for easy identification during the live observations. After 60 min of recording piglet behaviour, all piglets from one sow were removed and taken to an adjoining room separated by a closed door, so as not to disturb the remaining sows and piglets in the farrowing room. Piglets were held by one handler with the tail facing outward. The second handler marked a length of 2 cm on the pigs' tail starting at the base on the pigs' tail and then either sham cut the pigs' tail or cut the tail using BT or CAUT. After tail docking, all piglets were retuned to their home pen at the same time and the behaviour of each individual pig was recorded using 1-min scan-samples (live observations) for 90 min. The observer sat directly behind the sow to prevent disturbing her as much as possible but still giving the observer a complete view of all piglets in the farrowing crate. Behaviour was recorded in the same way prior to and after tail docking. Behaviours measured included lying alone, lying touching the sow or other piglets, massaging, nursing, scooting, aggressive encounters, standing, sitting and walking (Table 1). All behaviours were mutually exclusive.

Statistical analysis

All data were tested for constant variance and departures from normal distribution. Data lacking normality were transformed logarithmically using log10. Data were subjected to analysis of variance using the mixed-model procedure of Statistical Analysis Systems Institute (SAS, 2004). All analyses were performed as two-tail tests. The piglet was the experimental unit. Litter was the block. The study was a random complete block design with three treatments (BT, CAUT and CON). For physiological measures, the main fixed effects were gender (two levels), treatment (three levels) and time (four levels). The random effects were litter (10 levels) and piglet (80 levels). The interactions between treatment and time (d.f. = 6), and treatment and litter (d.f. = 18), were included in the model. The model had a repeated structure on time allowing, incorporation of heterogeneity of variances across time. A total of 10 litters and 80 piglets (CON, $n = 40$; BT, $n = 20$; CAUT, $n = 20$) were examined. Behaviour data were recorded as the percentage of observations of a given behaviour over the entire sampling period and analysed using analysis of variance using the mixed-model procedure of SAS. The 150 min behaviour observation period was divided into 10- and 15-min periods. For behavioural measures, the main fixed effects were gender (two levels), treatment (three levels) and period (10 levels). The random effects were litter (10 levels) and piglet (80 levels). The interactions between treatment and period (d.f. = 18), and treatment and litter (d.f. = 18) were included in the model. The model had a repeated structure on time, allowing incorporation of heterogeneity of variances across time. A total of 10 litters and 80 piglets (CON, $n = 40$; BT, $n = 20$; CAUT, $n = 20$) were examined.

Results

Experiment 1: physiology

Total white blood cell (WBC) counts differed among treatments over time (Figure 1). Total WBCs were lower ($P < 0.05$) among BT and CAUT piglets compared with CON piglets 30 min after tail docking. Total WBCs counts were lower ($P < 0.01$) among BT and CAUT piglets at 30 min compared with baseline (0 min) counts but returned to baseline values by 60 min after tail docking. Total WBC counts of CON piglets did not change ($P > 0.05$) over time. No litter-by-treatment interactions were found.

WBC measures differed over time ($P > 0.01$) regardless of treatment (Table 2). The number and percentage of neutrophils increased ($P > 0.01$) over time. Conversely, the number and percentage of lymphocytes decreased ($P > 0.01$) over time. The neutrophil-to-lymphocyte ratio increased ($P > 0.005$) over time. No treatment or litter-by-treatment interactions were found.

https://doi.org/10.1017/S1751731107001450 Published online by Cambridge University Press

Cortisol concentrations differed among treatments over time (Figure 2). Cortisol concentrations were higher ($P < 0.05$) among BT-docked piglets compared with CON and CAUT piglets 60 min after tail docking. Cortisol concentrations did not differ ($P > 0.05$) between CAUT and CON piglets at this time. No other differences ($P > 0.05$) in cortisol concentrations were observed among treatments at 0, 30 or 90 min. No litter-by-treatment interactions were found.

Experiment 2: behaviour

Cautery and BT piglets spent more ($P < 0.001$) time scooting compared with CON piglets between 0 and 15 min post tail docking (Figure 3). Piglets docked using CAUT and BT spent more time ($P < 0.05$) scooting compared with CON piglets between 31 and 45 min post tail docking.

Prior to tail docking, BT spent more ($P < 0.01$) time sitting and CAUT tended ($P < 0.07$) to spend more time sitting compared with CON piglets. Piglets tail docked using CAUT spent more ($P < 0.01$) time sitting and BT-docked piglets tended ($P < 0.08$) to spend more time sitting than CON piglets between 0 and 15 min post tail docking (Figure 4).

No other treatment, period-by-treatment or litter-by-treatment interactions were found for the other behaviours measured in this study.




Figure 1 Total white blood cell (WBC) counts ($10^3/\mu l$) of piglets tail docked using cautery (CAUT; ◆), blunt trauma cutting (BT; ▲) or sham-docked controls (CON; ■) at 0, 30, 60 and 90 min after tail docking or control handling. At each time, least-square means accompanied by an * are different at $P < 0.05$.

Discussion

This study was designed to assess the acute physiological and behavioural response of piglets to tail docking. Furthermore, we wanted to compare two methods of tail docking to determine whether one method would cause less stress compared with the other. Tail docking by the conventional BT method was shown to cause an acute cortisol and behavioural response in piglets. However, tail




Figure 2 Cortisol concentrations (ng/ml) of piglets tail docked using cautery (CAUT; ◆), blunt trauma cutting (BT; ▲), or sham-docked controls (CON; ■) at 0, 30, 60 and 90 min after tail docking or control handling. At each time, least-square means accompanied by an * are different at $P < 0.05$.




Figure 3 The percentage of time spent performing scooting behaviour by piglets tail docked using cautery (CAUT), blunt trauma cutting (BT) or sham-docked controls (CON) over a 150-min period. At each time, least-square means accompanied by subscripts are different at $P < 0.07$.

Table 2 *Comparison of white blood cell values over time of piglets tail-docked using cautery (CAUT), blunt trauma cutting (BT) or sham-docked controls (CON)*

	Time (min after tail docking or sham)				
	0	30	60	90	*P* value
n	80	80	80	80	
Neutrophil count ($10^3/\mu l$)	4.3 ± 0.29^{a}	4.5 ± 0.24^{b}	5.2 ± 0.25^{b}	5.4 ± 0.35^{b}	0.01
Lymphocyte count ($10^3/\mu l$)	9.8 ± 0.57^{a}	7.5 ± 0.76^{b}	7.74 ± 0.65^{b}	6.81 ± 0.78^{b}	0.01
Neutrophils (%)	31.6 ± 2.18^{a}	41.1 ± 2.22^{b}	43.1 ± 2.25^{bc}	47.9 ± 2.63^{c}	0.001
Lymphocytes (%)	64.6 ± 2.32^{a}	55.5 ± 2.50^{b}	52.7 ± 2.51^{bc}	48.2 ± 2.78^{c}	0.001
Neutrophil : lymphocyte ratio	0.7 ± 0.07^{a}	1.0 ± 0.10^{b}	1.1 ± 0.10^{b}	1.3 ± 0.13^{b}	0.005

[a,b,c,d]Least-square means with different superscripts within a row are different ($P < 0.05$).

https://doi.org/10.1017/S1751731107001450 Published online by Cambridge University Press




Figure 4 The percentage of time spent performing sitting behaviour by piglets tail docked using cautery (CAUT), blunt trauma cutting (BT), or sham-docked controls (CON) over a 150-min period. At each time, least-square means accompanied by subscripts are different at $P < 0.08$.

docking using a CAUT reduced the cortisol response to tail docking compared with the conventional method using BT.

Total WBC counts declined in piglets tail docked using BT or CAUT 30 min post tail docking. Reduction in the total WBC numbers in piglets experiencing tail-docking stress may be the result of leucocyte trafficking. Dhabhar and McEwen (1997) hypothesised that during stress leucocytes move out of the peripheral blood and into the skin, lymph nodes and bone marrow to prepare the animal for a potential assault. The number of lymphocytes decreased immediately after a 2 h restraint test, and after 30 s of handling in mice (Bowers *et al.*, 2007), suggesting that leucocyte trafficking can occur quickly in response to an acute stress. Leucocyte trafficking in response to tail docking in pigs may potentially benefit the piglet by preparing its body for possible future stressors.

Cortisol concentrations were influenced by tail docking over time. At 60 min after tail docking, cortisol concentrations were higher in BT piglets compared with CON and CAUT piglets. The cortisol response to tail docking was similar between CON and CAUT piglets. Prunier *et al.* (2005) also showed that cortisol concentrations did not differ between piglets tail docked using a heated docking iron and control-handled piglets for up to 180 min after tail docking. ACTH was also similar between piglets tail-docked using a heated docking iron and non-docked control piglets (Prunier *et al.*, 2005). The stress response to different tail-docking methods has been compared in lambs and lambs tail-docked using a heated docking iron produced similar cortisol and behaviour (total active behaviours) levels as control-handled lambs (Graham *et al.*, 1997). Intense heat associated with the CAUT may cause third-degree burns and thereby destroy the nociceptors in the immediate area, thereby reducing the perception of pain experienced by these animals. It would be interesting to determine whether these CAUT piglets experience more distress later on when the nociceptors regenerate compared with piglets docked using the BT method. Neuromas were found to be present in the tail stump of docked pigs (Simonsen *et al.*, 1991) and tail-docked heifers showed increased sensitivity to heat and cold (Eicher *et al.*, 2006), suggesting that these animals may experience increased sensitivity to pain and chronic discomfort due to tail docking. However, the present study was designed to determine the acute physiological and behavioural effects of tail docking, not the long-term consequences. It has been suggested that cautery may delay wound healing, which could possibly lead to chronic infections (Graham *et al.*, 1997). However, in an unpublished study carried out in this laboratory, wound healing was assessed daily in piglets tail docked using CAUT or BT up to the time they were moved into the nursery. Wound healing did not differ between these two methods over time. Therefore, if tail docking using CAUT reduces the acute stress response to tail docking and has no long-term detrimental effects on the health or pain and/or discomfort experienced by the piglets, this may be a practical alternative to reduce the stress caused by this procedure.

Behaviour of individual piglets was recorded prior to and after tail docking. Tail-docked piglets spent more time scooting than CON piglets. It appeared as though piglets performed this behaviour to relieve themselves of the discomfort caused by having their tails removed as scooting behaviour is not a natural behaviour performed by piglets and was not recorded prior to tail docking or by CON piglets. Tail-docked piglets performed scooting behaviour for up to 45 min after tail docking, but no scooting behaviour was recorded after 45 min among tail-docked piglets (regardless of the tail-docking method). Ninety minutes after tail docking, cortisol concentrations were similar among CON, CAUT and BT pigs. The absence of scooting behaviour in combination with CON level cortisol concentrations in tail-docked pigs 90 min after tail docking suggests that these animals experienced transient stress due to tail docking, which was minimal 90 min after tail docking. However, these animals may experience chronic pain or stress due to increased sensitivity in the tail stump owing to neuroma formation (Simonsen *et al.*, 1991; Eicher *et al.*, 2006). Further studies are needed to determine whether tail docking of piglets results in chronic pain or stress due to this procedure.

Piglets tail-docked using CAUT spent more time sitting than CON and BT pigs, up to 15 min after tail docking. Sitting is a natural behaviour performed by pigs, but in the situation of tail-docked piglets, sitting may have been carried out as a means to relieve the sensation caused by the wound of tail docking or to protect the wounded site. Sitting behaviour was also performed by CAUT and BT piglets prior to tail docking; therefore, it is difficult to determine whether sitting behaviour performed after tail docking by CAUT and BT pigs was in response to having been tail docked or was simply a natural behaviour. The major behavioural changes (tail wagging, tail jamming and vocalisation) in response to tail docking in piglets occurred in the first minute after tail docking (Noonan *et al.*, 1994). In the present study, behavioural observations started when the piglets were returned to the home pen and piglets were only returned to the home pen once all the piglets from that

https://doi.org/10.1017/S1751731107001450 Published online by Cambridge University Press

litter had been tail docked; for this reason, the initial behavioural response to tail docking was not recorded. It appears as though the optimum behavioural indices of tail docking stress in piglets are tail wagging, tail jamming, posterior scooting and vocalisation, which occur more frequently within the first few minutes after tail docking.

Tail docking is a routine management practice carried out on pig farms to control tail biting. Analgesics and anaesthetics are not routinely used for pain relief for tail docking on commercial pig farms in the US. Therefore, it would be beneficial to the welfare of piglets to determine other methods of tail docking that may reduce the acute stress caused by this procedure. Behaviour appeared to be similar between piglets tail docked using CAUT and BT cutting after tail docking; however, cortisol concentrations were lower in CAUT piglets (in fact similar to CON piglets) than in piglets docked using BT cutting at 60 min post tail docking. Tail docking using a heated CAUT may reduce the acute physiological stress response caused by tail docking as compared with conventional tail docking using BT cutting. Behaviour did not differ between piglets tail docked using the conventional BT or CAUT docking method. Further studies are needed to confirm these findings and to determine the possible chronic stress caused by tail docking in piglets.

Acknowledgements

This research was supported by the National Pork Board, Des Moines, IA, USA. The authors thank Lindsey Hulbert, Matt Massey and Mila Billette for their assistance with this project.

References

Bowers SL, Bilbo SD, Dhabhar FS and Nelson RJ 2007. Stressor-specific alterations in corticosterone and immune responses in mice. Brain, Behavior, and Immunity (in press).

Dhabhar FS and McEwen BS 1997. Acute stress enhances while chronic stress suppresses cell-mediated immunity *in vivo*: a potential role for leukocyte trafficking. Brain, Behavior, and Immunity 11, 286–306.

Eicher SD, Morrow-Tesch JL, Albright JL, Dailey JW, Young CR and Stanker LH 2000. Tail-docking influences on behavioral, immunological, and endocrine responses in dairy heifers. Journal of Dairy Science 83, 1456–1462.

Eicher SD, Cheng HW, Sorrells AD and Schutz MM 2006. Behavioral and physiological indicators of sensitivity or chronic pain following tail docking. Journal of Dairy Science 89, 3047–3051.

Federation of Animal Science Societies 1999. Guide for the care and use of agricultural animals in agriculture and teaching, 1st revised edition FASS, Savoy, IL, USA.

Fraser D 1987. Mineral deficient diets and the pig's attraction to blood: implications to tail-biting. Canadian Journal of Animal Science 67, 909–918.

Fraser D and Rushen J 1987. Attraction to blood as a factor in tail-biting by pigs. Applied Animal Behaviour Science 17, 61–68.

Graham MJ, Kent JE and Molony V 1997. Effects of four analgesic treatments on the behavioural and cortisol response of 3-week-old lambs to tail docking. Veterinary Journal 153, 87–97.

Graham MJ, Kent JE and Molony V 2002. The influence of the site of application on the behavioural responses of lambs to tail docking by rubber ring. Veterinary Journal 164, 240–243.

Guy JH, Rowlinson P, Chadwick JP and Ellis M 2002. Behaviour of two genotypes of growing–finishing pig in three different housing systems. Applied Animal Behaviour Science 75, 193–206.

Hurnik JF, Webster AB and Siegel PB 1995. Dictionary of farm animal behavior. Iowa State University Press, Ames.

Jankevicius ML and Widowski TM 2003. Does balancing for color affect pigs' preference for different flavored tail-models? Applied Animal Behaviour Science 84, 159–165.

Jankevicius ML and Widowski TM 2004. The effect of ACTH on pigs' attraction to salt or blood-flavored tail-models. Applied Animal Behaviour Science 87, 55–68.

Kent JE, Molony V and Graham MJ 1998. Comparison of methods for the reduction of acute pain produced by rubber ring castration or tail docking of week-old lambs. Veterinary Journal 155, 39–51.

McGlone JJ 1985. A quantitative ethogram of aggressive and submissive behaviors in recently regrouped pigs. Journal of Animal Science 61, 559–565.

McGlone JJ and Nicholson RI 1992. Effects of limited floor and feeder space on pig performance and tail biting. Texas Tech University Agricultural Sciences Technical Report T-5-317, 26–27.

McGlone JJ, Sells J, Harris S and Hurst RJ 1990. Cannibalism in growing pigs: effects of tail docking and housing system on behavior, performance and immune function. Texas Tech University Agricultural Sciences Technical Report. T-5-283, 69–71.

Mellor DJ and Holmes RJ 1988. Rings versus knife for docking and castration of lambs. Australian Veterinary Journal 65, 403–404.

Molony V and Kent JE 1997. Assessment of acute pain in farm animals using behavioral and physiological measurements. Journal of Animal Science 75, 266–272.

Noonan GJ, Rand JS, Priest J, Ainscow J and Blackshaw JK 1994. Behavioural observations of piglets undergoing tail docking, teeth clipping and ear notching. Applied Animal Behaviour Science 39, 203–213.

Petrie NJ, Mellor DJ, Stafford KJ, Bruce RA and Ward RN 1996. Cortisol response of calves to two methods of tail docking used with or without local anaesthetic. New Zealand Veterinary Journal 44, 4–8.

Prunier A, Mounier AM and Hay M 2005. Effects of castration, tooth resection, or tail docking on plasma metabolites and stress hormones in young pigs. Journal of Animal Science 83, 216–222.

Schreiner DA and Ruegg PL 2002. Responses to tail docking in calves and heifers. Journal of Dairy Science 85, 3287–3296.

Simonsen HB, Klinken L and Bindseil E 1991. Histopathology of intact and docked pigtails. British Veterinary Journal 147, 407–412.

Statistical Analysis Systems Institute 2004. SAS/STAT user's guide version 9.1. SAS Institute Inc., Cary, NC, USA.

Walker PK and Bilkei G 2006. Tail-biting in outdoor pig production. Veterinary Journal 171, 367–369.

https://doi.org/10.1017/S1751731107001450 Published online by Cambridge University Press