Seed Science Research (2001) **11**, 335–343 DOI: 10.1079/SSR200190

Intrapopulation variation in *Abutilon theophrasti* seed mass and its relationship to seed germinability

Hameed A. Baloch, Antonio DiTommaso[1]* and Alan K. Watson

Department of Plant Science, Macdonald Campus of McGill University, Ste-Anne-de-Bellevue, Québec, Canada H9X 3V9

Abstract

The relationship between seed mass variation and germinability in ten *Abutilon theophrasti* Medic. plants from a single agricultural population was examined under controlled environmental conditions. Seeds were collected in the autumn of 1995 and dry-stored in paper bags at 4°C until the start of trials in the spring of 1998. For each plant, between 549 and 1000 randomly selected seeds were separated into seven mass fractions based on individual seed mass, with classes ranging from <6.0 to >11.0 mg. Subjecting the seeds to a cold stratification (4°C) period for 7 d followed by a 21-d alternating day/night (25/14°C) temperature and 14-h photoperiod regime, resulted in 75% overall germination, 24% dormancy and 1% non-viable seeds. The majority of the seeds germinated within 7 d of being exposed to the alternating temperature/light regimes. There was a significant ($P < 0.001$) difference in mean seed mass between the ten plants, with mean mass ranging from 8.8 to 9.6 mg. For nine of the ten plants, the greatest proportion of seeds occurred in the 9.0–9.9 mg mass fraction, while the lowest proportion of seeds was generally found within the 6.0–6.9 mg mass fraction. Most seeds (96%) having a mass below 6.0 mg were non-viable and, of the viable seeds, none germinated. There was a significant ($P < 0.0001$) relationship between seed mass and total germination, but not rate of germination. Germination peaked for seed mass fractions comprising the greatest proportion of total seeds (8.0–9.9 mg) and was lowest for seeds with high or low mass. There was a weak trend of lower germinability for heavier seeds (>10.0 mg) compared with lighter seeds. Maternal source had a significant effect ($P < 0.05$) on total germination and germination rate. Findings from this study suggest that intrapopulation variation in *Abutilon theophrasti* seed mass and its influence on germinability of seeds may play a significant role in maintaining a variable germination pattern and persistent seed bank in this troublesome annual agricultural weed.

Keywords: *Abutilon theophrasti*, intrapopulation variation, seed bank, seed dormancy, seed germination, seed mass, velvetleaf

Introduction

Seed size plays an important role in determining the successful establishment of individual plants within a population (Roach and Wulff, 1987; Westoby *et al*., 1992; Milberg and Lamont, 1997; Kidson and Westoby, 2000). Seed size variation may be due to trade-offs in resource allocation between seed size and number (Venable, 1992) or as a response to environmental heterogeneity (Janzen, 1977). Intraspecific seed size variation and its relation to seedling survival have been demonstrated in several plant species (Meyer *et al*., 1995; Andersson and Milberg, 1998).

The ability of a seed to germinate is affected by several factors, including genetic origin, age and growing environment of the parent during seed maturation, and seed position on the plant (Bello *et al*., 1995; Dekker *et al*., 1996; Andersson and Milberg, 1998; Wright *et al*., 1999). Germination and dormancy of seeds can also be influenced by seed size, shape and surface orientation (Harper, 1977; Moles *et al*., 2000). Seed dormancy has generally been viewed as an important physiological adaptation to variable, unpredictable environments (Cohen, 1966; Fenner, 1985; Allen and Meyer, 1998).

Intrapopulation variation in seed mass and germination has been examined for a few species (e.g. Harper and Obeid, 1967; Stamp, 1990; Zammit and Zedler, 1990; Evans and Cabin, 1995; Milberg *et al*., 1996). In species exhibiting seed dimorphism, such as

*Correspondence
Tel: 607–254–4702
Fax: 607–255–8615
Email: ad97@cornell.edu
[1] Present address: Department of Crop and Soil Sciences, Cornell University, Ithaca, NY 14853, USA

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

Bidens pilosa and *Cakile edentula*, differences in seed mass correlated with differences in germination (Forsyth and Brown, 1982; Zhang, 1993). The relationship between seed mass and germination in plant species that do not produce dimorphic seeds is not as evident, and often the confounding effect of environmental factors on seed size and germination has further complicated matters (Milberg *et al*., 1996). The relatively few studies that have attempted to relate individual seed size fractions to viability and germination rates have provided inconsistent results, showing both that seeds of greater mass exhibit higher (Zammit and Zedler, 1990) and lower (Stamp, 1990) germination and rates of germination than seeds of lower mass. Milberg *et al*. (1996) reported that there were large variations in germination rate between nine seed mass fractions in *Lithospermum arvense* and *Anchusa arvensis*. Moreover, the relationship between germination and seed mass differed in both species and between populations of the same species. Large seeds of *Lithospermum* always germinated well, whereas small seeds generally had a poor percentage germination. However, for one of the *Anchusa* populations, germination tended to decrease with increasing seed mass. Research by Harper and Obeid (1967) on fibre and oil seed flax cultivars demonstrated that seeds of intermediate size germinated faster than seeds from extreme size classes. Few studies have examined intrapopulation germinability and dormancy characteristics of seeds from different size fractions for species exhibiting strong physical dormancy.

Abutilon theophrasti (velvetleaf) is a large, self-pollinated annual weed infesting maize and soybean crops, particularly in the mid-western and north-eastern regions of the United States, eastern Canada and the eastern Mediterranean regions. Each plant is capable of producing up to 17,000 seeds that exhibit a high degree of physical dormancy due to the presence of a thick, hard seed coat that is impermeable to water (Warwick and Black, 1988; Baskin and Baskin, 1989). Under optimal conditions for germination, hard seeds cannot imbibe water and are unable to germinate. However, physical scarification, resulting from regular cycles of freezing and thawing or from placement in boiling water for brief periods, disrupts the seed coat and increases germination in hard-seeded species (Lacroix and Staniforth, 1964; Zanin *et al*., 1989; Li *et al*., 1999).

In work by Zhang and Hamill (1997), *Abutilon theophrasti* seeds, having a mass ranging from 4.8 to 11.7 mg, did not exhibit significant differences in germination and dormancy. Surprisingly, in that study, only 39% of seeds germinated, while a large portion of seeds (40%) were found to be non-viable. The remaining seeds (21%) were viable but dormant. Given the low viability of seeds in the Zhang and Hamill (1997) study, it was unclear whether the lack of a relationship between seed size and germinability of seeds in *Abutilon theophrasti* is a general feature of this species or is simply specific to the population tested. Moreover, no attempt was made to relate seed size variation to germinability for individual plants. This may be especially important given the numerous instances where within-plant variation in seed size contributes most to the total seed mass variation within a population (Pitelka *et al*., 1983; Thompson, 1984; Milberg *et al*., 1996). To date, the relationship between intrapopulation seed mass variation and seed germinability in *Abutilon theophrasti* is not well understood and requires further study.

Thus, the objectives of this study were: (1) to determine the variability in seed mass, total germination and rate of germination for the seeds of ten *Abutilon theophrasti* plants from a single agricultural population; and (2) to determine the relationship between seed mass, total germination and rate of germination for seeds from these ten plants.

Materials and methods

Mature capsules of *Abutilon theophrasti* were collected in late September 1995 from ten individually marked plants located within a 3 ha maize field at the Macdonald Campus Farm in Ste-Anne-de-Bellevue, Québec, Canada. Plants were selected randomly within the field but were at least 30 m apart from each other. Capsules were collected randomly on individual plants and were considered mature for harvest when their colour had changed from light green to dark black. Under typical growing conditions of south-western Québec, capsules mature from early August until early October. Thus, the late-September collection date permitted the harvesting of capsules that matured over the normal time period for this region. Capsules were dry-stored at 4°C in paper bags until the start of the experiment in spring 1998. Seeds from the ten plants were obtained by separating them from chaff by hand. The total number of seeds collected from each plant ranged from 549 to 1000, depending on the number of seeds produced by each plant and availability at the time of harvest. Seeds from each plant were then weighed individually to the nearest 0.1 mg and grouped into one of seven seed mass fractions. All seeds were placed on moistened filter paper in 9-cm-diameter plastic Petri dishes containing 8 ml of distilled water. No more than 50 seeds per mass fraction were placed within each Petri dish. Petri dishes were subsequently wrapped with Parafilm™ to prevent water evaporation during the experimental period. All seeds within Petri dishes were then subjected to a low-temperature (4°C) moist stratification treatment for 1 week in the dark. Seeds were subjected to this treatment because it simulated

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

the early season, cold, moist soil conditions that seeds would typically experience under Québec conditions in March and April. At the end of the stratification period, seeds were incubated in a controlled environment chamber programmed to provide 25/14°C day/night temperatures, and a 14-h photoperiod at a photosynthetically active radiation (PAR) of 400 μmol m^{-2} s^{-1} using fluorescent lamps. These environmental conditions are typical of south-western Québec field conditions in late May, when the first *Abutilon theophrasti* germination flush generally occurs. The number of germinated seeds having a radicle at least 2 mm long was recorded daily for a period of 21 days, and all germinated seeds were removed from the Petri dishes. At the end of the experiment, all non-germinated seeds were initially examined for viability by pressing them gently with forceps. Seeds that were crushed under this moderate pressure were considered to be dead. The remaining seeds were then placed in boiling water for 10 s, removed and placed on moistened filter paper, and returned to the original controlled environment conditions for a period of 48 h. All seeds that germinated following this additional germination period were characterized as being viable but dormant, while seeds that did not germinate during this 48-h period were considered to be dead. Placement of seeds in boiling water for a short period of time overcomes physical dormancy in this hard-seeded species (Lacroix and Staniforth, 1964; Zanin *et al*., 1989).

The mean time to germination was determined for each plant and for each of the seven seed mass fractions by first multiplying the total number of seeds germinating on the first day by 1, the total number of seeds germinating on the second day by 2, and so on. The average number of days that were required for seeds to germinate was then determined by dividing the sum of these terms by the total number of seeds that germinated (Milberg *et al*., 1996).

Germination, dormancy and viability data were square root transformed to homogenize variances prior to using the GLM analysis procedure in SAS (Cary, NC, USA). The analysis of variance (ANOVA) procedure was modified in order to take into account the potential heterogeneity of variances and lack of independence over time (from day 1 to day 21) of the temporal repeated measures of individual plants and seed mass (Dutilleul, 1998). The model used for the modified univariate ANOVA was derived by adding the time effects plus all the interactions to the terms of the standard ANOVA model and using the REPEATED statement of the GLM procedure. Most seeds below 6.0 mg were non-viable and thus were not included in the data analyses. Relationships between seed mass, germinability and dormancy were determined by both linear and non-linear regression analyses. The Kruskall–Wallis one-way ANOVA on ranks was used to determine differences in mean seed mass of plants.

Results

Seed mass variation between plants

For the ten plants combined, 75% of seeds had a mass within the 8.0–8.9 mg and 9.0–9.9 mg ranges, while only 1.5% of seeds had a mass within the two extreme seed mass fractions of <6.0 mg and >11.0 mg (Fig. 1). The variation in seed mass for each of the ten plants was large, with the greatest differences occurring for the 8.0–8.9 and 9.0–9.9 mg seed mass fractions (Fig. 1). Median seed mass differed significantly ($P < 0.0001$) among the ten plants, with seeds from individual 3 having the lowest median mass (8.8 mg) and seeds from individual 10 having the highest median mass (9.8 mg) (Fig. 2).

Between plant germination characteristics

After the 21-d experimental period, seed germination differed significantly ($P < 0.05$) among the ten plants (Table 1). For all plants and mass fractions combined, 75% of seeds germinated. In contrast, seed germination for individuals 6 and 8 were 57% and 92%, respectively (Fig. 3). Overall, 24% of *Abutilon* seeds did not germinate under the controlled environment conditions used, and 1.0% of seeds were non-viable.

Seventy-five per cent of seeds germinated within the first 7 d of the experiment, regardless of maternal source (Fig. 3). Germination declined sharply for all plants 10 d after the start of the experiment. The rate of germination differed significantly ($P < 0.05$) between plants (Fig. 4). Across all seed mass fractions, the average number of days required for seeds to germinate ranged from 3.2 d in individual 1 to 5.6 d in individual 7, with an overall population mean of 4.6 d. Seeds from individuals 6 and 4 exhibited the greatest and least variation in germination rate across the seed size fractions, respectively.

Relationship between seed mass and germination characteristics

Seed size had a highly significant ($P < 0.0001$) impact on total germination (Table 1). The highest germination levels (>85%) were observed for the 7.0–7.9 and 8.0–8.9 mg seed mass fractions, and the lowest levels for the lightest seed mass fraction (<6.0 mg), in which no seeds germinated. Only 58% of all seeds in the 6.0–6.9 mg seed mass fraction germinated. Most non-viable seeds (96%) had a seed mass below 7.0 mg. A weak positive trend was observed between mean seed mass and the number of

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press




Figure 1. Cumulative frequency distributions of seed mass within seven seed fractions for ten *Abutilon theophrasti* plants collected from a single agricultural population. Curves are based on the individual masses of 1000 seeds, except for plants 1 (898), 5 (900), 8 (549), 9 (713) and 10 (820).




Figure 2. Median (solid line), mean (dotted line) and 25th and 75th percentiles of seed mass for each of ten *Abutilon theophrasti* plants from a single agricultural population. Bars above and below each box indicate the 90th and 10th percentiles, respectively.

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press




Figure 3. Cumulative germination (%) of *Abutilon theophrasti* seeds over a 21-d period for each of ten plants collected from a single agricultural population. A total of 549–1000 seeds were tested for each plant.




Figure 4. Mean number of days to germination for seeds collected from ten *Abutilon theophrasti* plants from a single agricultural population. Bars indicate ±SE of the means, and values above bars indicate the total number of seeds that germinated for each plant.

seeds not germinating when data for all ten plants were combined (Fig. 5).

The average number of days required for germination was not affected by seed mass (data not shown), although heavier seeds (>11.0 mg) did require, on average, 1.5 additional days to germinate. Interestingly, seeds germinating on the first day of the study had a substantially greater mass (9.7 mg) than seeds germinating during the following 12 days (Fig. 6). There was a trend of increasing time to germination for heavier seeds from day 4 to day 14 of the study (Fig. 6).

Discussion

The important differences in seed mass distribution found between the ten *Abutilon theophrasti* plants were likely due to a number of environmental effects rather than genotypic variation (Zhang and Hamill, 1997). It is plausible that the microenvironments in which the ten plants grew differed in their ability to supply essential resources such as nutrients and light. These differences would have resulted in plants producing seeds that were particularly large or small. It is also possible that the variation in seed mass observed among the ten plants may have been due to differential seed filling in seeds harvested from different branch locations on parental plants (Gutterman, 1992). Zhang and Hamill (1996) reported that mean seed weight in *Abutilon theophrasti* plants varied depending on population, maternal environment and their interaction. However, photosynthetic rate and number of leaves and flowers did not differ among the different populations. Baloch (2001) reported that, under favourable growing conditions, seed mass in *Abutilon theophrasti* was influenced by the branch location of capsules on

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

Table 1. Summary of ANOVA for *Abutilon theophrasti* total germination as affected by plant source, seed mass fraction and time of sampling (21 d)[a]. Seeds having a mass below 6.0 mg were omitted from this analysis as no seeds germinated

Source	df[b]	F-value	Probability	Adjusted Prob>F[c] G–G	Adjusted Prob>F[c] H–F
(a) Between-subject effects					
Plant source	9	2.16	0.0433		
Seed mass	5	22.89	<0.0001		
Error	45				
(b) Within-subject effects					
Time	20	28.11	<0.0001	<0.0001	<0.0001
Time × plant source	180	2.41	<0.0001	<0.0001	<0.0001
Time × seed mass	100	1.61	0.0003	0.0213	0.0062
Error	900				

[a]The Greenhouse and Geisser epsilon is a multiplicate factor by which the number of degrees of freedom of the F test for time-related effects are reduced in order to take the autocorrelation and heteroscedasticity over time into account (Dutilleul, 1998).
[b]df = Degrees of freedom.
[c]The G–G (Greenhouse–Geisser) and H–F (Huynh–Feldt) adjustment values are also shown, G–G = Greenhouse–Geisser epsilon = 0.3319; H–F= Huynh–Feldt epsilon = 0.5165.




Figure 5. Relationship between the proportion of seeds not germinating and mean seed mass for seeds collected from ten *Abutilon theophrasti* plants from a single agricultural population. Data for seeds from only six seed mass fractions were used since, for the lowest fraction (i.e. <6.0 mg), most seeds were non-viable and no seeds germinated. Only mass fractions having more than five seeds were used in the analysis, $y = -0.47 + 0.08x$; $R^2 = 0.31$.




Figure 6. Mean mass (±SE) of *Abutilon theophrasti* seeds that germinated on each of the observed dates. No germination occurred after day 17. Data for the ten *Abutilon* plants were combined. Values near solid circles are the total number of seeds that germinated on that day.

parent plants. That is, seeds collected from early maturing capsules (basipetal) had greater mean weights than seeds harvested from late-maturing (acropetal) capsules. Interestingly, under the stressful conditions of fungal infection and high density, capsules harvested from different branch positions had seeds of similar mean weight. Plants grown under these conditions produced fewer capsules having a larger seed mass than plants grown in the absence of disease and at lower density. Vaughton and Ramsey (1998) also reported that mean seed mass in *Banksia marginata* was greater from early produced flowers lower down on the main stem than from later-produced flowers on the upper part of the main stem. Kane and Cavers (1992) noted that mean seed mass in *Panicum miliaceum* decreased over the growing season, both on individual panicles and all inflorescences of the plant. In contrast, Obeso (1993) observed no variation in mean seed mass at different fruit locations in *Asphodelus albus*.

Within the agricultural population sampled, the ten *Abutilon theophrasti* plants exhibited a large degree

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

of variation in the germinability of their seeds (from 57 to 92%), despite a normal distribution in seed mass of the plants (data not shown). Hence, this variability in germination between the plants was likely not due to differences in seed mass frequency distributions. This finding suggests that differences in the maternal growing environment in this agricultural field (e.g. full sun versus shade) may have resulted in seeds from different plants having divergent germination abilities (Fenner, 1991; Bello *et al*., 1995). For instance, Bello *et al*. (1995) found that *Abutilon theophrasti* plants grown under full sunlight produced smaller-sized seeds that were 20% less likely to germinate under optimal environmental conditions than larger seeds produced from shaded plants. It was unclear from their study whether the decrease in seed dormancy for shaded plants resulted in fewer seeds having hard seed coats or whether some other dormancy mechanism was responsible. In the present study, variation in germination ability for the ten *Abutilon theophrasti* plants may also have been due, in part, to the harvesting of mature capsules on different branch positions. Research on other plant species has revealed large variability in germination of seeds collected from different branch positions on a single plant (Gutterman, 1992; Dekker *et al*., 1996). No attempt was made in the present study to harvest mature capsules from similar positions on each of the plants.

The rate of germination for seeds of the ten *Abutilon theophrasti* plants also differed over the 21-d experimental period. There was no obvious trend observed between the average amount of time required for germination and median seed mass for each of the ten plants. For instance, seeds of individual 1 had a median mass of 9.7 mg and required an average 3.2 d to achieve germination, whereas seeds of individual 7 had a similar median mass (9.6 mg) but required nearly 2.5 d more than individual 1 to achieve germination (5.6 d).

Zhang and Hamill (1997) reported no differences between the germination ability of *Abutilon theophrasti* seeds having a mass ranging from 4.8 mg to 11.7 mg. In the present study, there was a weak trend of lower germination ability for seeds having relatively high mass (i.e. >11 mg). The production of relatively large (>11.0 mg) dormant seeds may be an adaptive feature in *Abutilon theophrasti* because it provides a source of seeds for the soil seed bank that will likely produce competitively superior seedlings when the seeds eventually germinate (Roach and Wulff, 1987; Kidson and Westoby, 2000). Although the total number of large seeds (e.g. >11.0 mg) produced by individual plants may be relatively small (19% of the nearly 9000 seeds collected in this study), many of these dormant seeds may be responsible for the annual fluctuations in *Abutilon theophrasti* emergence patterns observed in the field (Warwick and Black, 1988; Cardina and Sparrow, 1997). Given that plants of this species may produce as many as 17,000 seeds per plant (Mitich, 1991), this would translate into approximately 3200 relatively large-sized seeds (>11.0 mg) produced per plant.

Variations in seed germinability and rate of germination between different plant populations of a species have been documented previously (e.g. Evans and Cabin, 1995; Schütz and Milberg, 1997). For example, Kane and Cavers (1992) reported that *Panicum miliaceum* seeds from two populations separated only by a road differed from each other in survival, germination and dormancy. Genetic uniformity within each population and heterogeneity in microenvironment were suggested as possible sources for the observed variations. Similarly, Milberg *et al*. (1996) reported that germinability and germination rate varied significantly between seed mass fractions in *Lithospermum arvense* and *Anchusa arvensis* and for different populations. Variability in parental growing environment was suggested to account for the differences in germination characteristics observed for the various seed mass classes. Hence, caution should be used when interpreting the results of the present study, given that seeds from only one agricultural population and for a single year were used (Andersson and Milberg, 1998).

Seeds of *Abutilon theophrasti* having a mass of less than 7 mg were largely non-viable (96%), although the proportion of these seeds for each of the plants was very low (i.e. 1–2% of the total number of seeds collected). These findings contrast with those of Zhang and Hamill (1997), where nearly 40% of the 1536 *Abutilon theophrasti* seeds collected were non-viable. Differences in growing environment, as well as duration and method of seed storage, may account for the observed disparity in results. That most *Abutilon theophrasti* seeds germinated within 7 d of being subjected to an alternating day/night temperature regime is consistent with other research (Zhang and Hamill, 1997). In the present study, heavier seeds generally required a longer period of time to germinate, although the effect was not significant. It is worth noting, however, that the mean mass of seeds (9.7 mg) germinating on the first day of the experiment was significantly greater than the mean mass of seeds germinating over the following 12 d. Thus, it would be expected that seedlings originating from these heavier seeds and first to emerge would have a competitive advantage over seedlings from lighter seeds that emerge later (Roach and Wulff, 1987; Gutterman, 1992; Lacey *et al*., 1997). However, Zhang and Hamill (1997) suggested that the adaptive advantage of *Abutilon theophrasti* plants produced from large seeds might not necessarily be reflected in the greater morphological and/or

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

physiological performance of these plants, but rather to their enhanced deleterious impact on neighbouring plants. Zhang and Hamill (1996) found a significant interaction between *Abutilon theophrasti* population and cropping system with respect to seed mass and germination rate. Plants produced significantly heavier seeds with higher germination rates when grown with a maize crop as compared with seeds produced from plants grown with a soybean crop. It was suggested that the production of smaller-sized seeds in soybean versus maize was the result of a genetically based discriminative selection under these two crop management systems.

Variation in mean seed mass and germinability for different *Abutilon theophrasti* agricultural populations needs further investigation. Future research should also be directed towards understanding more fully the role that parental growing environment and capsule branch position play in determining seed mass variation and germinability in this troublesome annual weed.

Acknowledgements

H.A.B. is grateful to the Government of Canada for financial support through a Canadian Commonwealth Scholarship. The authors also thank C.L. Mohler, R.F. Masangkay and two anonymous reviewers for their helpful comments on earlier versions of the manuscript.

References

Allen, P.S. and Meyer, S.E. (1998) Ecological aspects of seed dormancy loss. *Seed Science Research* **8**, 183–191.

Andersson, L. and Milberg, P. (1998) Variation in seed dormancy among mother plants, populations and years of seed collection. *Seed Science Research* **8**, 29–38.

Baloch, H.A. (2001) Effects of seed size and a fungal pathogen, *Colletotrichum coccodes*, on population dynamics of velvetleaf (*Abutilon theophrasti* Medic.). PhD Thesis, McGill University, Montreal, Canada.

Baskin, J.M. and Baskin, C.C. (1989) Physiology of dormancy and germination in relation to seed bank ecology. pp. 53–66 *in* Leck, M.A.; Parker, V.T.; Simpson, R.L. (Eds) *Ecology of soil seed bank*. New York, Academic Press.

Bello, I.A., Owen, M.D.K. and Hatterman-Valenti, H.M. (1995) Effect of shade on velvetleaf (*Abutilon theophrasti*) growth, seed production, and dormancy. *Weed Technology* **9**, 452–455.

Cardina, J. and Sparrow, D.H. (1997) Temporal changes in velvetleaf (*Abutilon theophrasti*) seed dormancy. *Weed Science* **45**, 61–66.

Cohen, D. (1966) Optimizing reproduction in a randomly varying environment. *Journal of Theoretical Biology* **12**, 119–129.

Dekker, J., Dekker, B., Hilhorst, H. and Karssen, C. (1996) Weedy adaptation in *Setaria* spp. IV. Changes in the germinative capacity of *S. faberii* (Poaceae) embryos with development from anthesis to after abscission. *American Journal of Botany* **83**, 979–991.

Dutilleul, P. (1998) Incorporating scale in study design and data analysis. pp. 387–425 *in* Peterson, D.L.; Parker, V.T. (Eds) *Ecological scale: Theory and applications*. New York, Columbia University Press.

Evans, A.S. and Cabin, R.J. (1995) Can dormancy affect the evolution of post-germination traits? The case of *Lesquerella fendleri*. *Ecology* **76**, 344–356.

Fenner, M. (1985) *Seed ecology*. London, Chapman & Hall.

Fenner, M. (1991) The effects of the parent environment on seed germinability. *Seed Science Research* **1**, 75–84.

Forsyth, C. and Brown, N.A.C. (1982) Germination of the dimorphic fruits of *Bidens pilosa* L. *New Phytologist* **90**, 151–164.

Gutterman, Y. (1992) Maternal effects on seeds during development. pp. 27–59 *in* Fenner, M. (Ed.) *Seeds: The ecology of regeneration in plant communities*. Wallingford, UK, CAB International.

Harper, J.L. (1977) *Population biology of plants*. London, Academic Press.

Harper, J.L. and Obeid, M. (1967) Influence of seed size and depth of sowing on the establishment and growth of varieties of fiber and oil seed flax. *Crop Science* **7**, 527–532.

Janzen, D.H. (1977) Variation in seed size within a crop of Costa Rican *Micuna andreana* (Leguminosae). *American Journal of Botany* **64**, 347–349.

Kane, M. and Cavers, P.B. (1992) Patterns of seed weight distribution and germination with time in a weedy biotype of proso millet (*Panicum miliaceum*). *Canadian Journal of Botany* **70**, 562–567.

Kidson, R. and Westoby, M. (2000) Seed mass and seedling dimensions in relation to seedling establishment. *Oecologia* **125**, 11–17.

Lacey, E.P., Smith, S. and Case, A.L. (1997) Parental effects on seed mass: Seed coat but not embryo/endosperm effects. *American Journal of Botany* **84**, 1617–1620.

Lacroix, L.J. and Staniforth, D.W. (1964) Seed dormancy in velvetleaf. *Weeds* **12**, 171–174.

Li, X., Baskin, J.M. and Baskin, C.C. (1999) Seed morphology and physical dormancy of several North American *Rhus* species (Anacardiaceae). *Seed Science Research* **9**, 247–258.

Meyer, S.E., Kitchen, S.G. and Carlson, S.L. (1995) Seed germination timing patterns in intermountain *Penstemon* (Scrophulariaceae). *American Journal of Botany* **82**, 377–389.

Milberg, P. and Lamont, B.B. (1997) Seed/cotyledon size and nutrient content play a major role in early performance of species on nutrient-poor soils. *New Phytologist* **137**, 665–672.

Milberg, P., Andersson, L., Elfverson, C. and Regnér, S. (1996) Germination characteristics of seeds differing in mass. *Seed Science Research* **6**, 191–197.

Mitich, L.W. (1991) Intriguing world of weeds. 33. Velvetleaf. *Weed Technology* **5**, 253–255.

Moles, A.T., Hodson, D.W. and Webb, C.J. (2000) Seed size and shape and persistence in the soil in the New Zealand flora. *Oikos* **89**, 541–545.

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

Obeso, J.R. (1993) Seed mass variation in the perennial herb *Asphodelus albus*: sources of variation and position effect. *Oecologia* **93**, 571–575.

Pitelka, L.F., Thayer, M.E. and Hansen, S.B. (1983) Variation in achene weight in *Aster acuminatus*. *Canadian Journal of Botany* **61**, 1415–1420.

Roach, D.A. and Wulff, R.D. (1987) Maternal effects in plants. *Annual Review of Ecology and Systematics* **18**, 209–235.

Schütz, W. and Milberg, P. (1997) Seed dormancy in *Carex canescens*: regional differences and ecological consequences. *Oikos* **78**, 420–428.

Stamp, N.E. (1990) Production and effect of seed size in a grassland annual (*Erodium brachycarpum*, Geraniaceae). *American Journal of Botany* **77**, 874–882.

Thompson, J.N. (1984) Variation among individual seed masses in *Lomatium grayi* (Umbelliferae) under controlled conditions: magnitude and partitioning of the variance. *Ecology* **65**, 626–631.

Vaughton, G. and Ramsey, M. (1998) Sources and consequences of seed mass variation in *Banksia marginata* (Proteaceae). *Journal of Ecology* **86**, 563–573.

Venable, D.L. (1992) Size-number trade-offs and the variation in seed size with plant resource status. *The American Naturalist* **140**, 287–304.

Warwick, S.I. and Black, L.D. (1988) The biology of Canadian weeds. 90. *Abutilon theophrasti*. *Canadian Journal of Plant Science* **68**, 1069–1085.

Westoby, M., Jurado, E. and Leishman, M. (1992) Comparative evolutionary ecology of seed size. *Trends in Ecology and Evolution* **7**, 368–372.

Wright, K.J., Seavers, G.P., Peters, N.C.B. and Marshall, M.A. (1999) Influence of soil moisture on the competitive ability and seed dormancy of *Sinapis arvensis* in spring wheat. *Weed Research* **39**, 309–317.

Zammit, C. and Zedler, P.H. (1990) Seed yield, seed size and germination behavior in the annual *Pogogyne abramsii*. *Oecologia* **84**, 24–28.

Zanin, G., Berti, A. and Sattin, M. (1989) Mais (*Zea mays* L.) *Abutilon theophrasti* Medicus: effetto della durata e del periodo di competizione. *Rivista di Agronomia* **23**, 185–192.

Zhang, J. (1993) Seed dimorphism in relation to germination and growth of *Cakile edentula*. *Canadian Journal of Botany* **71**, 1231–1235.

Zhang, J. and Hamill, A.S. (1996) Responses of *Abutilon theophrasti* to agricultural management systems. *Weed Research* **36**, 471–481.

Zhang, J. and Hamill, A.S. (1997) Seed weight, intraspecific competition, and plant performance in *Abutilon theophrasti*. *Canadian Journal of Botany* **75**, 1614–1620.

Received 14 March 2001
accepted after revision 14 May 2001
© CAB *International* 2001

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press

Seed Dispersal and Frugivory: Ecology, Evolution and Conservation

Edited by **D J Levey**, *Department of Zoology, University of Florida, Gainesville, Florida, USA,* and **W R Silva**, *Departamento de Zoologia, Instituto de Biologia, UNICAMP, Campinas, Brazil* and **M Galetti**, *Departamento de Ecologia, Universidade Estadual Paulista, Sao Paulo, Brazil*

January 2002 **544 pages Hardback**
ISBN 0 85199 525 X
£75.00 (US$140.00)

Readership: Plant and animal ecology, seed science.

Until recently, the production of fruits by plants, their consumption by animals (frugivory) and the relevance of these to seed dispersal have attracted less attention than topics such as pollination biology. However, since the 1970s they have started to gain more prominence and now give rise to more research funding, seminal papers and international symposiums. This book contains chapters adapted from the Third International Symposium-Workshop on Frugivores and Seed Dispersal held in August 2000 in Rio Quente, Brazil.

Contents include:

Historical and Theoretical Perspectives
- Maintenance of tree diversity in tropical forests, *J Terborgh, N Pitman, M Silman, H Schichter, and P Nunez V.*
- Dissemination limitation and the origin and maintenance of species-rich tropical forests, *E W Schupp and T Milleron*
- Assessing recruitment limitation: Concepts, methods, and case studies from a tropical forest , *H C Muller-Landau, et al*
- Have frugivores influenced the evolution of fruit traits in New Zealand? *J M Lord, A S Markey and J Marshall*
- Mechanistic models for tree seed dispersal by wind in dense forests and open landscapes, *R Nathan, H S Horn, J Chave, and S A Levin*
- The role of vertebrates in the diversification of new world mistletoes, *C Restrepo, S Sargent, D J Levey, and D M Watson*

Plant Strategies
- Mistletoes as parasites and seed-dispersing birds as disease vectors: current understanding, challenges, and opportunities, *J E Aukema and C Martínez del Rio*
- Secondary metabolites of ripe fleshy fruits: ecology and phylogeny in the genus Solanum, *M L Cipollini, et al*
- The seed dispersers and fruit syndromes of Myrtaceae in the Brazilian Atlantic Forest, *M A Pizo*
- Are plant species that need gaps for recruitment more attractive to seed-dispersing birds and ants than other species? *C C Horvitz, et al*
- The role of fruit traits in determining fruit removal in east Mediterranean ecosystems, *I Izhaki*
- Seed dispersal of mimetic fruits: parasitism, mutualism, or exaptation? *M Galetti*
- Secondary dispersal of Jeffery pine seeds by rodent scatter hoarders: the roles of pilfering, recaching, and variable environment, *S B VanderWall*
- The role of seed size in dispersal by a scatterhoarding rodent, *P A Jansen, et al*
- Mast seeding and predator-mediated indirect interactions in a forest community: evidence from post- dispersal fate of rodent-generated caches, *K Hoshizaki and P E Hulme*

Animal Strategies
- Seasonality of fruiting and food hoarding by rodents in Neotropical forests: consequences for seed dispersal and seedling recruitment *P Forget, D S Hammond, T Milleron, and R Thomas*
- Seed eaters: seed dispersal, destruction, and demography, *P E Hulme*
- Plant-animal co-evolution: Is it thwarted by spatial and temporal variation in animal foraging? *C A Chapman and L J Chapman*
- The frugivorous diet of the maned wolf, *Chrysocyon brachyurus* in Brazil: ecology, and conservation, *J C Motta jnr and K Martins*

Consequences of Seed Dispersal
- Frugivore-generated seed shadows: A landscape view of demographic and genetic effects, *P Jordano and J A Godoy*
- Contributions of seed dispersal and demography to recruitment limitation in a Costa Rican cloud forest, *K G Murray and J M Garcia-C*
- A meta-analysis of gut treatment on seed germination, *A Traveset and M Verdú*
- Seed dispersal effectiveness by Cercopithecus monkeys: implications for seed input into degraded areas, *B A Kaplin and J E Lambert*
- Exploring the link between animal frugivory and plant strategies: the case of primate fruit-processing and post-dispersal seed fate, *J E Lambert*

Conservation, Biodiversity, and Management
- Extinct pigeons and declining bat populations: Are large seeds still being dispersed in the tropical Pacific? *K R McConkey and D R Drake*
- Potential consequences of extinction of frugivorous birds for shrubs of a tropical wet forest, *B A Loiselle and J G Blake*
- Primate frugivory in two species-rich Neotropical Forests: implications for the demography of large-seeded plants in overhunted areas, *C A Peres and M van Roosmalen*
- Patterns of fruit-frugivore interactions in two Atlantic Forest bird communities of southeastern Brazil: implications for conservation, *W R Silva, P De Marco, É Hasui, and V S M Gomes*
- Limitations of animal seed dispersal for enhancing forest succession on degraded lands, *R S Duncan and C A Chapman*
- Frugivory and seed dispersal in degraded tropical east Asian landscapes, *R T Corlett*
- Behavioral and ecological considerations for managing bird damage to cultivated fruits, *M L Avery*
- Harvest and management of forest fruits by humans: implications for fruit-frugivore interactions, *S M Moegenburg*

CABI Publishing, CAB International,
Wallingford, Oxon, OX10 8DE, UK
Tel:+44(0)1491 832111 Fax:+44(0)1491 829292
Email: orders@cabi.org www.cabi-publishing.org/bookshop
CABI Publishing, CAB International,
10 East 40th Street, Suite 3203, New York, NY 10016, USA
Tel:+1 212 481 7018 Fax:+1 212 686 7993
Email: cabi-nao@cabi.org www.cabi-publishing.org/bookshop

For further information or to order please contact CABI *Publishing*, UK or an exclusive CABI *Publishing* distributor in your area. For pre-paid orders in the UK, please add £2.75 for the 1st book and 60p for each additional book ordered (up to max. of 10). For pre-paid orders elsewhere, please add £4.00 for the 1st book and £1.00 for each additional book. For orders not pre-paid, postage and packing will be charged according to the weight of the book.

https://doi.org/10.1079/SSR200190 Published online by Cambridge University Press