Math. Proc. Camb. Phil. Soc. (2008), **144**, 457 © 2008 *Cambridge Philosophical Society*
doi:10.1017/S0305004107000643 *Printed in the United Kingdom*
First published online 11 February 2008

On Bernstein–Heinz–Chern–Flanders inequalities

BY J. L. M. BARBOSA, G. P. BESSA AND J. F. MONTENEGRO
Department of Mathematics, Universidade Federal do Ceará, 60455-760
Fortaleza-CE, Brazil.
e-mail: lucas@mat.ufc.br, bessa@mat.ufc.br, fabio@mat.ufc.br

(*Received* 17 *January* 2006; *revised* 18 *December* 2006)

Abstract

We give an interpretation of the Chern–Heinz inequalities for graphs in order to extend them to transversally oriented codimension one C^2-foliations of Riemannian manifolds. It contains Salavessa's work on mean curvature of graphs and fully generalizes results of Barbosa–Kenmotsu–Oshikiri [**3**] and Barbosa–Gomes–Silveira [**2**] about foliations of 3-dimensional Riemannian manifolds by constant mean curvature surfaces. This point of view of the Chern–Heinz inequalities can be applied to prove a Haymann–Makai–Osserman inequality (lower bounds of the fundamental tones of bounded open subsets $\Omega \subset \mathbb{R}^2$ in terms of its inradius) for embedded tubular neighbourhoods of simple curves of $\mathbb{R}^n$.

1. *Introduction*

Let $G(z)$ be the graph of a C^2-function $z = z(x, y)$ defined on $x^2 + y^2 < r^2$ with mean curvature $H(x, y)$ and Gaussian curvature $K(x, y)$. Heinz [**16**] showed that, if $|H(x, y)| \geqslant b > 0$ then $b \leqslant 1/r$ and if $K(x, y) \geqslant b > 0$ then $b^{1/2} \leqslant 1/r$. As observed by Heinz, these inequalities are improvements of inequalities of the form $r \cdot \varrho(b) \leqslant 1$, for some constant $\varrho(b)$ depending on b, implicit in Bernstein's work [**4**, **5**].

Chern [**10**] and Flanders [**13**], independently, extended these inequalities to graphs of C^2-functions $z = z(x_1, \ldots, x_n)$ defined on bounded domains $\Omega \subset \mathbb{R}^n$ with smooth boundaries $\partial\Omega$, showing that, if the mean curvature satisfies $|H(x_1, \ldots, x_n)| \geqslant b > 0$, then $n \cdot b \leqslant \mathrm{vol}_{n-1}(\partial\Omega)/\mathrm{vol}_n(\Omega)$, and if the scalar curvature $S(x_1, \ldots, x_n) \geqslant b > 0$, then $\sqrt{n(n-1)b} \leqslant \mathrm{vol}_{n-1}(\partial\Omega)/\mathrm{vol}_n(\Omega)$. These inequalities are known as the Chern–Heinz inequalities for graphs.

An immediate corollary of the Chern–Heinz inequalities is that a graph of a C^2-function z: $\mathbb{R}^n \to \mathbb{R}$ has constant mean curvature H if and only if $H = 0$, and it has constant nonnegative scalar curvature $S \geqslant 0$ if and only if $S = 0$.

Isabel Salavessa in [**22**] extended the Chern–Heinz inequality to graphs $G(f)$ of smooth functions f: $M \to \mathbb{R}$. Showing that $n \cdot \inf_\Omega |H_{G(f)}| \leqslant \mathrm{vol}_{n-1}(\partial\Omega)/\mathrm{vol}_n(\Omega)$ for every oriented compact domain $\Omega \subset M$, where $n = \dim M$. In particular, if M has Cheeger constant $h(M) = 0$ then the graph $G(f)$ is minimal. Recall that the Cheeger constant is given by $h(M) = \inf_\Omega(\mathrm{vol}_{n-1}(\partial\Omega)/\mathrm{vol}_n(\Omega))$, where the infimum is taken over all relatively compact subsets Ω of M with smooth boundary, see [**8**].

We can look at a graph of a smooth function z: $\Omega \to \mathbb{R}$ as a leaf of a transversally oriented codimension one smooth foliation of $\Omega \times \mathbb{R}$ obtained by vertical translation of the

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

graph. The Chern–Heinz inequalities say that for this particular foliation of $\Omega \times \mathbb{R}$, the infimum of the mean curvatures of these leaves are bounded above by the $h(\Omega)$. In fact, the infimum of the mean curvatures of the leaves of any transversally oriented codimension one C^2-foliation of $\Omega \times \mathbb{R}$ is bounded above by $h(\Omega)$.

In this paper, we show that the Chern–Heinz inequalities hold for any transversally oriented codimension one C^2-foliations of open subsets Ω of Riemannian manifolds M in the sense that the infimum of the mean curvature of the leaves are bounded above by the fundamental tone $\lambda^*(\Omega)$ of Ω. Here, the upper bound for the infimum of the mean curvatures of the leaves is given in terms of $\lambda^*(\Omega)$, although it is possible to express it in terms of the Cheeger constant $h(\Omega)$. Using this notation we state our main results.

THEOREM 1·1. *Let $\mathcal{F}$ be a transversely oriented codimension one C^2-foliation of a connected open set Ω of an $(n+1)$-dimensional Riemannian manifold M. Then*

$$2\sqrt{\lambda^*(\Omega)} \geqslant n \cdot \inf_{F\in\mathcal{F}} \inf_{x\in F} |H^F(x)|,$$

where H^F stands for the mean curvature function of the leaf F.

THEOREM 1·2. *Let $\mathcal{F}$ be a transversely oriented codimension one C^2 foliation of a complete Riemannian manifold M with nonnegative Ricci curvature. Suppose that the leaves are complete hypersurfaces with the same constant mean curvature H. Then $H = 0$ and each leaf is stable. If furthermore a leaf F is compact then it is totally geodesic and the Ricci curvature of M in the direction normal to F is zero.*

2. *Proof of Theorem* 1·1 *and its consequences*

Let us recall that the fundamental tone $\lambda^*(\Omega)$ of an open subset $\Omega \subset M$ of a smooth Riemannian manifold M is given by

$$\lambda^*(\Omega) = \inf\left\{ \frac{\int_\Omega |\nabla f|^2}{\int_\Omega f^2},\ f \in H_0^1(\Omega) \setminus \{0\} \right\}, \tag{2·1}$$

where $H_0^1(\Omega)$ is the completion of $C_0^\infty(\Omega)$ with respect to the norm $\|\varphi\|_\Omega^2 = \int_\Omega \varphi^2 + \int_\Omega |\nabla\varphi|^2$. We recall that Bessa and Montenegro [**7**] proved that for an open set $\Omega \subset M$ one has

$$\lambda^*(\Omega) \geqslant c(\Omega)^2/4, \tag{2·2}$$

for $c(\Omega) = \sup_{X\in\mathcal{X}(\Omega)} (\inf_\Omega \frac{\operatorname{div} X}{\|X\|_\infty})$, where $\mathcal{X}(\Omega)$ is the set of all smooth vector fields in Ω such that $\|X\|_\infty = \sup_x |X(x)| < \infty$.

To prove Theorem 1·1 we proceed as follows. Let Ω be an open set of a Riemannian manifold M and $\mathcal{F}$ a codimension one transversally oriented C^2-foliation. This just means we may choose a continuous unit vector field η on M that is normal to the leaves of $\mathcal{F}$. Represent by $H^F(x)$ the value of the mean curvature of the leaf F at x computed with respect to η. Set $b = \inf_{F\in\mathcal{F}} \inf_{x\in F} |H^F(x)|$. If $b = 0$ there is nothing to prove. Assume then, that $b > 0$. This implies that H^F does not change sign. Hence, we may choose the unit vector field η in such way that $H^F(x) > 0$ for any $x \in \Omega$. It is well known that $\operatorname{div}\eta = n \cdot H^F$. Therefore, $\inf_\Omega \operatorname{div}\eta = n \cdot \inf_\Omega H_F(x) \geqslant n \cdot b$. Since $\|\eta\|_\infty = 1$ and applying (2·2) to estimate $\lambda^*(\Omega)$ we obtain

$$2\sqrt{\lambda^*(\Omega)} \geqslant c(\Omega) \geqslant \inf_\Omega \frac{\operatorname{div}\eta}{\|\eta\|_\infty} = n \cdot \inf_{x\in\Omega} H_F(x) \geqslant n \cdot b.$$

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

This proves Theorem 1·1.

This theorem has a number of interesting consequences, stated below as corollaries. It imposes strong restrictions for the existence of foliations by constant mean curvature hypersurfaces on open sets with zero fundamental tone or with Ricci curvature bounded below, see Corollaries 2·1 and 2·2.

COROLLARY 2·1. *Let $\mathcal{F}$ be a transversely oriented codimension one C^2-foliation of a Riemannian manifold M for which $\lambda^*(M) = 0$. If the leaves of $\mathcal{F}$ have the same constant mean curvature then they are minimal submanifolds of M.*

Remark. The class of Riemannian manifolds M with $\lambda^*(M) = 0$ is quite large. Besides all closed Riemannian manifolds, it contains all open Riemannian manifolds with asymptotically nonnegative Ricci curvature, see [**9**]. Recall that an open Riemannian manifold M has asymptotically nonnegative Ricci curvature if $Ric_M(x) \geqslant -\psi(dist_M(x_0, x))$, for a continuous function $\psi\colon [0, \infty) \to [0, \infty)$ with $\lim_{t\to\infty} \psi(t) = 0$, $x_0 \in M$.

In [**3**], Barbosa–Kenmotsu–Oshikiri considered a transversely oriented codimension-one C^2-foliation $\mathcal{F}$ of an open subset Ω of an n-dimensional complete Riemannian manifold M with Ricci curvature satisfying $Ric_M \geqslant (n-1)\kappa$. They assumed that the leaves were complete oriented hypersurfaces with the same constant mean curvature H. In their theorem 3·1 they showed that, if M is compact and $\kappa = 0$ then the leaves of $\mathcal{F}$ are totally geodesic and the Ricci curvature of M is zero in the directions normal to the leaves. The remark above, Corollary 2·1 coupled with Theorem 1·2 implies as a particular case, the Barbosa–Kenmotsu–Oshikiri main result, theorem 3·1.

In [**3**, theorem 38], they investigated the case when M is the simply connected space form of constant sectional curvature κ. When $\kappa \leqslant 0$ and $H \geqslant (n-1)\sqrt{-\kappa}$ they concluded that $H = (n-1)\sqrt{-\kappa}$. The following corollary of Theorem 1·1 extends theorem 3·8 of Barbosa–Kenmotsu–Oshikiri to n-dimensional Riemannian manifolds with Ricci curvature satisfying $\mathrm{Ric}_M \geqslant (n-1)\kappa$.

COROLLARY 2·2. *Let $\mathcal{F}$ be transversely oriented codimension-one C^2-foliation of a complete n-dimensional Riemannian manifold M with Ricci curvature* $\mathrm{Ric}_M \geqslant (n-1)\kappa$. *Then*:

(i) $2\sqrt{\lambda^*(\mathbb{M}^n(\kappa))} \geqslant \inf_{F\in\mathcal{F}} \inf_{x\in F} |H_F(x)|$, *where $\mathbb{M}^n(\kappa)$ is the simply connected space form of constant sectional curvature κ and dimension n*;

(ii) *If $|H_F| \geqslant b > 0$ then $\kappa = -a^2$ for some positive a satisfying $(n-1)\cdot a \geqslant b$.*

Let $B_M(r)$ be a geodesic ball of radius r in the Riemannian Manifold M whose Ricci curvature satisfies $\mathrm{Ric}_M \geqslant (n-1)\kappa$, and let $B_{\mathbb{M}(\kappa)}(r)$ be the geodesic ball of radius r in the simply connected n-dimensional space $\mathbb{M}(\kappa)$ of constant sectional curvature κ. By Cheng's Comparison Theorem (see [**9**]) we know that

$$\lambda^*(B_M(r)) \leqslant \lambda^*\big(B_{\mathbb{M}(\kappa)}(r)\big)\,.$$

Since the fundamental tone of M can be also given by $\lambda^*(M) = \lim_{r\to\infty} \lambda^*(B_M(r))$, it follows that $\lambda^*(M) \leqslant \lambda^*(\mathbb{M}(\kappa))$. This and Theorem 1·1 gives us:

$$\inf_{F\in\mathcal{F}} \inf_{x\in F} |H_F(x)| \leqslant 2\sqrt{\lambda^*(M)} \leqslant 2\sqrt{\lambda^*(\mathbb{M}(\kappa))}$$

and (i) is proved.

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

To obtain (ii) observe that, if $|H_F| \geqslant b > 0$ then, from (i) we have $\sqrt{\lambda^*(\mathbb{M}(\kappa))} \geqslant b/2$. But, $\lambda^*(\mathbb{M}(\kappa)) \neq 0$ if and only if $\kappa = -a^2$ in which case $\lambda^*(\mathbb{M}(\kappa)) = (n-1)^2 a^2/4$. Thus the inequality $\sqrt{\lambda^*(\mathbb{M}(\kappa))} \geqslant b/2$ can occur only when $\kappa = -a^2$ and $(n-1)a \geqslant b$. This completes the proof of the Corollary.

Theorem 1·1 also has a version for scalar curvature provided the ambient manifold has nonpositive sectional curvature.

COROLLARY 2·3. *Let M be an $(n+1)$-dimensional Riemannian manifold with nonpositive sectional curvature and let $\mathcal{F}$ be a transversely oriented codimension one C^2-foliation of a connected open set $\Omega \subset M$. Suppose that the scalar curvature S_F of each leaf if nonnegative. Then*

$$\sqrt{\inf S} \leqslant 2\sqrt{\lambda^*(\Omega)}.$$

In particular, if $\lambda^*(M) = 0$, $\Omega = M$, and all the leaves have the same has constant nonnegative scalar curvature $S > 0$, then $S = 0$.

If $\inf S = 0$ there is nothing to prove. So, we assume that $\inf S = c > 0$. Let $p \in F$ and $\{e_1, \ldots e_n\}$ be an orthonormal basis for the tangent space T_pF of the leaf $F \in \mathcal{F}$. The Gauss equation for the plane generated by e_i, e_j is:

$$\tilde{K}(e_i, e_j) = \overline{K}(e_i, e_j) + \langle B(e_i, e_i),\ B(e_j, e_j)\rangle - |B(e_i, e_j)|^2,$$

where $\tilde{K}$ represents the Gaussian curvature of F and $\overline{K}$ is the one of the ambient space $\Omega \times \mathbb{R}$. Adding these equations results:

$$S(p) = \sum_{i,j} \overline{K}(e_i, e_j) + n^2 H^2 - \|B\|^2.$$

Since the sectional curvatures K of M are, by hypothesis, nonpositive, then $\overline{K} \leqslant 0$. It follows that

$$S(p) \leqslant n^2 H^2$$

and, since $S \geqslant c > 0$ then $H \geqslant \sqrt{c}/n > 0$. By Theorem 1·1 we have that

$$2\sqrt{\lambda^*(\Omega)} \geqslant n \inf_{F\in\mathcal{F}} \inf_{x\in F} |H_f(x)| \geqslant \sqrt{c} = \sqrt{\inf S}.$$

This proves Corollary 2·3.

We turn to higher order mean curvatures. Recall that if $\psi\colon N \to M$ is an n-dimensional oriented hypersurface of M and $k_1, \ldots, k_n$ are the principal curvatures at $p \in N$, then the r-th mean curvatures H_r of $\psi(N)$ at $\psi(p)$ are defined by the identity

$$(1+tk_1)(1+tk_2)\cdots(1+tk_n) = 1 + \binom{n}{1} H_1 t + \binom{n}{2} H_2 t^2 + \cdots + \binom{n}{n} H_n t^n$$

for all real numbers t. Thus, H_1 is the mean curvature of ψ, H_n is the Gauss–Kronecker curvature. Since we always have $H_1^2 \geqslant H_2$, the following version of Theorem 1·1 for the 2nd mean curvature H_2, is direct.

COROLLARY 2·4. *Let $\mathcal{F}$ be a transversely oriented codimension one C^2-foliation of a connected open set Ω of a Riemannian manifold M. Suppose that the leaves have the second mean curvature $H_2 \geqslant 0$. Then*

$$2\sqrt{\lambda^*(\Omega)} \geqslant n \cdot \inf_{F\in\mathcal{F}} \inf_{x\in F} \left(H_2^F\right)^{1/2}(x)$$

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

where H_2^F stands for the second mean curvature function of the leaf F. In particular, if $\lambda^*(M) = 0$, $\Omega = M$ and all the leaves have the same constant 2nd mean curvature $H_2 \geqslant 0$, then $H_2 = 0$.

If the principal curvatures at a point $q \in \psi(N)$ are positive, $k_i(q) > 0$ and if $H_r \geqslant 0$ then Gårding inequalities, $H_k^{1/k} \leqslant H_1$, $H_k^{(k-1)/k} \leqslant H_{k-1}$, $k = 1, \dots r$ hold everywhere, see [**16**, **18**]. It was observed in the proof of [**13**, theorem 1·3] that Gårding inequalities are still true under the weaker hypothesis $k_i(q) \geqslant 0$, if $H_r > 0$. The statement of the next corollary could be about foliations but we preferred to make it about graphs. It generalizes [**13**, corollary 1·7].

COROLLARY 2·5. *Let M be a complete Riemannian manifold with $\lambda^*(M) = 0$ and $z\colon M \to \mathbb{R}$ be a C^2-function whose graph $G(z) \subset M \times \mathbb{R}$ has constant nonnegative rth mean curvature $H_r \geqslant 0$. If $r \geqslant 3$ suppose that there is a point $q \in G(z)$ with all principal curvature $k_i(q) \geqslant 0$. Then $H_r = 0$.*

The cases $r = 1$ and $r = 2$ are covered by Theorem 1·1 and Corollary 2·1. Now, suppose that $r \geqslant 3$ and the graph has a point q with principal curvatures $k_i(q) \geqslant 0$. If $H_r \geqslant 0$ is constant then we may suppose that $H_r > 0$, otherwise there is nothing to prove. Then we have that $H_1 \geqslant H_r^{1/r} > 0$. Since $\lambda^*(M) = 0$, by Theorem 1·1, $0 = (2/n)\sqrt{\lambda^*(M)} \geqslant \inf_M H_1 \geqslant H_r^{1/r} > 0$. A contradiction.

3. *Proof of Theorem* 1·2

Let M be a complete Riemannian manifold with Ricci curvature satisfying $\mathrm{Ric}_M \geqslant 0$. Then we have

$$\lambda^*(M) = 0. \tag{3·1}$$

Let $\mathcal{F}$ be transversely oriented codimension-one C^2-foliation of M. Then, there exists a unit vector field η in M normal to all the leaves of $\mathcal{F}$. Let H_F represent the mean curvature of the leaf F computed with respect to η. Assume the leaves of $\mathcal{F}$ are complete hypersurfaces with the same constant mean curvature H, that is, $H_F = H$, where H is a constant. It then follows from (3·1) and Theorem 1·1 that each leaf $F \in \mathcal{F}$ is minimal.

Fix a leaf F. On F consider the Jacobi operator:

$$L = \triangle + \mathrm{Ric}(\eta) + |A|^2 \tag{3·2}$$

where $\triangle$ represents the Laplace operator in F, A represents the second fundamental form of F computed with respect to η, and Ric (η) is the Ricci curvature of M in the direction of η.

Let C be a compact set of F with smooth boundary ∂C. Take another compact set C' such that $C \subset C'$. We may assume that the only solution of the problem

$$\begin{aligned} L(v) &= 0 \qquad \text{in } C', \\ v &= 0 \qquad \text{in } \partial C' \end{aligned}$$

is the function $v \equiv 0$. This means that zero is not an eigenvalue of the operator L in C'. Thus there exists a positive function $u \in C^\infty(C')$ with

$$\begin{aligned} L(u) &= 1 \qquad \text{in } C', \\ u &= 0 \qquad \text{in } \partial C'. \end{aligned}$$

Consider now the variation $\psi_t\colon C' \to M$ of C' given by $\psi_t(x) = \exp_x(t\,u(x)\eta(x))$ for

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

$|t| < \epsilon$. Represent by $H(t)$ the mean curvature of the immersion ψ_t. It has derivative at $t = 0$ given by $2H'(0) = L(u) = 1$ on C' (see [**21**]). Since $H(0) = 0$, then, for small $0 < t < \epsilon$, $H(t) > 0$. Now we analyze some possibilities for the behavior of u.

If u is positive at some interior point of C' then $\psi_t(C')$ must have a tangency point with some leaf of $\mathcal{F}$, and this is not possible by the maximum principle. Thus $u \leqslant 0$.

If $u \leqslant 0$ and it is zero at an interior point of C' we have that $u \equiv 0$ and this is impossible since $Lu = 1$. Thus $u < 0$ in the interior of C' and $u = 0$ on $\partial C'$. Setting $w = -u$ we have that $w > 0$ in the interior of C' with $L(w) < 0$.

Let u_1 be the first eigenfunction of C, i.e.,

$$\begin{aligned} L(u_1) + \lambda_1(C)u_1 &= 0 \qquad \text{on } C, \\ u_1 &= 0 \qquad \text{on } \partial C. \end{aligned}$$

Furthermore we know that u_1 is not zero in the interior of C. Hence, we may assume it is positive in the interior of C.

Suppose by contradiction that $\lambda_1(C) < 0$. Define in C the function $h = w - t\,u_1$ This is possible since $C \subset C'$. Observe that on ∂C we have $h = w > 0$ and independent of t. Hence, we may choose t such that $h \geqslant 0$ and is zero at some point in the interior of C. Observe that

$$L(h) = L(w) - tL(u_1) \leqslant t\lambda_1(C)u_1 < 0$$

on C'. Then $\triangle h < 0$ and h has a minimum in the interior of C. By the maximum principle h is constant. Since $h = 0$ at one point, then it is everywhere zero. But this is a contradiction since, as we have seen, $h > 0$ at ∂C.

This contradiction shows that $\lambda_1(C) \geqslant 0$. Suppose that F is not stable. Then there exists a function $f\colon F \to \mathbb{R}$ such that $\int_F f = 0$ and $L(f) + \bar{\lambda}_1 f = 0$ with $\bar{\lambda}_1 < 0$. Here $\bar{\lambda}_1$ is the first eigenvalue of the twisted problem:

$$\begin{aligned} L(f) + \bar{\lambda} f &= 0 \qquad \text{on } F, \\ \int_F f &= 0. \end{aligned}$$

But then $\bar{\lambda}_1 < 0$ would be the first eigenvalue λ_1 of L on the set $\{x \in F;\ f(x) > 0\}$. But, exhausting this set with a family of compact sets we may show that $\lambda_1 \geqslant 0$. This contradiction shows that F is stable as we wished to prove.

Assume now that F is compact. Since F is stable, by a result of Fisher–Colbrie [**12**] there is a positive function $g\colon F \to \mathbb{R}$ solution to

$$\triangle g + (\mathrm{Ric}(\eta) + |A|^2)g = 0.$$

Integrating this equation over F we obtain

$$0 = \int_F \triangle g + (\mathrm{Ric}(\eta) + |A|^2)g = \int_F (\mathrm{Ric}(\eta) + |A|^2)g.$$

Since $\mathrm{Ric}(\eta) \geqslant 0$ and $g > 0$ then we conclude that

$$\mathrm{Ric}(\eta) = 0$$

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

and

$$|A|^2 = 0.$$

Consequently F is totally geodesic and the Ricci curvature of M in direction normal to F is zero. This concludes the proof of Theorem 1·2.

Remark. For $n = 3$, Schoen [**23**] showed that a complete stable minimal surface in a 3-dimensional Riemannian manifold with nonnegative Ricci curvature is totally geodesic. Thus, for $n = 3$, Theorem 1·2 fully generalizes [**3**, theorem 3·1]. *A similar stability result as Theorem* 1·2 was proved by Bessa, Jorge and Oliveira for minimal immersions with bounded curvature in [**6**].

4. *Haymann–Makai–Osserman inequality*

Recall that the inradius $\rho(\Omega)$ of a connected open set Ω of a Riemannian manifold M is defined as $\rho(\Omega) = \sup\{r > 0;\ B_M(r) \subset \Omega\}$, where $B_M(r)$ represents a geodesic ball of radius r of M. In [**18**], Makai proved that the fundamental tone $\lambda^*(\Omega)$ of a simply connected bounded domain $\Omega \subset \mathbb{R}^2$ with inradius ρ and smooth boundary is bounded below by $\lambda_1(\Omega) \geqslant 1/4\rho^2$. Unaware of Makai's result, Haymann [**15**] proved years later that $\lambda_1(\Omega) \geqslant 1/900\rho^2$. Osserman [**20**] among other things improved Haymann's estimate back to $\lambda_1(\Omega) \geqslant 1/4\rho^2$. Recently, the Haymann–Makai–Osserman inequality was improved by Bañelos–Carroll in [**1**] to $0.6197/\rho^2$.

Our next result extends of Haymann–Makai–Osserman inequality to embedded tubular neighbourhoods of simple smooth curves in $\mathbb{R}^n$ with variable radius.

THEOREM 4·1. *Let $\gamma\colon I = (\alpha, \beta) \subset \mathbb{R} \to \mathbb{R}^n$ be a simple smooth curve and $T_\gamma(\rho(t))$ be an embedded tubular neighbourhood of γ with variable radius $\rho(t)$ and smooth boundary $\partial T_\gamma(\rho(t))$. Let $\rho_o = \sup_t \rho(t) > 0$ be its* inradius. *Then*

$$\lambda^*(T_\gamma(\rho(t))) \geqslant \frac{(n-1)^2}{4\rho_o^2}. \tag{4·1}$$

Consider the family of balls of $\mathbb{R}^n$ $B_t = B_{\mathbb{R}^n}(\gamma(t), \rho_o)$, $t \in I$ with center at $\gamma(t)$ and radius ρ_o. The set $\{S_t = \partial B_t \cap T_\gamma(\rho(t))\}$ is a smooth codimension one transversally oriented foliation of $T_\gamma(\rho(t)) \setminus (B_\alpha \cup B_\beta)$. Pushing the family B_t little further one can fill $T_\gamma(\rho(t))$ with a smooth codimension one transversally oriented foliation such that the mean curvature of the leaves is constant $1/\rho_o$. The Theorem 1·1 yields that $\lambda^*(T_\gamma(\rho)) \geqslant (n-1)^2/4\rho^2$.

Acknowledgements. The research and writing of this work was partially supported by CNPq-Brazil.

REFERENCES

[**1**] R. BAÑUELOS and T. CARROLL. Brownnian motion and the fundamental frequence of a drum. *Duke Math. J.* **75** (1994), 575–602.
[**2**] J. L. M. BARBOSA, J. M. GOMES and A. M. SILVEIRA. Foliations of 3-dimensional space forms by surfaces with constant mean curvature *Bol. Soc. Bras. Mat.* **18** (1987), 1–12.
[**3**] J. L. M. BARBOSA, K. KENMOTSU and O. OSHIKIRI. Foliations by hypersurfaces with constant mean curvature. *Math. Z.* **207** (1991), 97–108.
[**4**] S. BERNSTEIN. Sur la généralisation du problème de Dirichlet. *Math. Ann.* **69** (1910), 82–136.
[**5**] S. BERNSTEIN. Sur les surfaces définies au moyen de leur courbure moyenne ou totale. *Ann. École Norm. Sup.* **27** (1909), 233–256.

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press

[**6**] G. P. BESSA, L. P. JORGE and G. OLIVEIRA–FILHO. Half-space theorems for minimal surfaces. *J. Diff. Geom.* **57** (2001), 493–508.
[**7**] G. P. BESSA and J. F. MONTENEGRO. Eigenvalue estimates for submanifolds with locally bounded mean curvature. *Ann. Global Anal. Geom.* **24** (2003), 279–290.
[**8**] J. CHEEGER. A. lower bound for the smallest eigenvalue of the Laplacian. *Problems in Analysis* (Princenton Univ. Press, 1970), 195–199.
[**9**] S. Y. CHENG. Eigenvalue comparison theorems and its geometric applications. *Math. Z.* **143** (1975), 289–297.
[**10**] S. S. CHERN. On the curvature of a piece of hypersurface in Euclidean space. *Abh. Math. Sem Hamburg* **29** (1965), 77–91.
[**11**] M. F. ELBERT. Constant positive 2-mean curvature hypersurfaces. *Illinois J. Math.* **46** (2002), 247–267.
[**12**] D. FISCHER–COLBRIE and R. SCHOEN. The structure of complete stable minimal surfaces in 3-manifolds of nonnegative scalar curvature. *Comm. Pure. Appl. Math.* **33** (1980), 199–211.
[**13**] H. FLANDERS. Remark on mean curvature. *J. London Math. Soc.* (2) **41** (1966), 364-366.
[**14**] F. FONTENELE and S. L. SILVA. Sharp estimates for size of balls in the complement of a hypersurface. *Geom. Dedicata* **115** (2005), 163–179.
[**15**] W. HAYMANN. Some bounds for principal frequency. *Applicable Anal.* **7** (1978), 247–254.
[**16**] E. HEINZ. Über Flächen mit eindeutiger projektion auf eine ebene, deren krümmungen durc ungleichungen eingschränkt sind. *Math. Ann.* **129** (1955), 451–454.
[**17**] L. GÅRDING. An inequality for hyperbolic polinomials. *J. Math. Mech.* **8** (1959), 957–965.
[**18**] E. MAKAI. A lower estimation of the principal frequencies of simply connected membranes. *Acta Math. Acad. Sci. Hungar.* **16** (1965), 319–366.
[**19**] S. MONTIEL and A. ROS. Compact hypersurfaces: the Alexandrov theorem for higher order mean curvatures. *Differential Geometry, a symposium in honor of Manfredo do Carmo.* B. Lawson and K. Teneblat, (Eds.). Pitman Monographs, **52** (Longman Scientific and Technical, 1991), 179–296.
[**20**] R. OSSERMAN. A note on Hymann's theorem on the bass note of a drum. *Comment. Math. Helv.* **52** (1977), 545–555.
[**21**] A. ROS. Compactness of spaces of properly embedded minimal surfaces with finite total curvature. *Indiana Univ. Math. J.* **44** (1995), 139–152.
[**22**] ISABEL M. C. SALAVESSA. Graphs with parallel mean curvature. *Proc. Amer. Math. Soc.* **107** (1989), 449–458.
[**23**] R. SCHOEN. Estimates for stable minimal surfaces in three dimensional manifolds. *Ann. of Math. Stud.* **103** (1983), 111–126.

https://doi.org/10.1017/S0305004107000643 Published online by Cambridge University Press