A.A.S. **2**, II, 217-232 (2007)

SOME FINITE TIME RUIN PROBLEMS

By D. C. M. Dickson

ABSTRACT

In the classical risk model, we use probabilistic arguments to write down expressions in terms of the density function of aggregate claims for joint density functions involving the time to ruin, the deficit at ruin and the surplus prior to ruin. We give some applications of these formulae in the cases when the individual claim amount distribution is exponential and Erlang(2).

KEYWORDS

Finite Time Ruin; Joint Density; Deficit at Ruin

CONTACT ADDRESS

David C. M. Dickson, Centre for Actuarial Studies, Department of Economics, University of Melbourne, Victoria 3010, Australia. E-mail: dcmd@unimelb.edu.au

1. Introduction

We consider the classical risk model in which the surplus process $\{U(t)\}_{t\geq 0}$ is given by:

$$U(t) = u + ct - S(t)$$

where u is the insurer's initial surplus, c is the rate of premium income per unit time, and $S(t)$ represents the aggregate amount of claims up to time t. The process $\{S(t)\}_{t\geq 0}$ is a compound Poisson process, with $S(t) = \sum_{i=1}^{N(t)} X_i$, where $\{X_i\}_{i=1}^{\infty}$ is a sequence of independent and identically distributed random variables, with X_i representing the amount of the ith claim, and $N(t)$ follows a Poisson distribution with parameter λt. We assume that the individual claim amounts have a continuous distribution with distribution function $F = 1 - \overline{F}$, density function f and mean m_1, and we adopt the notation $G(x, t) = \Pr(S(t) \leq x)$, with $g(x, t) = \frac{\partial}{\partial x} G(x, t)$.

The finite time ruin probability is defined as:

$$\psi(u, t) = \Pr(U(s) < 0 \text{ for some } s,\ 0 < s \leq t)$$

and let $\delta(u, t) = 1 - \psi(u, t)$. Prahbu's (1961) formula for $\delta(u, t)$ is:

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$\delta(u, t) = G(u + ct, t) - c \int_0^t g(u + cs, s)\delta(0, t - s)ds \tag{1}$$

with:

$$\delta(0, t) = e^{-\lambda t} + \int_0^{ct} \delta(0, t, x)dx$$

where:

$$\delta(0, t, x) = \frac{x}{ct} g(ct - x, t)$$

is the density associated with non-ruin from initial surplus zero over $[0, t]$ and a surplus of x at time t. (See Gerber, 1979, p112).

Although extensive numerical work has been done using Prahbu's formula, much of which is summarised in Seal (1978), few explicit solutions for $\delta(u, t)$ have arisen from this formula, presumably because the use of Prahbu's formula requires explicit expressions for G and g. The notable exception is when F is the exponential distribution function.

Dickson & Willmot (2005) derived an expression for the density of the time to ruin, T, defined as:

$$T = \inf\{t: U(t) < 0\}$$

with $T = \infty$ if $U(t) > 0$ for all $t > 0$. Their expression was found by inverting the Laplace transform of T. It is not a concise solution, but is suitable for computational purposes, as illustrated by Dickson & Willmot (2005) and Willmot & Woo (2007). Our main purpose in this paper is to identify alternative expressions for the density of the time to ruin. We do this in Section 2 by applying a probabilistic argument, and we can confirm the validity of the argument by differentiating equation (1). Based on the probabilistic argument, we then give formulae in Section 3 for some joint distributions relating to the time to ruin, the deficit at ruin and the surplus prior to ruin. Some illustrations of the application of these formulae are given in Sections 4 and 5, and conclusions are presented in Section 6.

2. The Density of the Time to Ruin

Let $w(u, t)$ denote the (defective) density of the time to ruin, so that $w(u, t) = \frac{\partial}{\partial t}\psi(u, t)$. We know from Dickson & Willmot (2005) that:

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$w(0,t) = \lambda e^{-\lambda t}\overline{F}(ct) + \lambda \int_0^{ct} \frac{x}{ct} g(ct-x,t)\overline{F}(x)dx \tag{2}$$

$$= \lambda e^{-\lambda t}\overline{F}(ct) + \lambda \int_0^{ct} \delta(0,t,x)\overline{F}(x)dx \tag{3}$$

and we can interpret this as follows. The expression $w(0,t)dt$ is interpreted as the probability of ruin occurring in the interval $(t, t+dt)$. The term $\lambda e^{-\lambda t}\overline{F}(ct)dt$ is interpreted as the probability that the first claim occurs in the interval $(t, t+dt)$, and that this claim exceeds ct, causing ruin. Similarly, for a fixed value of x, $\lambda\delta(0,t,x)\overline{F}(x)dx\,dt$ is interpreted as the probability of:

(a) non-ruin over $[0,t]$ with $U(t) = x$; and
(b) a claim occurring in the interval $(t, t+dt)$ of amount greater than x, causing ruin.

Extending this interpretation to an initial surplus greater than zero, we get:

$$w(u,t) = \lambda e^{-\lambda t}\overline{F}(u+ct) + \lambda \int_0^{u+ct} \delta(u,t,x)\overline{F}(x)dx \tag{4}$$

where $\delta(u,t,x)$ is the density associated with non-ruin from an initial surplus u over $[0,t]$ and a surplus of x at time t. Extending formula (3.13) of Dickson & Waters (2006) from $t=1$ to a general value for t, we obtain:

$$\begin{aligned}\delta(u,t,x) = {} & g(u+ct-x,t) - cI(t > x/c)\int_0^{t-x/c} g(u+cs,s)\delta(0,t-s,x)ds \\ & - I(t > x/c)g(u+ct-x, t-x/c)e^{-\lambda x/c}\end{aligned} \tag{5}$$

where I is the indicator function. Formula (4) is not the most useful formula for computing $w(u,t)$, but its structure allows us to write down a joint density function, and this will be done in the next section.

A more convenient formula for computing $w(u,t)$ is:

$$\begin{aligned}w(u,t) = {} & \lambda e^{-\lambda t}\overline{F}(u+ct) + \lambda \int_0^{u+ct} g(u+ct-x,t)\overline{F}(x)dx \\ & - c\int_0^t g(u+cs,s)w(0,t-s)ds\end{aligned} \tag{6}$$

which follows by inserting formula (5) for $\delta(u,t,x)$ in formula (4). We can use this equation to compute $w(u,t)$ for certain individual claim amount distributions, as illustrated in Sections 4 and 5.

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

3. Joint Densities

Now let $Y \equiv |U(T)|$ denote the deficit at ruin and let $w_{Y,T}(u, y, t)$ denote the (defective) joint density of Y and T. Dickson & Willmot (2005, Section 3) obtain formula (3) by inverting the Laplace transform of T. If we apply the same approach to the bivariate Laplace transform of Y and T, straightforward manipulations give:

$$w_{Y,T}(0, y, t) = \lambda e^{-\lambda t} f(ct + y) + \lambda \int_0^{ct} \delta(0, t, x) f(x + y) dx. \tag{7}$$

The interpretation here is as for formula (3), except that, for $x \leq ct$, $\overline{F}(x)$ in formula (3) is replaced by $f(x + y)$, i.e. rather than requiring a claim above x to cause ruin, we require a claim of amount $x + y$ to cause ruin with a deficit of y. By adapting this interpretation to an initial surplus $u > 0$, we obtain:

$$w_{Y,T}(u, y, t) = \lambda e^{-\lambda t} f(u + ct + y) + \lambda \int_0^{u+ct} \delta(u, t, x) f(x + y) dx. \tag{8}$$

Inserting for $\delta(u, t, x)$, similar manipulations to those in the previous section yield:

$$\begin{aligned} w_{Y,T}(u, y, t) = \lambda e^{-\lambda t} f(u + ct + y) + \lambda \int_0^{u+ct} g(u + ct - x, t) f(x + y) dx \\ - c \int_0^t g(u + c(t - s), t - s) w_{Y,T}(0, y, s) ds. \end{aligned} \tag{9}$$

Formula (9) provides an alternative means to the transform inversion method of calculating the joint density of Y and T presented in Dickson (2008). Its main advantage is that it is a simple integral expression in terms of g, which makes it suitable for direct application for certain forms of f, or for numerical integration. Its main disadvantage is that the third term is of a similar form to the integral term in Prahbu's formula.

Further, if we let $X \equiv U(T^-)$ denote the surplus prior to ruin and define $w_{X,Y,T}(u, x, y, t)$ to be the (defective) joint density of X, Y and T, we see that, for $0 < x < u + ct$:

$$w_{X,Y,T}(u, x, y, t) = \lambda \delta(u, t, x) f(x + y)$$

with density of $\lambda e^{-\lambda t} f(u + ct + y)$ for $x = u + ct$. This argument is given by Wu *et al.* (2003), but they use a different formula for $\delta(u, t, x)$.

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

Integrating out y, we obtain the joint density of the surplus prior to ruin and the time to ruin as:

$$w_{X,T}(u, x, t) = \lambda\delta(u, t, x)\overline{F}(x)$$

for $0 < x < u + ct$, with density of $\lambda e^{-\lambda t}\overline{F}(u + ct)$ for $x = u + ct$. This leads to the relationship stated in Gerber & Shiu (1997):

$$w_{X,Y,T}(u, x, y, t) = w_{X,T}(u, x, t)\frac{f(x + y)}{\overline{F}(x)}.$$

4. Exponential Claims

Let us first consider the case when $\overline{F}(x) = e^{-\alpha x}$, where $\alpha > 0$, so that $f(x) = \alpha\overline{F}(x)$. Then, inserting for f in equation (8), this gives:

$$\begin{aligned} w_{Y,T}(u, y, t) &= \lambda e^{-\lambda t}\alpha e^{-\alpha(u+ct+y)} + \lambda\int_0^{u+ct}\delta(u, t, x)\alpha e^{-\alpha(x+y)}dx \\ &= \alpha e^{-\alpha y}\left(\lambda e^{-\lambda t}\overline{F}(u + ct) + \lambda\int_0^{u+ct}\delta(u, t, x)\overline{F}(x)dx\right) \\ &= \alpha e^{-\alpha y}w(u, t) \end{aligned}$$

which is a well-known result — see, for example, Gerber (1979).

Consider equation (6) for $w(u, t)$. The first two terms give:

$$\begin{aligned} &\lambda e^{-\lambda t}\overline{F}(u + ct) + \lambda\int_0^{u+ct} g(u + ct - x, t)\overline{F}(x)dx \\ &= \lambda e^{-\lambda t-\alpha u-\alpha ct} + \lambda\sum_{m=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^m}{m!}\int_0^{u+ct}\frac{\alpha^m x^{m-1}e^{-\alpha x}}{\Gamma(m)}e^{-\alpha(u+ct-x)}dx \\ &= \lambda e^{-\lambda t-\alpha u-\alpha ct}\left(1 + \sum_{m=1}^{\infty}\frac{(\alpha\lambda t)^m(u + ct)^m}{m!m!}\right) \\ &= \lambda e^{-\lambda t-\alpha u-\alpha ct}I_0\left(\sqrt{4\alpha\lambda t(u + ct)}\right) \end{aligned}$$

where:

$$I_v(t) = \sum_{n=0}^{\infty}\frac{(t/2)^{2n+v}}{n!(n + v)!}$$

is the modified Bessel function of order v.

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

To obtain the third term of equation (6), we first find $w(0,t)$ from equation (2), as:

$$\begin{aligned} w(0,t) &= \lambda e^{-\lambda t}\overline{F}(ct) + \lambda\int_0^{ct}\frac{x}{ct}g(ct-x,t)\overline{F}(x)dx \\ &= \lambda e^{-\lambda t}e^{-\alpha ct} + \lambda\sum_{m=1}^{\infty}e^{-\lambda t}\frac{(\lambda t)^m}{m!}\int_0^{ct}\frac{x}{ct}\frac{\alpha^m(ct-x)^{m-1}e^{-\alpha(ct-x)}}{\Gamma(m)}e^{-\alpha x}dx \\ &= \lambda e^{-\lambda t}e^{-\alpha ct} + \lambda\sum_{m=1}^{\infty}e^{-\lambda t}e^{-\alpha ct}\frac{(\alpha\lambda t)^m}{m!\Gamma(m)}\frac{\Gamma(2)\Gamma(m)}{\Gamma(m+2)}(ct)^m \\ &= \lambda e^{-\lambda t}e^{-\alpha ct}\sum_{m=0}^{\infty}\frac{\left(\alpha\lambda ct^2\right)^m}{m!(m+1)!} \\ &= \lambda e^{-\lambda t}e^{-\alpha ct}\eta_1(t) \end{aligned}$$

where:

$$\eta_1(t) = \sum_{m=0}^{\infty}\frac{(\alpha\lambda ct^2)^m}{m!(m+1)!} = \frac{1}{\sqrt{\alpha\lambda ct^2}}I_1\left(\sqrt{4\alpha\lambda ct^2}\right).$$

Thus, (minus) the third term in equation (6) is:

$$\begin{aligned} & c\int_0^t g(u+c(t-s),t-s)\lambda e^{-(\lambda+\alpha c)s}\eta_1(s)ds \\ &= \lambda c\int_0^t\sum_{n=1}^{\infty}e^{-\lambda(t-s)}\frac{(\lambda(t-s))^n}{n!}\frac{\alpha^n(u+c(t-s))^{n-1}}{(n-1)!}e^{-\alpha(u+c(t-s))}e^{-(\lambda+\alpha c)s}\eta_1(s)ds \\ &= \lambda c e^{-\alpha u-(\lambda+\alpha c)t}\int_0^t\sum_{n=1}^{\infty}\frac{(\lambda(t-s))^n}{n!}\frac{\alpha^n(u+c(t-s))^{n-1}}{(n-1)!}\eta_1(s)ds. \end{aligned} \tag{10}$$

We can evaluate (10) by observing that the integral is the convolution of:

$$\eta_2(t) = \sum_{n=1}^{\infty}\frac{(\alpha\lambda t)^n}{n!}\frac{(u+ct)^{n-1}}{(n-1)!} = \frac{\alpha\lambda t}{\sqrt{\alpha\lambda t(u+ct)}}I_1\left(\sqrt{4\alpha\lambda t(u+ct)}\right)$$

and η_1, and we can find this from the following result, taken from Erdélyi (1954, p201).

Result 4.1. Let:

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$\phi(t) = \frac{t^{v/2}}{(t+\beta)^{v/2}} I_v\left(A\sqrt{t^2+\beta t}\right).$$

Then:

$$\tilde{\phi}(s) = \int_0^\infty e^{-st}\phi(t)dt = \frac{A^v}{\sqrt{s^2-A^2}} \frac{1}{\left(s+\sqrt{s^2-A^2}\right)^v} \exp\left\{\frac{\beta}{2}\left(s-\sqrt{s^2-A^2}\right)\right\}. \quad \square$$

To apply this result we note that we can write:

$$\eta_2(t) = \frac{\sqrt{\alpha\lambda/c}\sqrt{t}}{\sqrt{\frac{u}{c}+t}} I_1\left(\sqrt{4\alpha\lambda c}\sqrt{\frac{u}{c}t+t^2}\right).$$

Then, by Result 4.1, with $v = 1$, $\beta = u/c$ and $A = \sqrt{4\alpha\lambda c}$:

$$\tilde{\eta}_2(s) = \sqrt{\alpha\lambda/c}\frac{A}{\sqrt{s^2-A^2}} \frac{1}{s+\sqrt{s^2-A^2}} \exp\left\{\frac{u}{2c}\left(s-\sqrt{s^2-A^2}\right)\right\}$$

and, from Gradshteyn & Ryzhik (1994, p1182), we have:

$$\tilde{\eta}_1(s) = \frac{2}{A}\frac{A}{s+\sqrt{s^2-A^2}}$$

which we can write as $\tilde{\eta}_1(s) = \left(s-\sqrt{s^2-A^2}\right)/(2\alpha\lambda c)$. Thus, we have:

$$\begin{aligned}\tilde{\eta}_1(s)\tilde{\eta}_2(s) &= \frac{\sqrt{\alpha\lambda/c}}{2\alpha\lambda c}\frac{A}{\sqrt{s^2-A^2}}\frac{s-\sqrt{s^2-A^2}}{s+\sqrt{s^2-A^2}} \exp\left\{\frac{u}{2c}\left(s-\sqrt{s^2-A^2}\right)\right\} \\ &= \frac{1}{2c\sqrt{\alpha\lambda c}}\frac{A}{\sqrt{s^2-A^2}}\frac{A^2}{\left(s+\sqrt{s^2-A^2}\right)^2} \exp\left\{\frac{u}{2c}\left(s-\sqrt{s^2-A^2}\right)\right\}\end{aligned}$$

which, by Result 4.1, is the Laplace transform of:

$$\frac{A}{2c\sqrt{\alpha\lambda c}}\frac{t}{t+u/c} I_2\left(\sqrt{4\alpha\lambda c}\sqrt{t^2+\frac{u}{c}t}\right) = \frac{t}{ct+u} I_2\left(\sqrt{4\alpha\lambda t(u+ct)}\right).$$

Hence, formula (10) becomes:

$$\lambda e^{-\alpha u-(\lambda+\alpha c)t}\frac{ct}{ct+u} I_2\left(\sqrt{4\alpha\lambda t(u+ct)}\right)$$

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

and we obtain:

$$w(u,t) = \lambda e^{-\alpha u-(\lambda+\alpha c)t}\left(I_0\left(\sqrt{4\alpha\lambda t(u+ct)}\right) - \frac{ct}{ct+u} I_2\left(\sqrt{4\alpha\lambda t(u+ct)}\right)\right) \qquad (11)$$

a result derived by Dickson *et al.* (2005) by complex inversion of the Laplace transform with respect to t of $w(u,t)$.

5. Erlang(2) Claims

Let us now consider the situation when $\overline{F}(x) = e^{-\alpha x}(1+\alpha x)$, so that:

$$g(x,t) = \sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2n}x^{2n-1}e^{-\alpha x}}{\Gamma(2n)}.$$

We know from Cheung *et al.* (2007) that:

$$w_{Y,T}(u,y,t) = k(u,t)\alpha e^{-\alpha y} + h(u,t)\alpha^2 y e^{-\alpha y} \qquad (12)$$

and the analysis below provides formulae for $h(u,t)$ and $k(u,t)$. We remark that solutions for $h(u,t)$ and $k(u,t)$ were obtained by Dickson (2008), using a very different approach. The formulae which we obtain below are quite different for $u > 0$, but the same for $u = 0$. However, the approach presented here also leads to alternative formulae for $h(0,t)$ and $k(0,t)$. These formulae are stated without proof at the end of this section.

We start by finding $w_{Y,T}(0,y,t)$. From equation (7) we have:

$$w_{Y,T}(0,y,t) = \lambda e^{-\lambda t}f(y+ct) + \frac{\lambda}{c}\int_0^{ct}\frac{x}{t}g(ct-x,t)f(y+x)dx,$$

and we will write:

$$\int_0^{ct}\frac{x}{t}g(ct-x,t)f(y+x)dx = \int_0^{ct}\frac{ct-x}{t}g(x,t)f(y+ct-x)dx.$$

Now:

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$
\begin{aligned}
&c\int_0^{ct} g(x,t)f(y+ct-x)dx \\
&= c\int_0^{ct}\sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2n}x^{2n-1}e^{-\alpha x}}{\Gamma(2n)}\alpha^2(y+ct-x)e^{-\alpha(y+ct-x)}dx \\
&= c\sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2(n+1)}}{\Gamma(2n)}e^{-\alpha(y+ct)}\int_0^{ct} x^{2n-1}(y+ct-x)dx \\
&= c\sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2(n+1)}}{\Gamma(2n)}e^{-\alpha(y+ct)}\left[\left((y+ct)\frac{(ct)^{2n}}{2n}-\frac{(ct)^{2n+1}}{2n+1}\right)\right] \\
&= c\sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2(n+1)}}{\Gamma(2n)}e^{-\alpha(y+ct)}\left[y\frac{(ct)^{2n}}{2n}+\frac{(ct)^{2n+1}}{2n(2n+1)}\right] \qquad (13)
\end{aligned}
$$

and similar manipulations give:

$$
\begin{aligned}
&\int_0^{ct}\frac{x}{t}g(x,t)f(y+ct-x)dx \\
&= \sum_{n=1}^{\infty} e^{-\lambda t}\frac{\lambda^n t^{n-1}}{n!}\frac{\alpha^{2(n+1)}}{\Gamma(2n)}e^{-\alpha(y+ct)}\left[y\frac{(ct)^{2n+1}}{2n+1}+\frac{(ct)^{2n+2}}{(2n+1)(2n+2)}\right]. \qquad (14)
\end{aligned}
$$

Combining the terms in y in (13) and (14) gives:

$$
\begin{aligned}
&\sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2(n+1)}}{\Gamma(2n)}e^{-\alpha(y+ct)}\left[cy\frac{(ct)^{2n}}{2n}-\frac{y}{t}\frac{(ct)^{2n+1}}{2n+1}\right] \\
&= c\alpha^2 y e^{-\alpha y}\sum_{n=1}^{\infty} e^{-(\lambda+c\alpha)t}\frac{(\lambda t)^n}{n!}\frac{(c\alpha t)^{2n}}{(2n+1)!}
\end{aligned}
$$

and combining the remaining terms gives:

$$
\begin{aligned}
&\sum_{n=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^n}{n!}\frac{\alpha^{2(n+1)}}{\Gamma(2n)}e^{-\alpha(y+ct)}\left[\frac{c(ct)^{2n+1}}{2n(2n+1)}-\frac{(ct)^{2n+2}}{t(2n+1)(2n+2)}\right] \\
&= 2c\alpha e^{-\alpha y}\sum_{n=1}^{\infty} e^{-(\lambda+c\alpha)t}\frac{(\lambda t)^n}{n!}\frac{(\alpha ct)^{2n+1}}{(2n+2)!}.
\end{aligned}
$$

Hence:

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$
\begin{aligned}
w_{Y,T}(0, y, t) &= \lambda e^{-\lambda t}\alpha^2(y + ct)e^{-\alpha(y+ct)} \\
&\quad + \frac{\lambda}{c}c\alpha^2 y e^{-\alpha y}\sum_{n=1}^{\infty} e^{-(\lambda+c\alpha)t}\frac{(\lambda t)^n}{n!}\frac{(c\alpha t)^{2n}}{(2n+1)!} \\
&\quad + \frac{\lambda}{c}2c\alpha e^{-\alpha y}\sum_{n=1}^{\infty} e^{-(\lambda+c\alpha)t}\frac{(\lambda t)^n}{n!}\frac{(\alpha ct)^{2n+1}}{(2n+2)!} \\
&= \lambda\alpha^2 y e^{-\alpha y}\sum_{n=0}^{\infty} e^{-(\lambda+c\alpha)t}\frac{(\lambda t)^n}{n!}\frac{(c\alpha t)^{2n}}{(2n+1)!} \\
&\quad + 2\lambda\alpha e^{-\alpha y}\sum_{n=0}^{\infty} e^{-(\lambda+c\alpha)t}\frac{(\lambda t)^n}{n!}\frac{(\alpha ct)^{2n+1}}{(2n+2)!}. \qquad (15)
\end{aligned}
$$

Noting that:

$$(2n+1)! = 4^n(1)_n\left(\tfrac{3}{2}\right)_n$$

and

$$(2n+2)! = 2\left[4^n\left(\tfrac{3}{2}\right)_n(2)_n\right]$$

where $(a)_n = \Gamma(a+n)/\Gamma(a)$ is Pochhammer's symbol, we can write:

$$
\begin{aligned}
w_{Y,T}(0, y, t) &= \alpha^2 y e^{-\alpha y}\lambda e^{-(\lambda+c\alpha)t}\,{}_0F_2\left(1, \frac{3}{2}; \frac{\lambda(\alpha c)^2 t^3}{4}\right) \\
&\quad + \alpha e^{-\alpha y}\lambda\alpha ct e^{-(\lambda+c\alpha)t}\,{}_0F_2\left(\frac{3}{2}, 2; \frac{\lambda(\alpha c)^2 t^3}{4}\right) \qquad (16)
\end{aligned}
$$

where:

$${}_0F_q(C_1, C_2, \ldots, C_q; Z) = \sum_{m=0}^{\infty}\frac{1}{(C_1)_m(C_2)_m\ldots(C_q)_m}\frac{Z^m}{m!}$$

is a generalised hypergeometric function. This form of solution is convenient for computational purposes.

Consider now $w_{Y,T}(u, y, t)$ and the first two terms in formula (9):

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$\lambda e^{-\lambda t} f(u+ct+y) + \lambda \int_0^{u+ct} g(u+ct-x,t) f(x+y)dx$$

$$= \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\alpha^2(u+ct+y)$$

$$+ \lambda \sum_{m=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^m}{m!}\int_0^{u+ct} f^{m^*}(x) f(u+ct+y-x)dx$$

$$= \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\alpha^2(u+ct+y)$$

$$+ \lambda \sum_{m=1}^{\infty} e^{-\lambda t}\frac{(\lambda t)^m}{m!}\int_0^{u+ct} \frac{\alpha^{2m}x^{2m-1}e^{-\alpha x}}{\Gamma(2m)}\alpha^2(u+ct+y-x)e^{-\alpha(u+ct+y-x)}dx$$

$$= \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\alpha^2(u+ct+y)$$

$$+ \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=1}^{\infty}\frac{(\lambda t)^m}{m!}\frac{\alpha^{2m+2}}{\Gamma(2m)}\int_0^{u+ct} x^{2m-1}(u+ct+y-x)dx$$

$$= \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\alpha^2(u+ct+y)$$

$$+ \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=1}^{\infty}\frac{(\lambda t)^m}{m!}\frac{\alpha^{2m+2}}{\Gamma(2m)}\left[(u+ct+y)\frac{(u+ct)^{2m}}{2m} - \frac{(u+ct)^{2m+1}}{2m+1}\right]$$

$$= \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\alpha^2(u+ct+y)$$

$$+ \lambda y e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=1}^{\infty}\frac{(\lambda t)^m}{m!}\frac{\alpha^{2m+2}}{\Gamma(2m)}\frac{(u+ct)^{2m}}{2m}$$

$$+ \lambda e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=1}^{\infty}\frac{(\lambda t)^m}{m!}\frac{\alpha^{2m+2}}{\Gamma(2m)}\frac{(u+ct)^{2m+1}}{2m(2m+1)}$$

$$= \lambda\alpha^2 y e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=0}^{\infty}\frac{(\lambda t)^m}{m!}\frac{[\alpha(u+ct)]^{2m}}{(2m)!}$$

$$+ \lambda\alpha e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=0}^{\infty}\frac{(\lambda t)^m}{m!}\frac{[\alpha(u+ct)]^{2m+1}}{(2m+1)!}$$

$$= \alpha^2 y e^{-\alpha y}\lambda e^{-\alpha u-(\lambda+\alpha c)t}\,{}_0F_2\left(\tfrac{1}{2},1;\lambda t(\alpha(u+ct))^2/4\right)$$

$$+ \alpha e^{-\alpha y}\alpha(u+ct)\lambda e^{-\alpha u-(\lambda+\alpha c)t}\,{}_0F_2\left(1,\tfrac{3}{2};\lambda t(\alpha(u+ct))^2/4\right). \tag{17}$$

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

Next, (minus) the third term of formula (9) is:

$$
\begin{aligned}
& c\int_0^t g(u+c(t-s),t-s)w_{Y,T}(0,y,s)ds \\
&= c\int_0^t g(u+c(t-s),t-s)\lambda\alpha^2 y e^{-\alpha y}\sum_{n=0}^{\infty} e^{-(\lambda+c\alpha)s}\frac{(\lambda s)^n}{n!}\frac{(c\alpha s)^{2n}}{(2n+1)!}ds \\
&\quad + c\int_0^t g(u+c(t-s),t-s)2\lambda\alpha e^{-\alpha y}\sum_{n=0}^{\infty} e^{-(\lambda+c\alpha)s}\frac{(\lambda s)^n}{n!}\frac{(\alpha c s)^{2n+1}}{(2n+2)!}ds \\
&= c\lambda\alpha^2 y e^{-\alpha y}\int_0^t \sum_{m=1}^{\infty} e^{-\lambda(t-s)}\frac{(\lambda(t-s))^m}{m!}f^{m*}(u+c(t-s)) \\
&\quad \times \sum_{n=0}^{\infty} e^{-(\lambda+c\alpha)s}\frac{(\lambda s)^n}{n!}\frac{(c\alpha s)^{2n}}{(2n+1)!}ds \\
&\quad + 2c\lambda\alpha e^{-\alpha y}\int_0^t \sum_{m=1}^{\infty} e^{-\lambda(t-s)}\frac{(\lambda(t-s))^m}{m!}f^{m*}(u+c(t-s)) \\
&\quad \times \sum_{n=0}^{\infty} e^{-(\lambda+c\alpha)s}\frac{(\lambda s)^n}{n!}\frac{(\alpha c s)^{2n+1}}{(2n+2)!}ds \\
&= c\lambda\alpha^2 y e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=1}^{\infty}\frac{\lambda^m\alpha^{2m}}{m!\Gamma(2m)} \\
&\quad \times \sum_{n=0}^{\infty}\frac{\lambda^n}{n!}\frac{(\alpha c)^{2n}}{(2n+1)!}\int_0^t (t-s)^m(u+c(t-s))^{2m-1}s^{3n}ds \\
&\quad + 2c\lambda\alpha e^{-\alpha(u+y)-(\lambda+\alpha c)t}\sum_{m=1}^{\infty}\frac{\lambda^m\alpha^{2m}}{m!\Gamma(2m)} \\
&\quad \times \sum_{n=0}^{\infty}\frac{\lambda^n}{n!}\frac{(\alpha c)^{2n+1}}{(2n+2)!}\int_0^t (t-s)^m(u+c(t-s))^{2m-1}s^{3n+1}ds.
\end{aligned}
$$

Using a binomial expansion of $(u+c(t-s))^m$, it is straightforward to show that for $r \geq 0$ and $m, n \geq 1$:

$$
\int_0^t (t-s)^n(u+c(t-s))^m s^r ds = \sum_{i=0}^{m}\binom{m}{i}c^i u^{m-i}t^{n+i+r+1}\frac{r!(n+i)!}{(n+r+i+1)!}.
$$

Thus, we can use expression (17) and the above development to write $h(u,t)$ and $k(u,t)$ from formula (12) as:

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

$$h(u,t)=\lambda e^{-\alpha u-(\lambda+\alpha c)t}\Bigg({}_0F_2\left(\tfrac{1}{2},1;\lambda t(\alpha(u+ct))^2/4\right)$$
$$-\frac{ct}{u}\sum_{m=1}^{\infty}\frac{\left(\lambda t\alpha^2u^2\right)^m}{m!\Gamma(2m)}\sum_{n=0}^{\infty}\frac{\left(\lambda\alpha^2c^2t^3\right)^n}{n!(2n+1)!}\sum_{i=0}^{2m-1}\binom{2m-1}{i}\frac{(ct/u)^i(3n)!(m+i)!}{(m+3n+i+1)!}\Bigg)$$

and

$$k(u,t)=\lambda e^{-\alpha u-(\lambda+\alpha c)t}\Bigg(\alpha(u+ct)\ {}_0F_2\left(1,\tfrac{3}{2};\lambda t(\alpha(u+ct))^2/4\right)$$
$$-\frac{2\alpha(ct)^2}{u}\sum_{m=1}^{\infty}\frac{\left(\lambda t\alpha^2u^2\right)^m}{m!\Gamma(2m)}\sum_{n=0}^{\infty}\frac{\left(\lambda\alpha^2c^2t^3\right)^n}{n!(2n+2)!}$$
$$\times\sum_{i=0}^{2m-1}\binom{2m-1}{i}\frac{(ct/u)^i(3n+1)!(m+i)!}{(m+3n+i+2)!}\Bigg).$$

Although these formulae for $h(u,t)$ and $k(u,t)$ are not particularly elegant, their implementation is straightforward, provided that we truncate the infinite sums at a suitable point. Given the form of the solution for $w(u,t)$ in the previous section, it would appear possible that the solutions for $h(u,t)$ and $k(u,t)$ could be expressed entirely in terms of generalised hypergeometric functions. Indeed, it can be shown that, by setting $u=0$ in the third term of formula (9), we can obtain an alternative to formula (16), namely:

$$w_{Y,T}(0,y,t)=h(0,t)\alpha^2ye^{-\alpha y}+k(0,t)\alpha e^{-\alpha y}$$

where:

$$h(0,t)=\lambda e^{-(\lambda+\alpha c)t}\left({}_0F_2\left(\frac{1}{2},1;\frac{\lambda\alpha^2c^2t^3}{4}\right)-\frac{\lambda\alpha^2c^2t^3}{3}\,{}_0F_2\left(2,\frac{5}{2};\frac{\lambda\alpha^2c^2t^3}{4}\right)\right)$$

and

$$k(0,t)=\alpha ct\lambda e^{-(\lambda+\alpha c)t}\left({}_0F_2\left(1,\frac{3}{2};\frac{\lambda\alpha^2c^2t^3}{4}\right)-\frac{\lambda\alpha^2c^2t^3}{12}\,{}_0F_2\left(\frac{5}{2},3;\frac{\lambda\alpha^2c^2t^3}{4}\right)\right).$$

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

6. Concluding Remarks

Liu & Zhao (2007) consider the joint distribution of the surplus prior to ruin, the deficit at ruin and the time to ruin in the compound binomial model. If, instead, we consider a discrete time compound Poisson risk model where the premium income per unit time is one and claims are distributed on the integers, as described in Dickson & Waters (1992), then there are counterparts of formulae in Sections 2 and 3 for this model. However, from a computational point of view, the recursive calculation scheme of Dickson & Waters (1992) (which is for the joint distribution of the deficit at ruin and the time to ruin, but can be extended to include the surplus prior to ruin) seems preferable.

Dickson & Willmot (2005) derived an expression for $w(u, t)$ when the individual claim amount distribution is an infinite mixture of Erlang distributions with the same scale parameter. Each of the individual claim amount distributions in Sections 4 and 5 is a special case of this distribution, but it seems rather difficult to condense Dickson & Willmot's expression to give a simple formula for $w(u, t)$ for comparison with the formulae presented in Sections 4 and 5. Indeed, we would conjecture that, if simplification of Dickson & Willmot's expression was possible, we would not obtain the formulae for $w(u, t)$ in Sections 4 and 5. To a large extent this conjecture is based on the fact that, in the case of exponential individual claims, Drekic & Willmot (2003) obtain a formula for $w(u, t)$ by inverting its Laplace transform, and this formula is expressed in terms of an infinite sum, where the summand contains a Bessel function. Whilst their result can be shown to be the same as formula (11) when $u = 0$, we are unable to show that the same is true for $u > 0$. Whilst Dickson & Willmot's formula has wide applicability — see Willmot & Woo (2007) — the formulae for $w(u, t)$ presented in Sections 4 and 5 seem preferable from a programming point of view for these individual claim amount distributions. Indeed, in the case of exponential claims, we can use supplied functions in a package such as Mathematica to ease computation.

The application of formulae (6) and (9) for individual claim amount distributions other than those considered in Sections 4 and 5 is feasible. We have chosen our examples for ease of presentation. The ideas extend to Erlang(n, α) claims where $n > 2$. In this case it is straightforward to solve directly for $w(u, t)$, while the derivation of $w(u, y, t)$ is a little tedious. We state, without proof, that, when the individual claim amount distribution is Erlang(n, α), the joint density of the time to ruin and the deficit at ruin is:

$$w(u, y, t) = \sum_{i=1}^{n} h_i(u, t) \frac{\alpha^i y^{i-1} e^{-\alpha y}}{\Gamma(i)}$$

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

where:

$$h_i(u,t) = \lambda e^{-\alpha u-(\lambda+c\alpha)t}\left(\sum_{m=0}^{\infty}\frac{(\lambda t)^m}{m!}\frac{(\alpha(u+ct))^{n(m+1)-i}}{\Gamma(n(m+1)-i+1)}\right.$$
$$-\frac{ct(n-i+1)}{u}\sum_{m=1}^{\infty}\frac{(\lambda t)^m(\alpha u)^{nm}}{m!\Gamma(nm)}\sum_{r=0}^{\infty}\frac{(\lambda t)^r}{r!}\frac{(\alpha ct)^{n(r+1)-i}}{(n(r+1)-i+1)!}$$
$$\left.\times\sum_{k=0}^{nm-1}\binom{nm-1}{k}\left(\frac{ct}{u}\right)^k\frac{(n(r+1)+r-i)!(m+k)!}{(m+k+n(r+1)+r-i+1)!}\right).$$

We know from Cheung *et al.* (2007) that, when the individual claim amount distribution is a finite mixture of Erlang distributions with the same scale parameter, then the joint density $w(u, y, t)$ decomposes in a similar way to the case of Erlang(n,α) claims. It therefore seems that the techniques presented above will equally apply to this individual claim amount distribution.

Acknowledgement

This paper has benefitted from helpful comments from an anonymous referee.

References

Abramowitz, M. & Stegun, I.A. (1965). *Handbook of mathematical functions.* Dover, New York.

Cheung, E.C.K., Dickson, D.C.M. & Drekic, S. (2007). Moments of discounted dividends for a threshhold strategy in the compound Poisson risk model. Unpublished manuscript.

Dickson, D.C.M. (2008). Some explicit solutions for the joint density of the time of ruin and the deficit at ruin. *ASTIN Bulletin*, **38**, 259-276.

Dickson, D.C.M. & Waters, H.R. (1992). The probability and severity of ruin in finite and in infinite time. *ASTIN Bulletin*, **22**, 177-190.

Dickson, D.C.M. & Waters, H.R. (2006). Optimal dynamic reinsurance. *ASTIN Bulletin*, **36**, 415-432.

Dickson, D.C.M. & Willmot, G.E. (2005). The density of the time to ruin in the classical Poisson risk model. *ASTIN Bulletin*, **35**, 45-60.

Dickson, D.C.M., Hughes, B.D. & Lianzeng, Z. (2005). The density of the time to ruin for a Sparre Andersen process with Erlang arrivals and exponential claims. *Scandinavian Actuarial Journal*, **5**, 358-376.

Drekic, S. & Willmot, G.E. (2003). On the density and moments of the time to ruin with exponential claims. *ASTIN Bulletin*, **33**, 11-21.

Erdélyi, A. (ed.) (1954). *Tables of integral transforms, Volume 1*. McGraw-Hill, New York.

Gerber, H.U. (1979). *An introduction to mathematical risk theory*. S.S. Huebner Foundation, Philadelphia, PA.

Gerber, H.U. & Shiu, E.S.W. (1997). The joint distribution of the time of ruin, the surplus immediately before ruin, and the deficit at ruin. *Insurance: Mathematics & Economics*, **21**, 129-137.

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press

GRADSHTEYN, I.S. & RYZHIK, I.M. (1994). *Table of integrals, series, and products* (Fifth edition). Academic Press, San Diego.

LIU, G. & ZHAO, J. (2007). Joint distributions of some actuarial random vectors in the compound binomial model. *Insurance: Mathematics & Economics*, **40**, 95-103.

PRABHU, N.U. (1961). On the ruin problem of collective risk theory. *Annals of Mathematical Statistics*, **32**, 757-764.

SEAL, H.L. (1978). *Survival probabilities*. John Wiley & Sons, New York.

WILLMOT, G.E. & WOO, J.K. (2007). On the class of Erlang mixtures with risk theoretic applications. *North American Actuarial Journal*, **11**(2), 99-115.

WU, R., WANG, G. & WEI, L. (2003). Joint distributions of some actuarial random vectors containing the time of ruin. *Insurance: Mathematics & Economics*, **33**, 147-161.

https://doi.org/10.1017/S1748499500000348 Published online by Cambridge University Press