Figures

I.1 Mosaic of photographic exposures of the A2713 field. (Reproduced with permission) *page* 3
1.1 Detail of the negative of a calibrated photographic exposure of a source on the sky, a grid of 21 × 19 pixels (picture elements). 33
2.1 Staff scientists and visiting astronomers at the 2.2-meter telescope at La Silla Observatory. (Photograph: Götz Hoeppe) 62
2.2 At twilight in the control room of the 2.2-meter telescope at La Silla Observatory. (Photograph: Götz Hoeppe) 65
2.3 At the BOB terminal. (Photograph: Götz Hoeppe) 66
2.4 Visiting astronomers inspecting air mass plots pertaining to three observing projects. (Photograph: Götz Hoeppe) 69
2.5 Air mass plot pertaining to Mary's observing program. (Photograph: Götz Hoeppe) 71
3.1 Flowchart of the MAMBO research group's data reductions as taken from a conference presentation of one of its members. (Reproduced with the authors' permission) 84
3.2 Scheme depicting basic steps of the MAMBO team's spectral energy distribution template-fitting technique. 85
3.3 Otfried pointing at an offset between galaxy photometric measurements and a spectral template (continuous line). (Photograph: Götz Hoeppe) 88
3.4 Otfried and Nadine visually assessing their flatfield frames. (Photograph: Götz Hoeppe) 90
3.5 Visualization of the differences between instrumental magnitudes and 2MASS catalog entries for stars in the A2713 field. (Photograph: Götz Hoeppe) 91
3.6 Otto's recipe to correct for the artifactual ring in the flatfield. (Photograph: Götz Hoeppe) 92

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 11 Nov 2026 at 10:03:24, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/9781009686754

3.7 Nadine's draft histogram, showing the number of galaxies as a function of redshift for all objects detected in the A2713 field. 97
3.8 Revised histogram as it appeared in the published paper. (© American Astronomical Society. Reproduced with permission) 98
3.9 Comparison of the galaxy number counts in a set of deep fields. (Figure 12 of Capak et al. 2007, 114; © American Astronomical Society. Reproduced with permission) 105
4.1 Nadine's draft histogram, showing the number of galaxies as a function of redshift for all objects detected in the A2713 field. See Transcript 4.2. 111
4.2 Peter explains a scatterplot that shows the redshifts of galaxies (vertical axis) over their red (R band) magnitudes (horizontal axis). See Transcript 4.4. (Photograph: Götz Hoeppe) 114
4.3 Peter demonstrates how Nadine may use the scatterplot to estimate the completeness of her galaxy catalog. See Transcripts 4.5 and 4.6. (Photograph: Götz Hoeppe) 116
4.4 Hertzsprung–Russell Diagram of four million stars from measurements of the European Space Agency's Gaia spacecraft. (© ESA/Gaia/DPAC 2018) 120
4.5 Figures of Arjen van der Wel's manuscript as discussed in the text. (© American Astronomical Society. Reproduced with permission) 125
4.6 Images of the galaxies in van der Wel's sample, made using Hubble Space Telescope I, J, and H exposures. (© American Astronomical Society/NASA/ESA/Hubble Space Telescope. Reproduced with permission) 128
4.7 Otto's sketch of the sequence of arithmetic operations (left) that Nadine was to apply to remedy her flatfield issue. See also Figure 3.6. (Photograph: Götz Hoeppe) 133
5.1 Presentation slide from David Spergel's lecture at New York University, showing two grayscale pixel maps of the microwave background fluctuation pattern of a patch in the sky. (© D. Spergel/Princeton University/ESA/Planck 2014) 139
5.2 False-color image of one of Nadine's flatfields. (Photograph: Götz Hoeppe) 148
5.3 Nadine briefing Otfried about her assessment of artifacts in the flatfield exposures, using paper printouts to summarize her findings. (Photograph: Götz Hoeppe) 149

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 11 Nov 2026 at 10:03:24, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/9781009686754

5.4 Otfried's schematic characterization and quantification of artifacts in the flatfield frames on his office blackboard. (Photograph: Götz Hoeppe) 150
8.1 The first entries of George Abell's (1958) catalog of 2,712 galaxy clusters in the northern and equatorial sky. (© American Astronomical Society. Reproduced with permission) 209
8.2 Structure of the 2MASS Redshift Survey (2MRS) catalog. (© American Astronomical Society. Reproduced with permission) 210

Downloaded from https://www.cambridge.org/core. UNIVERSIDAD AUTONOMA METROPOLITANA, on 11 Nov 2026 at 10:03:24, subject to the Cambridge Core terms of use, available at https://www.cambridge.org/core/terms. https://doi.org/10.1017/9781009686754