Combinatorics, Probability and Computing (2007) 16, 733–746. © 2007 Cambridge University Press
doi:10.1017/S0963548306008388 First published online 23 January 2007 Printed in the United Kingdom

On the Chromatic Number of Random Graphs with a Fixed Degree Sequence

ALAN FRIEZE[1†], MICHAEL KRIVELEVICH[2‡] and CLIFF SMYTH[3]

[1]Department of Mathematical Sciences, Carnegie Mellon University, Pittsburgh PA15213, USA
(e-mail: alan@random.math.cmu.edu)

[2]Department of Mathematics, Raymond and Beverly Sackler Faculty of Exact Sciences,
Tel Aviv University, Tel Aviv 69978, Israel
(e-mail: krivelev@post.tau.ac.il)

[3]Department of Mathematics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
(e-mail: csmyth@math.mit.edu)

Received 25 January 2006; revised 14 November 2006

Let $\mathbf{d} = 1 \leqslant d_1 \leqslant d_2 \leqslant \cdots \leqslant d_n$ be a non-decreasing sequence of n positive integers, whose sum is even. Let $\mathscr{G}_{n,\mathbf{d}}$ denote the set of graphs with vertex set $[n] = \{1, 2, \ldots, n\}$ in which the degree of vertex i is d_i. Let $G_{n,\mathbf{d}}$ be chosen uniformly at random from $\mathscr{G}_{n,\mathbf{d}}$. Let $d = (d_1 + d_2 + \cdots + d_n)/n$ be the average degree. We give a condition on $\mathbf{d}$ under which we can show that w.h.p. the chromatic number of $\mathscr{G}_{n,\mathbf{d}}$ is $\Theta(d/\ln d)$. This condition is satisfied by graphs with exponential tails as well those with power law tails.

1. Introduction

Let $\mathbf{d} = 1 \leqslant d_1 \leqslant d_2 \leqslant \cdots \leqslant d_n$ be a fixed non-decreasing sequence of n positive integers, whose sum is even. Let $\mathscr{G}_{n,\mathbf{d}}$ denote the set of graphs with vertex set $[n] = \{1, 2, \ldots, n\}$ in which the degree of vertex i is d_i. Let $G_{n,\mathbf{d}}$ be chosen uniformly at random from $\mathscr{G}_{n,\mathbf{d}}$. When $d_i = r$ for $i \in [n]$ then this models a random r-regular graph $G_{n,r}$ and there is a large literature on this subject. We refer the reader to the survey by Wormald [18] for an excellent summary. By now we know much about the structure of random regular graphs.

For general $\mathbf{d}$, less is known. In many, but not all, cases we can estimate $|\mathscr{G}_{n,\mathbf{d}}|$. See Bender and Canfield [5] and McKay and Wormald [15, 13]. We have the *configuration model* to study them: see Bollobás [6]. We know something of their connectivity properties: see Molloy and Reed [16, 17]. (See also Cooper and Frieze [8] for the connectivity properties

† Supported in part by NSF grant CCR-0200945.
‡ Research supported in part by USA–Israel BSF grant 2002-133 and by grants 64/01 and 526/05 from the Israel Science Foundation.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

of random *digraphs* with a fixed degree sequence.) They have been used in the context of *massive graph models* of telephone networks and the Web: see Aiello, Chung and Lu [3].

In this paper we will concern ourselves with the chromatic number of $G_{n,\mathbf{d}}$. What should we expect? For r-regular graphs it is known that when r is large $\chi(G_{n,r}) \sim \frac{r}{2\ln r}$ w.h.p.;[1] see Frieze and Łuczak [10] and Achlioptas and Moore [1] (see also Cooper, Frieze, Reed and Riordan [9] for extensions of some of these results to the case where $r = r(n) \leqslant n^{1-\eta}$, for an arbitrary small constant $\eta > 0$). For the random graph $G_{n,p}$ where the average degree $d = (n-1)p$ is large, it is known that $\chi(G_{n,p}) \sim \frac{d}{2\ln d}$: see Bollobás [7], Łuczak [12] and Achlioptas and Naor [2]. So if we let

$$d = \frac{d_1 + d_2 + \cdots + d_n}{n}$$

be the average degree in $G_{n,\mathbf{d}}$, then we might hope to prove that $\chi(G_{n,\mathbf{d}}) \sim \frac{d}{2\ln d}$ w.h.p. This is too much to expect given the variety of possible degree sequences. Indeed if $v = \frac{n}{\ln n}$ and

$$d_i = \begin{cases} 1 & i \leqslant n - v, \\ d\ln n & i > n - v, \end{cases}$$

then the average degree $\sim d$, but w.h.p. the sub-graph H induced by the v largest degree vertices has average degree close to $d\ln n$ when d is large. Then w.h.p. $\chi(H) \sim \frac{d\ln n}{2\ln\ln n}$ for $\ln d = o(\ln\ln n)$ and so we will have to be less ambitious in our goals.[2] Let

$$D_k = d_n + d_{n-1} + \cdots + d_{n-k+1}$$

be the sum of the k largest degrees.

Let

$$M_1 = D_n = dn \quad \text{and} \quad M_2 = \sum_{i=1}^{n} d_i(d_i - 1) \leqslant \Delta M_1 \quad \text{where } \Delta = d_n.$$

Theorem 1.1.

1 *Suppose that there exist constants* $1/2 < \alpha < 1$, $\epsilon, K_0 > 0$ *and* $\omega = \omega(n) \to \infty$ *such that:*

(a)

$$D_k \leqslant K_0 dn(k/n)^{\alpha} \tag{1.1}$$

for $k \leqslant \epsilon n$,

(b) $\Delta^5 \leqslant M_2/\omega$.

Then there exists b_1 *dependent only on* α, ϵ, K_0 *such that w.h.p.*

$$\chi(G_{n,\mathbf{d}}) \leqslant b_1 \frac{d}{\ln d}.$$

[1] A sequence of events $\mathscr{E}_n$ occurs *with high probability* (w.h.p.) if $\lim_{n\to\infty} \mathbf{Pr}(\mathscr{E}_n) = 1$.

[2] The methods of [9] can be used to verify this claim.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

2 *Suppose only that* $\Delta^4 \leqslant M_1/\omega$ (*a weaker condition than* 1(b))*; then there exists* b_2 *such that w.h.p.*

$$\chi(G_{n,\mathbf{d}}) \geqslant b_2 \frac{d}{\ln d}.$$

Notice that if condition (a) of the theorem holds for α, ϵ, K_0 then it also holds when K_0 is replaced by a larger constant. As we increase K_0 our bound b_1 will decrease. We are therefore justified in assuming throughout that K_0 is sufficiently large that some inequalities are valid.

Condition (b) is chosen so that we can use the results of [14]. It may be possible to prove our results under the less stringent conditions of [13], but there are difficulties, as will be pointed out later.

A referee has suggested that the quantity $d' = \max_k D_k^2/dnk$ (which can be viewed as the probabilistic version of the 'maximum average degree') might be a better predictor of chromatic number. This could well be true and it makes for an interesting research question. We also suspect that if $d_i \sim df(i/n)$, where f is some 'nice' real valued function on $[0,1]$ such that $\int_0^1 f(x)dx = 1$, then we might be able to determine the chromatic number of $G_{n,\mathbf{d}}$ asymptotically.

The proof of the upper bound is given in Section 3. The proof of the lower bound is given in Section 4. We have made no attempt to optimize constants.

2. Degree sequences that satisfy (1.1)

It is natural to ask whether there many types of degree sequence that satisfy the conditions of the first part of the theorem.

We first consider a degree sequence with a *power law* tail. For integer $\ell \geqslant 1$ we let ν_ℓ denote the number of vertices of degree ℓ. Our assumption is that there are some constants $A > 0$ and $\gamma > 3$ such that, for $\ell \geqslant (A/\epsilon)^{1/(\gamma-1)}$,

$$\nu_\ell \leqslant \begin{cases} 0 & \ell \leqslant 1, \\ \lfloor Ad\ell^{-\gamma} n \rfloor & 2 \leqslant \ell \leqslant n^{1/5}/\ln n, \\ 0 & \ell > n^{1/5}/\ln n. \end{cases}$$

Here we have $\alpha = \frac{\gamma-2}{\gamma-1} > 1/2$.

We have forced condition (b) and so we only need to check condition (a). Suppose now that $k = \kappa n \leqslant \epsilon n$. Then

$$D_k \leqslant Adn \sum_{\ell \geqslant r} \ell^{-(\gamma-1)} = (1+\delta_r)\frac{Adn}{\gamma-2} r^{2-\gamma},$$

where $\delta_r \to 0$ as $r \to \infty$, and

$$r = \min\left\{ \rho : Ad \sum_{\ell \geqslant \rho} \ell^{-\gamma} \geqslant \kappa \right\} = (1+\delta'_\kappa)\left(\frac{Ad}{(\gamma-1)\kappa}\right)^{1/(\gamma-1)},$$

where $\delta'_\kappa \to 0$ as $\kappa \to 0$. Thus,

$$D_k \leqslant (1+\delta_r)(1+\delta'_\kappa)^{2-\gamma}(\gamma-2)^{-1} A \left(\frac{\gamma-1}{Ad}\right)^{\alpha} dn\kappa^{\alpha},$$

and we have the condition of Theorem 1.1.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

A similar argument holds if we assume that our degree sequence has an *exponential* tail, namely, for some constants $A > 0$ and $0 < \epsilon \ll \gamma < 1$, we have for $\ell \geqslant \lfloor \ln_{1/\gamma}(Ad/\epsilon) \rfloor$

$$\nu_\ell \leqslant Ad\gamma^\ell n.$$

Note that w.h.p. the degree sequence of $G_{n,p}$, $p = c/n$, c constant, satisfies this condition.

Suppose now that $k = \kappa n \leqslant \epsilon n$. Then

$$D_k \leqslant Adn \sum_{\ell \geqslant r} \ell \gamma^\ell \leqslant Adn \int_{x=r-1}^{\infty} x\gamma^x dx = \frac{Adn\gamma^{r-1}}{\ln 1/\gamma}\left(r - 1 + \frac{1}{\ln 1/\gamma}\right),$$

where

$$r = \min\left\{\rho : Ad\sum_{\ell \geqslant \rho} \ell\gamma^\ell \geqslant \kappa\right\} \sim \ln_{1/\gamma}\left(\frac{Ad}{\kappa \ln 1/\gamma}\right)$$

as $\epsilon \to 0$.

Thus

$$D_k \leqslant 2\kappa n,$$

and we have the condition of Theorem 1.1 with room to spare.

3. Upper bound

3.1. Configurations

We will work initially in the configuration model and then show how our results can be justified in the uniform model. Let $W = [nd]$ be our set of *points* and let $W_i = [d_1 + \cdots + d_{i-1} + 1, d_1 + \cdots + d_i]$, $i \in [n]$, partition W. The function $\phi : W \to [n]$ is defined by $w \in W_{\phi(w)}$. Given a pairing F (*i.e.*, a partition of W into $m = dn/2$ pairs) we obtain a (multi-)graph G_F with vertex set $[n]$ and an edge $(\phi(u), \phi(v))$ for each $\{u, v\} \in F$. Choosing a pairing F uniformly at random from among all possible pairings of the points of W produces a random (multi-)graph G_F.

This model is valuable because of the following easily proven fact. Suppose $G \in \mathscr{G}_{n,\mathbf{d}}$. Then

$$\mathbf{Pr}(G_F = G \mid G_F \text{ is simple}) = \frac{1}{|\mathscr{G}_{n,\mathbf{d}}|}.$$

It follows that if G is chosen randomly from $\mathscr{G}_{n,\mathbf{d}}$, then for any graph property $\mathscr{P}$

$$\mathbf{Pr}(G \in \mathscr{P}) \leqslant \frac{\mathbf{Pr}(G_F \in \mathscr{P})}{\mathbf{Pr}(G_F \text{ is simple})}. \tag{3.1}$$

Before starting the proof of the upper bound in Theorem 1.1, we introduce some notation that we will use throughout the course of the proof. Let

$$\theta = \max\left\{d^{2/(2\alpha+1)}, \frac{16}{2\alpha - 1}\right\}$$

and

$$\epsilon_1 = \min\left\{\epsilon, \left(\frac{(2\alpha - 1)\theta}{K_1 d}\right)^{2/(2\alpha-1)}\right\} \tag{3.2}$$

for some sufficiently large K_1.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

For a set $U \subseteq V(G_F)$ we denote by $D(U)$ the sum of the degrees of U in G_F.

We first prove the following result.

Lemma 3.1. *With high probability G_F does not contain a set of vertices S with $k = |S| \leqslant \epsilon_1 n$, such that S induces a sub-graph $G[S]$ with minimum degree at least θ.*

Proof. Suppose that there exists a vertex set S of size $k \in [\theta, \epsilon_1 n]$ that induces a graph $G[S]$ with minimum degree at least θ. Let S_1 be the $k/2$ lowest degree vertices of S and let $S_2 = S \setminus S_1$. For vertex sets S, T let $e(S)$ denote the number of edges contained in S and let $e(S : T)$ denote the number of edges joining S and T.

At least one of the following two events must occur:

$\mathscr{E}_1 : e(S_1 : S_2) \geqslant \theta k/4$,
$\mathscr{E}_2 : e(S_1) \geqslant \theta k/8$.

This follows because

$$2e(S_1) + e(S_1 : S_2) = \sum_{v \in S_1} d_{G[S]}(v) \geqslant \theta k/2.$$

Now $k \leqslant \epsilon_1 n$, and so

$$dn - \theta k/2 \geqslant dn - \theta \epsilon_1 n/2 \geqslant \frac{99}{100} dn,$$

provided

$$K_1 \geqslant 16 \left(\frac{1600}{2\alpha - 1} \right)^{\alpha - 1/2}.$$

(We have chosen K_1 sufficiently large that, whichever of $d^{2/(2\alpha+1)}, \frac{16}{2\alpha-1}$ is the larger, $\theta \epsilon_1 < 1/100$.)

And so

$$\begin{aligned}
\mathbf{Pr}(\mathscr{E}_1) &\leqslant \sum_{k=\theta}^{\epsilon_1 n} \sum_{|S|=k} \binom{D(S_1)}{\theta k/4} \left(\frac{D(S_2)}{dn - \theta k/2} \right)^{\theta k/4} \\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n} \binom{n}{k} \binom{D_{k/2}}{\theta k/4} \left(\frac{D_{k/2}}{dn - \theta k/2} \right)^{\theta k/4} \\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n} \left(\frac{en}{k} \right)^k \left(\frac{5eD_{k/2}^2}{\theta k dn} \right)^{\theta k/4} \\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n} \left(\frac{en}{k} \right)^k \left(\frac{5eK_0^2 d}{2^{2\alpha}\theta} \cdot \left(\frac{k}{n} \right)^{2\alpha - 1} \right)^{\theta k/4} \\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n} \left(\left(\frac{k}{n} \right)^{(2\alpha-1)\theta/4 - 1} \left(\frac{5e^2 K_0^2 d}{2^{2\alpha}\theta} \right)^{\theta/4} \right)^k \\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n} \left(\left(\frac{k}{n} \right)^{3(2\alpha-1)\theta/16} \left(\frac{5e^2 K_0^2 d}{2^{2\alpha}\theta} \right)^{\theta/4} \right)^k
\end{aligned}$$

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

$$\leqslant \sum_{k=\theta}^{\epsilon_1 n}\left(\left(\frac{k}{n}\right)^{(2\alpha-1)/4}\frac{5e^2K_0^2}{2^{2\alpha}}\right)^{\theta k/4}$$
$$= o(1), \tag{3.3}$$

if we choose $K_1 \geqslant 100e^4K_0^4$ (we used the assumption $k/n \leqslant \epsilon_1 \leqslant ((2\alpha-1)\theta/(K_1 d))^{2/(2\alpha-1)} \leqslant (16/K_1)^{2/(2\alpha-1)}$ in the last line).

Explanation of (3.3). At least $\theta k/4$ of the points corresponding to vertices of S_1 are to be paired with points corresponding to vertices of S_2. When pairing points, the probability of pairing with a point corresponding to S_2 is never more than $\frac{D(S_2)}{dn-\ell}$, given up to $\ell/2-1$ previous pairings.

In the following estimate, $\ell = \ell(S_1)$ is the total degree of the subgraph $G[S_1]$. Here $\theta k/4 \leqslant \ell \leqslant K_0\epsilon_1^\alpha dn$. Now the maximum degree in S_1 is at most $\frac{D_{k/2}}{k/2} \leqslant K_0 d(2n/k)^{1-\alpha}$ and so, if $a_1, a_2, \dots a_{k/2}$ is the degree sequence of $G[S_1]$ (more explanation below), then

$$\begin{aligned}
\mathbf{Pr}(\mathscr{E}_2) &\leqslant \sum_{k=\theta}^{\epsilon_1 n}\sum_{S_1=\{i_1<\cdots<i_{k/2}\}}\sum_{\ell=\theta k/4}^{K_0\epsilon_1^\alpha dn}\sum_{a_1+\cdots+a_{k/2}=\ell}\frac{\binom{\binom{k/2}{2}}{\ell/2}}{(dn-\ell)^{\ell/2}}\prod_{t=1}^{k/2}d_{i_t}^{a_t}\\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n}\sum_{S_1=\{i_1<\cdots<i_{k/2}\}}\sum_{\ell=\theta k/4}^{K_0\epsilon_1^\alpha dn}\sum_{a_1+\cdots+a_{k/2}=\ell}\left(\frac{k^2}{\ell dn}\right)^{\ell/2}\prod_{t=1}^{k/2}(K_0 d(2n/k)^{(1-\alpha)})^{a_t}\\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n}\sum_{S_1=\{i_1<\cdots<i_{k/2}\}}\sum_{\ell=\theta k/4}^{K_0\epsilon_1^\alpha dn}\sum_{a_1+\cdots+a_{k/2}=\ell}\left(\frac{16K_0^2dk^{2\alpha-1}}{\theta n^{2\alpha-1}}\right)^{\ell/2}\\
&\leqslant \sum_{k=\theta}^{\epsilon_1 n}\sum_{S_1=\{i_1<\cdots<i_{k/2}\}}\sum_{\ell=\theta k/4}^{K_0\epsilon_1^\alpha dn}\left(\frac{64K_0^2dk^{2\alpha-1}}{\theta n^{2\alpha-1}}\right)^{\ell/2}\\
&\leqslant dn\sum_{k=\theta}^{\epsilon_1 n}\binom{n}{k/2}\left(\frac{64K_0^2dk^{2\alpha-1}}{\theta n^{2\alpha-1}}\right)^{\theta k/8}\\
&\leqslant dn\sum_{k=\theta}^{\epsilon_1 n}\left(\frac{2en}{k}\left(\frac{64K_0^2dk^{2\alpha-1}}{\theta n^{2\alpha-1}}\right)^{\theta/4}\right)^{k/2}\\
&\leqslant dn\sum_{k=\theta}^{\epsilon_1 n}\left(\frac{2^{12}K_0^4d^2k^{2\alpha-1}}{\theta^2 n^{2\alpha-1}}\right)^{\theta k/16}\\
&= o(1).
\end{aligned} \tag{3.4}$$

Here we choose $K_1 \geqslant 2^{10}K_0^2$ so that $2^{12}K_0^4d^2\epsilon_1^{2\alpha-1} < \theta^2/2$.

Explanation of (3.4). Fix $S_1 = \{i_1 < \cdots < i_{k/2}\}$. Now fix the degrees $a_1, \dots, a_{k/2}$ of $i_1, \dots, i_{k/2}$. A crude upper bound on the number of graphs with this set of degrees is

$$\binom{\binom{k/2}{2}}{\ell/2}.$$

We then use the fact that the probability an edge exists between two vertices of degrees d_r, d_s is never more than $\frac{d_r d_s}{dn-\ell}$, given up to $\ell/2-1$ previous pairings. □

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

Now let

$$t_1 = 1 + \lceil \log_3(2/\epsilon_1) \rceil$$

and

$$I_t = \left\{ n - \frac{n}{3^t} + 1, \ldots, n - \frac{n}{3^{t+1}} \right\} \quad \text{for } t = 0, 1, \ldots, t_1.$$

Let G_t be the sub-graph of G_F induced by I_t and for $k \in I_t$ let $\tilde{d}_k$ be the degree of k in G_t.

Let

$$\Delta_t = \frac{K_2 d}{2^{(2\alpha-1)t}},$$

where $K_2 = 3\epsilon^{-1}2^{(2\alpha-1)t_0} + K_2' + K_2''$,

$$t_0 = \lfloor \log_3 1/\epsilon \rfloor,$$

and K_2', K_2'' are defined below.

For $0 \leqslant t \leqslant t_1$ let $B_t = \{k \in I_t : \tilde{d}_k \geqslant \Delta_t\}$, $Z_t = |B_t|$ and $Z = Z_0 + \cdots + Z_{t_1}$.

Lemma 3.2. *$Z \leqslant \epsilon_1 n/2$ w.h.p.*

Proof. If $k \in I_t$, $t \leqslant t_0$, then there are at least $n - |\bigcup_{i=1}^{t} I_i| \geqslant 3\epsilon n$ vertices of degree at least d_k, and thus $d_k \leqslant 3\epsilon^{-1}d$ and so $B_t = \emptyset$, $0 \leqslant t \leqslant t_0$. For $t > t_0$ and $k \in I_t$ we see that $\tilde{d}_k$ is stochastically dominated by the binomial $B\left(d_k, \frac{D_{n/3^t}}{dn}\right)$. Furthermore, since $k \geqslant n - \epsilon n$, we have

$$d_k \leqslant \frac{D_{n/3^{t+1}}}{n/3^{t+1}} \leqslant K_0 3^{(t+1)(1-\alpha)} d \tag{3.5}$$

and

$$\frac{D_{n/3^t}}{dn} \leqslant K_0 3^{-\alpha t}.$$

So, if $K_2' = K_0^2 e^2 3^{1-\alpha}$ then

$$\mathbf{Pr}\left(\tilde{d}_k \geqslant \Delta_t\right) \leqslant \left(\frac{K_0^2 3^{1-\alpha} e}{K_2'(3/2)^{(2\alpha-1)t}} \right)^{\Delta_t} \leqslant e^{-\Delta_t}.$$

(We used $\mathbf{Pr}[B(n,p) \geqslant k] \leqslant \binom{n}{k} p^k \leqslant \left(\frac{enp}{k}\right)^k$.)

Thus,

$$\mathbf{E}(Z) \leqslant n \sum_{t=1}^{t_1} 3^{-t} e^{-\Delta_t} < n e^{-\Delta_{t_1}}.$$

Now,

$$2^{(2\alpha-1)t_1} \leqslant 2^{(2\alpha-1)\left(2+\log_3(2/\epsilon)+\frac{2}{2\alpha-1}\log_3\left(\frac{2K_1 d}{(2\alpha-1)\theta}\right)\right)} \leqslant 2^{2+\log_3(2/\epsilon)+\frac{2}{2\alpha-1}\log_3(2K_1/(2\alpha-1))} d^{\gamma},$$

where

$$\gamma = \frac{1}{2\alpha+1}\frac{\ln 4}{\ln 3} \in \left(\frac{2}{5}, \frac{2}{3}\right). \tag{3.6}$$

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

So, if

$$K_2'' = d^{-1/3}2^{2+\log_3(2/\epsilon)+\frac{2}{2\alpha-1}\log_3(2K_1/(2\alpha-1))}\left(\ln 4 + \frac{2}{2\alpha-1}\ln(K_1 d/(2\alpha-1))\right),$$

then

$$\mathbf{E}(Z) < e^{-d^{2/3-\gamma}(2/(2\alpha-1))\ln(K_1 d/(2\alpha-1))}n/4 \leqslant \epsilon_1 n/4.$$

It is straightforward to show that Z is concentrated about its mean. A random pairing can be obtained by choosing a random permutation $\omega = (w_1, w_2, \ldots, w_{dn})$ of W and pairing w_{2i-1} with w_{2i}, for $1 \leqslant i \leqslant dn/2$. Interchanging two elements in ω can change Z by at most two. Lemma 11 of [11] states:

Let S be a set with $|S| = N$. Let Ω be the set of $N!$ permutations of S. Let ω be chosen uniformly from Ω. Let $Z = Z(\omega)$ be such that $|Z(\omega) - Z(\omega')| \leqslant 1$ when ω' is obtained from ω by interchanging two elements of the permutation.

$$\mathbf{Pr}(|Z - \mathbf{E}(Z)| \geqslant t) \leqslant 2e^{-2t^2/N}. \tag{3.7}$$

Using (3.7) with $N = dn$ and $t = (dn)^{1/2}\ln n$ yields the lemma. □

We now appeal to a result of Alon, Krivelevich and Sudakov [4] to show that we can colour the graphs $\Gamma_t = G_t - B_t$ with few colours. The main result of that paper is as follows.

Theorem 3.3. *There exists an absolute positive constant C_0 such that the following holds. Let G be a graph with maximum degree Δ in which the neighbourhood of any vertex v spans at most Δ^2/f edges. Then the chromatic number of G is at most $C_0\frac{\Delta}{\ln f}$.*

We use this to prove the following lemma.

Lemma 3.4. *There exists $\beta > 0$ such that*

$$\mathbf{Pr}\left(\exists t \in [0, t_1] : \chi(\Gamma_t) \geqslant \frac{C_0\Delta_t}{\beta \ln d}\right) = o(1).$$

Proof. Fix t and $v \in I_t$ and condition on the neighbours of v in G_F for which $v \notin B_t$. In so doing we fix at most Δ pairs of the configuration. With this conditioning, the number of edges ξ_v in the Γ_t-neighbourhood of v is stochastically dominated by the Binomial distribution $B(\Delta_t^2/2, \Delta/(dn - 2\Delta))$. So, if $\beta = 4/3 - 2\gamma > 0$ (γ as in (3.6)) then

$$\mathbf{Pr}\left(\xi_v \geqslant \frac{\Delta_t^2}{d^\beta}\right) \leqslant \binom{\Delta_t^2/2}{\Delta_t^2/d^\beta}\left(\frac{2\Delta}{dn}\right)^{\Delta_t^2/d^\beta} \leqslant \left(\frac{e\Delta}{d^{1-\beta}n}\right)^{\Delta_t^2/d^\beta}.$$

But $\Delta \leqslant n^{1/3}$ and $\Delta_t^2/d^\beta \geqslant 10$ for sufficiently large K_1 and so

$$\mathbf{Pr}\left(\xi_v \geqslant \frac{\Delta_t^2}{d^\beta}\right) = o(1/n).$$

Applying Theorem 3.3 with $\Delta = \Delta_t$ and $f = d^\beta$ yields the lemma. □

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

Lemma 3.5. *If F is chosen uniformly from the set Ω of pairings then w.h.p.*

$$\chi(G_F) \leqslant b_1 \frac{d}{\ln d}.$$

Proof. Let $B_t, G_t, \Gamma_t, t \in [0, t_1]$, be as defined above. It follows from Lemma 3.4 that w.h.p. the number of colours needed to colour $\bigcup_{t=0}^{t_1} \Gamma_t$ is at most

$$\sum_{t=0}^{t_1} \frac{C_0 K_2 d}{2^{(2\alpha-1)t} \beta \ln d} = O\left(\frac{d}{\ln d}\right).$$

Finally note that the number of vertices in $B = \bigcup_{t=0}^{t_1} B_t \cup \{i \geqslant n - \frac{n}{3^{t_1+1}}\}$ is w.h.p. less than ϵn and then, by Lemma 3.1, the graph induced by B is $\theta = d^{2/(2\alpha+1)}$-degenerate and so can be coloured with at most $\theta + 1 = O(d/\ln d)$ colours. □

3.2. From configurations to graphs

It is at this point that we appeal to some results from McKay and Wormald [14]. Where possible, we will use the terminology and notation of that paper. A *loop* of a pairing F is a pair $\{u, v\}$ such that $\phi(u) = \phi(v)$. A *double pair* of F is a pair $\{u_1, v_1\}, \{u_2, v_2\} \in F$ such that $\phi(u_1) = \phi(u_2)$ and $\phi(v_1) = \phi(v_2)$. A *double loop* of F is a pair of pairs $\{u_1, v_1\}, \{u_2, v_2\}$ such that $\phi(u_1) = \phi(v_1) = \phi(u_2) = \phi(v_2)$. A *triple pair* is a triple of pairs $\{u_i, v_i\}$, $i = 1, 2, 3$ such that $\phi(u_1) = \phi(u_2) = \phi(u_3)$ and $\phi(v_1) = \phi(v_2) = \phi(v_3)$.

In the lemmas that follow, we will assume always that condition (b) of Theorem 1.1 holds.

Lemma 3.6. (Lemma 2 of [14]) *The probability that F contains at least one triple pair is $O(\Delta^2 M_2^2 / M_1^3) = o(1)$ and the probability of at least one double loop is $O(\Delta^2 M_2 / M_1^2) = o(1)$.*

Now let l denote the number of loops and r denote the number of double pairs in F.

Lemma 3.7. (Lemma 3 of [14]) *If $\lambda(n) \to \infty$ then w.h.p.*

$$l \leqslant 2\Delta + \lambda \quad \text{and} \quad r \leqslant \Delta^2 + \lambda. \tag{3.8}$$

We define the following two operations on a pairing. If $\phi(u) = i$ then we say that u is in *cell i*.

I *l-switching*. Take pairs $\{p_1, p_6\}, \{p_2, p_3\}, \{p_4, p_5\}$ where $\{p_2, p_3\}$ is a loop, and $p_1, \ldots, p_6$ are in five different cells. Replace these pairs by $\{p_1, p_2\}, \{p_3, p_4\}, \{p_5, p_6\}$. In this operation, none of the pairs created or destroyed is permitted to be part of a double pair. (See Figure 1.)

II *r-switching*. Take pairs $\{p_1, p_5\}, \{p_2, p_6\}, \{p_3, p_7\}, \{p_4, p_8\}$ where $\phi(p_2) = \phi(p_3)$ and $\phi(p_6) = f(p_7)$, but the cells containing $p_1, p_2, p_4, p_5, p_6, p_8$ are all distinct. Replace these pairs by $\{p_1, p_2\}, \{p_3, p_4\}, \{p_5, p_6\}, \{p_7, p_8\}$. In this operation, none of the pairs created or destroyed (other than the pairs $\{p_2, p_6\}, \{p_3, p_7\}$) is permitted to be part of a multiple pair. (See Figure 2.)

A *forward l*-switching is an l-switching as described, and a *backward l*-switching is the reverse operation. We use the same convention for r-switchings. Note that a forward

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press




Figure 1.




Figure 2.

l-switching always reduces the number of loops by one and does not create or destroy double pairs. Similarly, a forward r-switching reduces the number of double pairs by one and neither creates nor destroys loops.

Now let $\mathscr{C}_{l,r}$ denote the set of pairings F with l loops, r double pairs and no triple pairs or double loops.

Lemma 3.8. (Lemma 4 of [14]) *Denote an operation taking an element of $\mathscr{C}_{i,j}$ to an element $\mathscr{C}_{k,l}$ by $\mathscr{C}_{i,j} \to \mathscr{C}_{k,l}$. For each of the following operations, we bound the number, m, of ways of applying the operation to a fixed F.*

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

(1) *Forward l-switching* $\mathscr{C}_{l,r} \to \mathscr{C}_{l-1,r}$*:*

$$2lM_1^2 \geqslant m \geqslant 2lM_1^2\left(1 - O\left(\frac{\Delta^2 + l + r}{M_1}\right)\right).$$

(2) *Backward l-switching* $\mathscr{C}_{l-1,r} \to \mathscr{C}_{l,r}$*:*

$$M_1M_2 \geqslant m \geqslant M_1M_2\left(1 - \frac{\Delta(6(l+2r)+\Delta l)}{M_2} - \frac{2\Delta(\Delta+2)}{M_1}\right).$$

(3) *Forward r-switching* $\mathscr{C}_{0,r} \to \mathscr{C}_{0,r-1}$*:*

$$4rM_1^2 \geqslant m \geqslant 4rM_1^2\left(1 - O\left(\frac{\Delta^2 + r}{M_1}\right)\right).$$

(4) *Backward r-switching* $\mathscr{C}_{0,r-1} \to \mathscr{C}_{0,r}$*:*

$$M_2^2 \geqslant m \geqslant M_2^2\left(1 - \frac{\Delta(16r + 9\Delta + 3 + \Delta^2)}{M_2}\right).$$

Now consider the following algorithm for generating a member of $\mathscr{G}_{n,\mathbf{d}}$:

1 Generate a random pairing F.
2 If there is a double loop or a triple pair, output $\perp$, construed as failure.
3 If the number of loops $l \geqslant 2\Delta + \log n$ or the number of double pairs $r \geqslant \Delta^2 + \log n$, output $\perp$, construed as failure.
4 $F_0 \leftarrow F$.
5 For $i = 1$ to l choose a random forward l-switching on F_{i-1}, creating $F_i \in \mathscr{C}_{l-i,r}$.
6 For $i = l+1$ to $l+r$ choose a random forward r-switching on F_{i-1}, creating $F_i \in \mathscr{C}_{0,r-(i-l)}$.
7 Output $G^* = G_{F_{l+r}} \in \mathscr{G}_{n,\mathbf{d}}$.

For each l, r satisfying (3.8), with $\lambda = \log n$, and $G \in \mathscr{G}_{n,\mathbf{d}}$, there are by Lemma 3.8(2), (4)

$$(M_1M_2)^l M_2^r\left(1 + O\left(\frac{\Delta l(\Delta l + r)}{M_2} + \frac{\Delta^2 l}{M_1} + \frac{\Delta r(\Delta^2 + r)}{M_2}\right)\right)$$

sequences of switchings which yield G. Each of these has probability

$$((2M_1^2)^l l!(4M_1^2)^r r!)^{-1}\left(1 + O\left(\frac{l(\Delta^2 + l + r)}{M_1} + \frac{r(\Delta^2 + r)}{M_1}\right)\right)$$

of being followed by the algorithm, given l, r.

Thus, if condition (b) holds, then w.h.p. the algorithm outputs a graph in $\mathscr{G}_{n,\mathbf{d}}$ and

$$\mathbf{Pr}(G^* = G) = (1 + o(1)) \sum_{l=0}^{2\Delta+\log n} \sum_{r=0}^{\Delta^2+\log n} \frac{M_1^l M_2^{r+l}}{2^l M_1^{2(l+r)} l!\, r!} \mathbf{Pr}(l \text{ loops}, r \text{ double pairs}),$$

and so, for $G_1, G_2 \in \mathscr{G}_{n,\mathbf{d}}$,

$$\mathbf{Pr}(G^* = G_1) = (1 + o(1))\mathbf{Pr}(G^* = G_2).$$

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

Given this, we only have to show that w.h.p. $\chi(G^*) = O(\chi(G_F))$. Thus, let H be the graph induced by the edges that are *added* by the switchings. We will show that w.h.p.

$$\Delta(H) \leqslant 5. \tag{3.9}$$

Since $\chi(G^*) \leqslant \chi(H)\chi(G_F) \leqslant (\Delta(H)+1)\chi(G_F)$, this will complete the proof of our upper bound.

Now every edge at distance $\geqslant 2$ from the loop or double edge can be used as one of the two edges destroyed by the two types of switching. Thus, vertex i has probability

$$O\left(\frac{d_i}{M_1 - \Delta^2}\right) = O\left(\frac{\Delta}{M_1}\right)$$

of having an H-edge created in any switching, regardless of the history of the switchings to this point.

Thus, for some constant $c > 0$, assuming due to Lemma 3.7 that G^* satisfies (3.8),

$$\mathbf{Pr}(\Delta(H) \geqslant 5) \leqslant n\binom{\Delta^2 + 2\Delta + 2\log n}{5}\left(\frac{c\Delta}{M_1}\right)^5 \leqslant nM_1^{5/2-15/4} = o(1),$$

and this completes the proof of the upper bound in Theorem 1.1.

Note that we cannot immediately use the stronger result of [15]. This is because in that paper Lemma 3.8(4) is replaced by an *average* bound.

4. Lower bound

We will work with the configuration model.

We first observe that

$$\mathbf{Pr}(G_F \text{ is simple}) \geqslant e^{-\Delta^2}. \tag{4.1}$$

This follows directly from the formula for the asymptotic number of labelled graphs with a degree sequence in which $\Delta = o(M_1^{1/4})$; see [15].

Let

$$t = b_2\frac{d}{\ln d},$$

where $b_2 > 0$ is a sufficiently small constant. Fix a partition $(V_1, \ldots, V_t)$ of the vertex set of G_F into t parts. We will show:

$$\mathbf{Pr}((V_1, \ldots, V_t) \text{ is a proper colouring of } G_F) \leqslant t^{-2n}.$$

This will be enough to beat both the union bound for the number of t-partitions (which is $\leqslant t^n$) and the inverse probability that G_F is simple (which is $\leqslant e^{\Delta^2}$; see (4.1)).

Observe first that

$$\mathbf{Pr}(V_i \text{ is independent in } G_F) \leqslant \left(1 - \frac{D(V_i)/2 - \Delta}{dn}\right)^{D(V_i)/2}.$$

Explanation. Match the first $D(V_i)/2$ points of V_i in the configuration model. When matching a point, it should be matched either to a point belonging to a vertex outside of V_i, or to a point of the same vertex.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

Now, if in the partition $(V_1,\dots,V_t)$ one of the sets V_i satisfies $D(V_i)\geqslant nd/4$, by the above argument we get (taking b_2 small enough if necessary):

$$\mathbf{Pr}((V_1,\dots,V_t) \text{ is a } t\text{-colouring}) \leqslant \mathbf{Pr}(V_i \text{ is independent}) \leqslant e^{-(1-o(1))nd/64} \ll t^{-2n}.$$

Assume therefore that $|V_i|\leqslant dn/4$ for all $1\leqslant i\leqslant t$. Assume w.l.o.g. that $D(V_1)\leqslant D(V_2)\leqslant \cdots \leqslant D(V_t)$. Choose $1\leqslant k\leqslant t$ so that

$$\frac{dn}{4}\leqslant \sum_{i=1}^{k} D(V_i)\leqslant \frac{dn}{2};$$

such a k exists as $D(V_i)\leqslant nd/4$ for all i.

We will estimate from above the probability that all of the sets V_i, $i=1,\dots,k$, are independent. In order to do this, we will expose the pairings of the points of V_i, each time choosing a set V_i that has at least half of its points unpaired, for as long as possible. Let σ describe the order in which the sets $V_1,\dots,V_k$ are exposed. Observe that

$$\mathbf{Pr}(V_{\sigma(i)} \text{ is independent} \mid V_{\sigma(1)},\dots,V_{\sigma(i-1)} \text{ are independent}) \leqslant \left(1-\frac{D(V_{\sigma(i)})/4-\Delta}{dn}\right)^{D(V_{\sigma(i)})/4} \leqslant e^{-\frac{D(V_{\sigma(i)})^2-4\Delta D(V_{\sigma(i)})}{16dn}}$$

(match the first half of yet unmatched points of $V_{\sigma(i)}$).

Suppose that we have in this way exposed the pairings of sets $V_{\sigma(1)},\dots,V_{\sigma(l)}$. If

$$\sum_{i=1}^{l} D(V_{\sigma(i)})\leqslant \frac{dn}{16},$$

then at this moment the number of paired points does not exceed $dn/8$, and therefore at least one of the unexposed sets V_i, $i\notin\{\sigma(1),\dots,\sigma(l)\}$, has at least half of its points unpaired, and we can proceed. Thus, if by the end of the process we have exposed pairings of sets $V_{\sigma(1)},\dots,V_{\sigma(l^*)}$, it follows that $\sum_{i=1}^{l^*} D(V_{\sigma(i)})\geqslant dn/16$ and therefore

$$\sum_{i=1}^{l^*}(D(V_{\sigma(i)})^2-4\Delta D(V_{\sigma(i)}))\geqslant \frac{(1-o(1))d^2n^2}{256l^*}\geqslant \frac{(1-o(1))d^2n^2}{256t}.$$

Hence

$$\mathbf{Pr}(V_{\sigma(1)},\dots,V_{\sigma(l^*)} \text{ are independent}) \leqslant \exp\left\{-\frac{\sum_{i=1}^{l^*}(D(V_{\sigma(i)})^2-4\Delta D(V_{\sigma(i)}))}{16dn}\right\}\leqslant \exp\left\{-\frac{dn}{4100t}\right\}.$$

Recalling the expression for t and choosing the constant b_2 there to be sufficiently small, we get the desired estimate.

This completes the proof of Theorem 1.1. □

Acknowledgement

We thank Paul Horn for pointing out an error in an earlier draft of the paper.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press

References

[1] Achlioptas, D. and Moore, C. (2004) The chromatic number of random regular graphs. In *Proc. RANDOM 2004*, pp. 219–228.

[2] Achlioptas, D. and Naor, A. (2005) The two possible values of the chromatic number of a random graph. *Ann. of Math.* **62** 1335–1351.

[3] Aiello, W., Chung, F. and Lu, L. (2001) A random graph model for power law graphs. *Experiment. Math.* **10** 53–66.

[4] Alon, N., Krivelevich, M. and Sudakov, B. (1999) Coloring graphs with sparse neighborhoods. *J. Combin. Theory Ser. B* **77** 73–82.

[5] Bender, E. A. and Canfield, E. R. (1978) The asymptotic number of labelled graphs with given degree sequences. *J. Combin. Theory Ser. A* **24** 296–307.

[6] Bollobás, B. (1980) A probabilistic proof of an asymptotic formula for the number of labelled regular graphs. *Europ. J. Combin.* **1** 311–316.

[7] Bollobás, B. (1988) The chromatic number of random graphs. *Combinatorica* **8** 49–55.

[8] Cooper, C. and Frieze, A. M. (2004) The size of the largest strongly connected component of a random digraph with a given degree sequence. *Combin. Probab. Comput.* **13** 319–338.

[9] Cooper, C., Frieze, A. M., Reed, B. A. and Riordan, O. (2002) Random regular graphs of non-constant degree: Independence and chromatic number. *Combin. Probab. Comput.* **11** 323–342.

[10] Frieze, A. M. and Łuczak, T. (1992) On the independence and chromatic numbers of random regular graphs. *J. Combin. Theory Ser. B* **54** 123–132.

[11] Frieze, A. M. and Pittel, B. G.(2004) Perfect matchings in random graphs with prescribed minimal degree. In *Mathematics and Computer Science III* (M. Drmota, P. Flajolet, D. Gardy and B. Gittenberger, eds), Trends in Mathematics, Birkhäuser, Basel, pp. 95–132.

[12] Łuczak, T. (1991) The chromatic number of random graphs. *Combinatorica* **11** 45–54.

[13] McKay, B. D. and Wormald, N. C. (1990) Asymptotic enumeration by degree sequence of graphs of high degree. *Europ. J. Combin.* **11** 565–580.

[14] McKay, B. D. and Wormald, N. C. (1990) Uniform generation of random regular graphs of moderate degree. *J. Algorithms* **11** 52–67.

[15] McKay, B. D. and Wormald, N. C. (1991) Asymptotic enumeration by degree sequence of graphs with degree $o(n^{1/2})$. *Combinatorica* **11** 369–382.

[16] Molloy, M. and Reed, B. A. (1995) A critical point for random graphs with a given degree sequence. *Random Struct. Alg.* **6** 161–180.

[17] Molloy, M. and Reed, B. A. (1998) The size of the largest component of a random graph on a fixed degree sequence. *Combin. Probab. Comput.* **7** 295–306.

[18] Wormald, N. C. (1999) Models of random regular graphs. In *Surveys in Combinatorics: Proc. 1999 British Combinatorial Conference* (J. D. Lamb and D. A. Preece, eds), Vol. 267 of *London Mathematical Society Lecture Note Series*, Cambridge University Press, Cambridge, pp. 239–298.

https://doi.org/10.1017/S0963548306008388 Published online by Cambridge University Press