Ergod. Th. & Dynam. Sys. (2017), **37**, 79–102
doi:10.1017/etds.2015.46
© *Cambridge University Press, 2015*

Multifractal analysis for weak Gibbs measures: from large deviations to irregular sets

THIAGO BOMFIM and PAULO VARANDAS

Departamento de Matemática, Universidade Federal da Bahia, Av. Ademar de Barros s/n, 40170-110 Salvador, Brazil
(e-mail: tbnunes@ufba.br, paulo.varandas@ufba.br)

(*Received* 23 *May* 2014 *and accepted in revised form* 20 *April* 2015)

Abstract. In this article we prove estimates for the topological pressure of the set of points whose Birkhoff time averages are far from the space averages corresponding to the unique equilibrium state that has a weak Gibbs property. In particular, if f has an expanding repeller and ϕ is a Hölder continuous potential, we prove that the topological pressure of the set of points whose accumulation values of Birkhoff averages belong to some interval $I \subset \mathbb{R}$ can be expressed in terms of the topological pressure of the whole system and the large deviations rate function. As a byproduct we deduce that most irregular sets for maps with the specification property have topological pressure strictly smaller than the whole system. Some extensions to a non-uniformly hyperbolic setting, level-2 irregular sets and hyperbolic flows are also given.

1. *Introduction*

Let $f : M \to M$ be a measurable transformation and μ an f-invariant and ergodic probability measure. The celebrated Birkhoff ergodic theorem asserts that for any given $\psi \in L^1(\mu)$ and for μ-almost every $x \in M$

$$\frac{1}{n} S_n \psi(x) := \frac{1}{n} \sum_{i=1}^{n-1} \psi \circ f^i(x) \to \int \psi \, d\mu$$

as n tends to infinity. On the other hand, despite the fact that from the ergodic point of view the set of points where the Birkhoff averages do not converge is negligible, it can be a topologically large set or have full dimension. To illustrate this fact, let us mention that if f is continuous and has the specification property, then the set of points where the Birkhoff averages do not converge is either empty or has total topological pressure with respect to any continuous potential (we refer the reader to e.g. [**35**] for details). The study of the topological pressure or dimension of these multifractal level sets can be traced back to Besicovitch and this topic has had contributions by many authors in recent years

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

(see [**2**, **11**, **12**, **15**, **19**, **20**, **22**, **27**, **30**, **32**, **34**, **35**, **39**] and references therein). Most commonly, given an observable ψ and the decomposition

$$M = \bigcup_{\alpha\in\mathbb{R}} M_\alpha \cup E_\psi$$

where $M_\alpha = \{x \in M : \lim_{n\to\infty} (1/n)S_n\psi(x) = \alpha\}$ and the irregular set E_ψ is the set of points for which the Birkhoff averages do not converge, one is interested in describing each of the previous sets from the topological, dimensional or ergodic point of view. Motivated by the aforementioned results of Thompson [**35**] and the special ergodic theorem proved by Kleptsyn *et al* [**23**], which prove that the Hausdorff dimension of the deviation set for the SRB (Sinai–Ruelle–Bowen) measure is smaller than the dimension of the manifold given a large deviations property, one of our aims in this article is to provide a multifractal description of more general sets.

Let us consider the sets

$$\overline{X}_{\mu,\psi,c} = \left\{x \in M : \limsup_{n\to\infty} \left|\frac{1}{n}\sum_{j=0}^{n-1} \psi(f^j(x)) - \int \psi\, d\mu\right| \geq c\right\}$$

and

$$\underline{X}_{\mu,\psi,c} = \left\{x \in M : \liminf_{n\to\infty} \left|\frac{1}{n}\sum_{j=0}^{n-1} \psi(f^j(x)) - \int \psi\, d\mu\right| \geq c\right\},$$

where μ is an f-invariant and ergodic probability measure, ψ is an observable and $c > 0$, and study them from the topological pressure viewpoint (more general sets will be defined later on). Clearly we have the inclusion $\underline{X}_{\mu,\psi,c} \subset \overline{X}_{\mu,\psi,c}$ for all $c > 0$. In many cases we are interested in studying $f|_\Lambda$ and we will consider the corresponding sets $\underline{X}_{\mu,\psi,c} \cap \Lambda$ and $\overline{X}_{\mu,\psi,c} \cap \Lambda$. When no confusion is possible, for notational simplicity we shall omit the dependence of the sets on ψ, μ and Λ and write simply $\underline{X}_c$ and $\overline{X}_c$. If $J \subset \mathbb{R}$ is an interval, we define $X(J)$ as the set of points x such that the limit $\lim (1/n)S_n\psi(x) \in J$ exists and let $X(c)$ denote the case when one considers the degenerate interval $J = [c, c]$. One motivation is to consider the decomposition of the set of points whose time averages do not converge to the space average by

$$M \setminus \left\{x : \frac{1}{n}\sum_{i=1}^{n-1} \psi \circ f^i(x) \to \int \psi\, d\mu\right\} = \bigcup_{c>0} \overline{X}_c = \bigcup_{c>0} \underline{X}_c$$

and to study the continuity, monotonicity and concavity of the pressure functions

$$c \mapsto P_{\overline{X}_c}(f, \phi) \quad \text{and} \quad c \mapsto P_{\underline{X}_c}(f, \phi).$$

Our first main purpose here is to study the previous functions in a context of dynamical systems admitting equilibrium states that exhibit a weak Gibbs property. Roughly, we prove that the previous topological pressure functions are bounded from above by the topological pressure of the dynamical system with an error term given by an exponential large deviations rate (see Theorem A and Corollaries A and B for precise statements). Furthermore, in a context of uniform hyperbolicity, we prove that $P_{\overline{X}_c}(f, \phi) = P_{\underline{X}_c}(f, \phi)$ and provide upper and lower bounds which allow us to conclude

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

that the pressure function $c \mapsto P_{\overline{X}_c}(f, \phi)$ is differentiable, concave and strictly decreasing in c and varies continuously along continuous parametrized families of dynamical systems (see Theorem B for the precise statement). Similar results for the multifractal analysis of level-2 sets, meaning the analysis of level and irregular sets for Birkhoff averages in the space of probability measures, are also obtained (see Theorem C). Hence, the connection between large deviations and multifractal analysis is revealed to be very fruitful.

Our second main purpose is to provide a finer description of the irregular set E_ψ. Although a dynamical system which admits some hyperbolicity and a unique equilibrium state (that has some weak Gibbs property) with exponential large deviations estimates can be proved to verify that the topological pressure of the sets $\underline{X}_{\mu,\psi,c}$ and $\overline{X}_{\mu,\psi,c}$ is strictly smaller than the topological pressure of the whole system, one cannot expect immediate estimates for the irregular set. In fact, the irregular set E_ψ may not be contained in either of the sets above for some fixed c. This motivates the decompositions $E_\psi = \bigcup_{c>0} \overline{E}_{\mu,\psi,c}$ and $E_\psi = \bigcup_{c>0} \underline{E}_{\mu,\psi,c}$ with

$$\overline{E}_{\mu,\psi,c} = E_\psi \cap \overline{X}_{\mu,\psi,c} \quad \text{and} \quad \underline{E}_{\mu,\psi,c} = E_\psi \cap \underline{X}_{\mu,\psi,c}.$$

In other words, the set $\underline{E}_{\mu,\psi,c}$ consists of those irregular points whose Birkhoff averages $(1/n)\sum_{j=0}^{n-1} \psi(f^j(x))$ remain at a distance larger than c from the time average $\int \psi \, d\mu$ for all large n. Finally, the set $\overline{E}_{\mu,\psi,c}$ consists of irregular points $x \in M$ whose Birkhoff averages $(1/n)\sum_{j=0}^{n-1} \psi(f^j(x))$ have infinitely many values of n for which the Birkhoff averages $(1/n)\sum_{j=0}^{n-1} \psi(f^j(x))$ remain at a distance larger than c. As a direct consequence of our results, even for maps with the specification property, irregular sets $\underline{E}_{\mu,\psi,c}$ and $\overline{E}_{\mu,\psi,c}$ have topological pressure strictly smaller than the topological pressure of the whole system. In fact, we can prove some lower bounds for the topological pressure of these irregular sets and, consequently, study its regularity as a function of the parameter c (cf. Corollary C). An extension to level-2 irregular sets is given in §2 while applications for hyperbolic maps and flows and non-uniformly hyperbolic dynamics are given in §4. In particular, an example from [**16**, **17**] is given that illustrates the case where the pressure function is discontinuous and not strictly decreasing. An extension of the current results to the study of irregular sets for non-additive sequences of observables was carried out in [**4**], while we expect that these results can be also extended to the class of partially hyperbolic diffeomorphisms in [**10**].

This article is organized as follows. In §2 we provide some definitions and state the main results. Section 3 is devoted to proofs of the results. Finally, in §4 we give a large number of applications and examples.

2. *Statement of the main results*

In this section we introduce some necessary notions and state the main results.

2.1. *(Weak) Gibbs measures.* In many cases equilibrium states arise as invariant measures absolutely continuous with respect to probability measures exhibiting some Gibbs property. Let us now describe this wide class of measures. Given $\varepsilon > 0$, $n \geq 1$ and $x \in M$, the *(n, ε)-dynamical ball* $B(x, n, \varepsilon)$ is the set of points $y \in M$ so that $d(f^j(x), f^j(y)) < \varepsilon$ for all $0 \leq j \leq n-1$.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

Definition 2.1. Given a potential ϕ we call a probability ν a *weak Gibbs measure* with respect to ϕ on $\Lambda \subset M$ if there exists $\varepsilon_0 > 0$ such that for every $0 < \varepsilon < \varepsilon_0$ there exists $K(\varepsilon) > 0$ and for ν-almost every x there exists a sequence $n_k(x) \to \infty$ such that

$$K(\varepsilon)^{-1} \leq \frac{\nu(B(x, n_k(x), \varepsilon))}{e^{-n_k(x)P + S_{n_k(x)}\phi(x)}} \leq K(\varepsilon),$$

where $S_n\phi = \sum_{j=0}^{n-1} \phi \circ f^j$ denotes the usual Birkhoff sum. If this condition holds for all positive integers n (independently of x) we call ν a *Gibbs measure* with respect to ϕ.

In this notion of weak Gibbs, one does not require the sequence of times to have positive density at infinity in the set of integers. Naturally, in applications, the most interesting case is when the value P in the previous expression coincides with the topological pressure $P_{\text{top}}(f, \phi)$. Such measures arise naturally in the context of (non-uniform) hyperbolic dynamics. Given a basic set Ω for a diffeomorphism f Axiom A (or Ω repeller to f), it is known that every potential ϕ satisfying

$$\text{there exists } A, \delta > 0 : \sup_{n \in \mathbb{N}} \gamma_n(\phi, \delta) \leq A, \tag{2.1}$$

where $\gamma_n(\phi, \delta) := \sup\{|S_n\phi(y) - S_n\phi(z)| : y, z \in B(x, n, \delta)\}$, admits a unique equilibrium state μ_ϕ and is a Gibbs measure. This condition, introduced by Bowen [**5**] to prove uniqueness of equilibrium states for expansive maps with the specification property, is known as the *Bowen condition*.

2.2. *Statement of the main results.* This section is devoted to the statement of the main results. For this purpose we shall introduce some definitions and notation. The first result provides a topological counterpart to the special ergodic theorem. In what follows, given a continuous function $\psi : M \to \mathbb{R}$, a probability measure μ and a closed set $I \subset \mathbb{R}$, we define

$$\overline{X}_I = \left\{ x \in M : \limsup_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n-1} \psi(f^j(x)) \in I \right\}$$

and, analogously,

$$\underline{X}_I = \left\{ x \in M : \liminf_{n \to \infty} \frac{1}{n} \sum_{j=0}^{n-1} \psi(f^j(x)) \in I \right\}.$$

Moreover, given $\delta > 0$, we denote by I_δ the δ-neighborhood of the set I. Finally, given a probability measure ν, let us define the large deviations upper bound:

$$L_{I,\nu} := -\limsup_{n \to +\infty} \frac{1}{n} \log \nu\left(\left\{ x \in \Lambda : \frac{1}{n} S_n\psi(x) \in I \right\} \right). \tag{2.2}$$

We are now in a position to state our first main result.

THEOREM A. *Let M be a compact metric space, $f : M \to M$ a continuous map, $\phi : M \to \mathbb{R}$ a continuous potential, ν a (not necessarily invariant) Gibbs measure on M and $\mu_\phi \ll \nu$ the unique equilibrium state of f with respect to ϕ. Then, for any continuous $\psi : M \to \mathbb{R}$, any closed interval $I \subset \mathbb{R}$ and any small δ,*

$$P_{\underline{X}_I}(f, \phi) \leq P_{\overline{X}_I}(f, \phi) \leq P_{\text{top}}(f, \phi) - L_{I_\delta,\nu} \leq P_{\text{top}}(f, \phi).$$

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

In fact, it follows from [**36**, Theorem 2.1] that, since ν is a (strong) Gibbs measure, if $\int \psi \, d\mu_\phi \notin I_\delta$ then the large deviations property $L_{I_\delta} > 0$ holds and, consequently, the topological pressure of the sets $\underline{X}_I$ and $\overline{X}_I$ is strictly smaller than $P_{\text{top}}(f, \phi)$. When no confusion is possible we shall omit the dependence of $L_{I,\nu}$ on ν. Our result is applicable to the case of topological repellers.

Remark 2.2. We notice that Theorem A also holds for bilateral subshifts of finite type and *locally Hölder* observables. In fact, given such an observable g there exists ψ that is constant along locally stable leaves (depending only on future coordinates of the shift) and such that $g = \psi + u \circ f - u$ for some continuous u (see [**6**]). Thus, $|(1/n)S_n g(x) - (1/n)S_n \psi(x)| \leq (2\|u\|_0)/n$ tends to zero (uniformly) as n approaches infinity and, consequently, $X_I(g) = X_I(\psi)$, $\overline{X}_I(g) = \overline{X}_I(\psi)$ and $\underline{X}_I(g) = \underline{X}_I(\psi)$ for all intervals $I \subset \mathbb{R}$. Hence, using the Gibbs property and replacing dynamic balls by cylinders associated with the Markov partition, the conclusions of Theorem A still hold.

Definition 2.3. Given a compact metric space (M, d) and a continuous open map $f : M \to M$ we say that an f-invariant set $\Lambda \subset M$ is a *repeller* for f if there are $C, \lambda, \varepsilon > 0$ so that $d(f^n(x), f^n(y)) \geq Ce^{\lambda n} d(x, y)$ for all $y \in B(x, n, \varepsilon)$ and $n \geq 1$.

It is clear that this holds for smooth expanding maps. Recall that an observable $\psi : M \to \mathbb{R}$ is *cohomologous to a constant* if there exists a constant c and a measurable function u such that $\psi = u \circ f - u + c$.

Definition 2.4. Given an observable $\psi : M \to \mathbb{R}$ and $t \in \mathbb{R}$, the *free energy* $\mathcal{E}_{f,\phi,\psi}$ is

$$\mathcal{E}_{f,\phi,\psi}(t) = \limsup_{n\to\infty} \frac{1}{n} \log \int e^{tS_n\psi} \, d\mu_{f,\phi}.$$

In many cases, e.g. when the transfer operator associated with the potential ϕ has a spectral gap property, the expression on the right-hand side converges to

$$\mathcal{E}_{f,\phi,\psi}(t) := \lim_{n\to\infty} \frac{1}{n} \log \int e^{tS_n\psi} \, d\mu_{f,\phi} = P_{\text{top}}(f, \phi + t\psi) - P_{\text{top}}(f, \phi).$$

If this is the case and the topological pressure is smooth, then $t \mapsto \mathcal{E}_{f,\phi,\psi}(t)$ is affine if ψ is cohomologous to a constant, and otherwise $t \mapsto \mathcal{E}_{f,\phi,\psi}(t)$ is strictly convex in some interval $J = [t_-, t_+]$ and one can associate with it the *'local' Legendre transform* $I_{f,\phi,\psi}$ given by

$$I_{f,\phi,\psi}(s) = \sup_{t\in J}\{s\, t - \mathcal{E}_{f,\phi,\psi}(t)\}$$

and well defined in the interval $[\mathcal{E}'_{f,\phi,\psi}(t_-), \mathcal{E}'_{f,\phi,\psi}(t_+)]$. The interval J may depend on f, ϕ and ψ and on the fact that $I_{f,\phi,\psi}(s)$ can often be proved to be a (local) level-1 large deviations *rate function* (see e.g. [**3**, **31**, **38**]): for all $[a, b] \subset J$

$$\limsup_{n\to\infty} \frac{1}{n} \log \nu_{f,\phi}\left(x \in M : \frac{1}{n} S_n\psi(x) \in [a, b]\right) \leq - \inf_{s\in[a,b]} I_{f,\phi,\psi}(s) \tag{2.3}$$

and

$$\liminf_{n\to\infty} \frac{1}{n} \log \nu_{f,\phi}\left(x \in M : \frac{1}{n} S_n\psi(x) \in (a, b)\right) \geq - \inf_{s\in(a,b)} I_{f,\phi,\psi}(s). \tag{2.4}$$

As a byproduct of our previous result, the large deviations property and the fact that equilibrium states associated to Hölder continuous potentials satisfy the Gibbs property, we deduce the following corollary.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

COROLLARY A. *Let $f : M \to M$ be a continuous map, $\Lambda \subset M$ be a transitive repeller, $\phi : M \to \mathbb{R}$ be a Hölder continuous potential and $\mu = \mu_{f,\phi}$ be the unique equilibrium state for $f|_\Lambda$ with respect to ϕ. Then, for any continuous observable $\psi : M \to \mathbb{R}$ and $c > 0$,*

$$P_{\underline{X}_c}(f, \phi) \leq P_{\overline{X}_c}(f, \phi) \leq P_{\text{top}}(f, \phi) - L_{c-\delta} < P_{\text{top}}(f, \phi)$$

for every small δ, where $L_c := L_{I_c}$ is defined as in (2.2) with respect to $I_c = (-\infty, \int \psi \, d\mu_\phi - c] \cup [\int \psi \, d\mu_\phi + c, +\infty)$.

Since, in the previous results, the topological pressure is strictly smaller than the topological pressure $P_{\text{top}}(f, \phi)$, this has particularly interesting applications in connection with the specification property. Recall that a system satisfies the *specification property* if for any $\varepsilon > 0$ there exists an integer $N = N(\varepsilon) \geq 1$ such that the following holds: for every $k \geq 1$, any points $x_1, \ldots, x_k$ and sequences of positive integers $n_1, \ldots, n_k$ and $p_1, \ldots, p_k$ with $p_i \geq N(\varepsilon)$, there exists a point x in M such that

$$d(f^j(x), f^j(x_1)) \leq \varepsilon \quad \text{for all } 0 \leq j \leq n_1$$

and

$$d(f^{j+n_1+p_1+\cdots+n_{i-1}+p_{i-1}}(x), f^j(x_i)) \leq \varepsilon$$

for every $2 \leq i \leq k$ and $0 \leq j \leq n_i$. We also obtain the following result.

COROLLARY B. *Let $f : M \to M$ be a continuous map admitting a transitive repeller $\Lambda \subset M$ and $\psi : M \to \mathbb{R}$ be such that the set of irregular points satisfies $E_\psi \neq \emptyset$. Then $P_{\text{top}}(f, \phi) = P_{E_\psi}(f, \phi) > P_{\overline{X}_c}(f, \phi)$ for every $c > 0$.*

In fact, it follows from [**35**] that a dynamical system with the specification property is such that irregular sets are either empty or have full topological pressure with respect to any continuous potential. Since the dynamical systems restricted to the transitive repeller satisfies the specification property, the first equality follows from [**35**]. In particular, using

$$E_\psi = \bigcup_{n \geq 1} [E_\psi \cap \overline{X}_{1/n}]$$

and also $P_{E_\psi}(f, \phi) = \sup_{n \geq 1} P_{E_\psi \cap \overline{X}_{1/n}}(f, \phi)$, the previous corollary roughly means that despite the set of irregular points having full topological pressure, the ones that give a larger contribution to the topological pressure are those with time averages which are infinitely often very close to the mean.

One could wonder if there could exist a strict inequality $P_{\underline{X}_c}(f, \phi) < P_{\overline{X}_c}(f, \phi)$ and what would be the regularity of the topological pressure of those subsets. The next theorem provides an answer to these questions under the assumption of uniform expansion.

THEOREM B. *Let $f : M \to M$ be a continuous map admitting a mixing repeller $\Lambda \subset M$, $\phi : M \to \mathbb{R}$ be a continuous potential so that μ_ϕ is the unique equilibrium state for f with respect to ϕ and $\mu_\phi \ll \nu$ where ν is a Gibbs measure. If ϕ, ψ satisfy the Bowen condition, ψ is not cohomologous to a constant and $\int \psi \, d\mu_{f,\phi} = 0$ then*

$$P_{\overline{X}_c}(f, \phi) \leq P_{\text{top}}(f, \phi) - \min\{I_{f,\phi,\psi}(-c), I_{f,\phi,\psi}(c)\}$$

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press




FIGURE 1. Continuity, monotonicity and concavity of the pressure function.

where $I_{f,\phi,\psi}$ is the large deviations rate function. If $0 \notin [c_1, c_2]$ and $c = \min\{|c_1|, |c_2|\}$ then either $\overline{X}_c = \emptyset$ or

$$P_{\overline{X}_c}(f,\phi) = P_{\underline{X}_c}(f,\phi) = P_{X(c_*)}(f,\phi) = P_{\mathrm{top}}(f,\phi) - I_{f,\phi,\psi}(c_*)$$

where

$$c_* = \begin{cases} c & \text{if } I_{f,\phi,\psi}(c) < I_{f,\phi,\psi}(-c), \\ -c & \text{otherwise.} \end{cases} \tag{2.5}$$

In particular, $\mathbb{R}_0^+ \ni c \mapsto P_{\overline{X}_c}(f,\phi)$ is differentiable, concave and strictly decreasing. Furthermore, the right-hand-side expression varies continuously with c and also varies continuously with ϕ, ψ in the C^α-topology. Moreover, if V is a compact metric space and $V \ni v \mapsto (f_v)_v$ is a continuous (in the C^1-topology) family of expanding maps on M then $v \mapsto P_{\overline{X}_c}(f_v,\phi)$ is also a continuous function.

We refer the reader to Figure 1 for an illustration of the properties of these pressure functions. Under the previous assumptions we can provide a more detailed description of the irregular sets $\overline{E}_c$ as follows.

COROLLARY C. *Let $f : M \to M$ be a continuous map admitting a mixing repeller $\Lambda \subset M$, $\phi : M \to \mathbb{R}$ be a continuous potential so that μ_ϕ is the unique equilibrium state for f with respect to ϕ and $\mu_\phi \ll \nu$ where ν is a Gibbs measure. Set $\overline{E}_c = \overline{X}_c \cap E_\psi$, the irregular set contained in $\overline{X}_c$. If $E_\psi \neq \emptyset$ then for every $c > 0$:*

(1) $P_{\mathrm{top}}(f,\phi) = P_{E_\psi}(f,\phi) > P_{\overline{X}_c}(f,\phi) \geq P_{\overline{E}_c}(f,\phi)$*; and*
(2) *if $\overline{E}_c \neq \emptyset$ then $P_{\overline{X}_c}(f,\phi) = P_{\overline{E}_c}(f,\phi)$ and $c \mapsto P_{\overline{E}_c}(f,\phi)$ is differentiable, concave and strictly decreasing.*

Actually, in this setting we can also provide estimates for irregular sets corresponding to empirical measures $\delta_{x,n} := (1/n)\sum_{j=0}^{n-1} \delta_{f^j(x)}$. Let $\mathcal{M}_1$ denote the set of probability measures on M and let d be any metric compatible with the weak* topology (e.g. $d(\mu,\nu) = \sum_{k\geq 1} 1/(2^k \|g_k\|_0)|\int g_k\, d\mu - \int g_k\, d\nu|$ for some countable and dense subset

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

$(g_k)_k$ of continuous observables). We say a *level-2 large deviations principle* holds for ν if there is a lower semicontinuous function $Q : \mathcal{M}_1 \to [0, +\infty]$ so that

$$\limsup_{n\to\infty} \frac{1}{n} \log \nu_{f,\phi}(x \in M : \delta_{x,n} \in U) \leq - \inf_{\eta\in U} Q(\eta)$$

for every closed set $U \subset \mathcal{M}_1$ and

$$\liminf_{n\to\infty} \frac{1}{n} \log \nu_{f,\phi}(x \in M : \delta_{x,n} \in V) \geq - \inf_{\eta\in V} Q(\eta)$$

for every open set $V \subset \mathcal{M}_1$. Level-2 large deviations principles in dynamical systems have been obtained, e.g. in [**13**, **14**]. Consider

$$\overline{Y}_{\mu,c} = \left\{x \in M : \limsup_{n\to\infty} d(\delta_{x,n}, \mu) \geq c\right\}, \quad \underline{Y}_{\mu,c} = \left\{x \in M : \liminf_{n\to\infty} d(\delta_{x,n}, \mu) \geq c\right\}$$

and, for $C \subset \mathcal{M}_1$, define $Y(C) := \{x \in M : \lim_{n\to+\infty} \delta_{x,n} \in C\}$.

Part of the strategy can be used to estimate the topological pressure of points with specified behavior of the empirical measures for dynamical systems that have the g-almost product structure and uniform separation property. These notions, introduced by Pfister and Sullivan [**28**], are strictly weaker than the specification property and the positive expansive property, respectively. In fact, the uniform separation property is true even for asymptotically entropy-expansive maps. Let us recall these notions.

Definition 2.5. Let M be a compact metric space and $f : M \to M$ be continuous. A non-decreasing unbounded map $g : \mathbb{N} \to \mathbb{N}$ is a *blow-up function* if $g(n) < n$ for all n and $\lim_{n\to+\infty} g(n)/n = 0$.

For any subset of integers $\Lambda \subset [0, N]$, we will use the family of distances in the metric space X given by $d_\Lambda(x, y) = \max\{d(f^i x, f^i y) : i \in \Lambda\}$ and consider the balls $B_\Lambda(x, \varepsilon) = \{y \in X : d_\Lambda(x, y) < \varepsilon\}$. Given a blow-up function g, $\varepsilon > 0$ and $n \geq 1$, the *g-mistake dynamical ball* $B_n(g; x, \varepsilon)$ of radius ε and length n associated with g is defined by

$$B_n(g; x, \varepsilon) = \{y \in X \mid y \in B_\Lambda(x, \varepsilon) \text{ for some } \Lambda \in I(g; n, \varepsilon)\} = \bigcup_{\Lambda\in I(g;n,\varepsilon)} B_\Lambda(x, \varepsilon)$$

where $I(g; n, \varepsilon) = \{\Lambda \subset [0, n-1] \cap \mathbb{N} \mid \#\Lambda \geq n - g(n)\}$. We are now in a position to define the g-almost product property.

Definition 2.6. Let g be a blow-up function. The continuous map $f : M \to M$ has the *g-almost product property* if there exists a non-increasing function $m : \mathbb{R}^+ \to \mathbb{N}$ such that for any $k \in \mathbb{N}$, any points $x_1, x_2, \ldots, x_k$, any positive $\varepsilon_1, \ldots, \varepsilon_k$ and any integers $n_i \geq m(\varepsilon_1)$ for $i = 1 \cdots k$, it holds that $\bigcap_{j=1}^k f^{-M_{j-1}} B_{n_j}(g; x_j, \varepsilon_j) \neq \emptyset$ where $M_0 = 0$ and $M_i = n_1 + n_2 + \cdots + n_i$, $i = 1, 2, \ldots, k-1$.

Given $\delta, \varepsilon > 0$ and $n \geq 1$ we say that two points $x, y \in X$ are (δ, n, ε)-*separated* if $\#\{0 \leq j \leq n-1 : d(f^j(x), f^j(y)) > \varepsilon\} \geq \delta n$. In addition, a set $E \subset X$ is (δ, n, ε)-separated if all pairs of distinct points in E are (δ, n, ε)-separated. This means that the moments at which the two pieces of orbit are ε-separated form a δ-proportion.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

Definition 2.7. A continuous map $f : M \to M$ has the *uniform separation property* if for any η there exists $\delta > 0$ and $\varepsilon > 0$ so that for any ergodic probability measure μ and any neighborhood F of μ in the space of all probability measures $\mathcal{M}_1$ there exists $n_{F,\mu,\eta} \geq 1$ such that

$$N(F; \delta, n, \varepsilon) \geq \exp[n(h_\mu(f) - \eta)]$$

for all $n \geq n_{F,\mu,\eta}$, where $N(F; \delta, n, \varepsilon)$ is the maximal cardinality of a (δ, n, ε)-separated subset of the set $\{x \in M : \delta_{x,n} \in F\}$.

Taking these notions in account, we also obtain the following result.

THEOREM C. *Let $f : M \to M$ and $\phi : M \to \mathbb{R}$ be continuous, ν be a (not necessarily invariant) Gibbs measure and assume $\mu = \mu_{f,\phi} \ll \nu$ is the unique equilibrium state for f with respect to ϕ. Assume the metric d on $\mathcal{M}_1$ has the following properties:*

(i) $d(\eta_1 + \eta, \eta_2 + \eta) = d(\eta_1, \eta_2)$ *for all* $\eta_1, \eta_2, \eta \in \mathcal{M}_1$*; and*

(ii) $d(t\eta_1, t\eta_2) = td(\eta_1, \eta_2)$ *for all* $\eta_1, \eta_2 \in \mathcal{M}_1$ *and* $t > 0$.

If a level-2 large deviations principle holds for ν then, for every $c > 0$,

$$P_{\overline{Y}_{\mu,c}}(f, \phi) \leq P_{\text{top}}(f, \phi) - \inf_{d(\eta,\mu)\geq c} Q(\eta) \leq P_{\text{top}}(f, \phi).$$

In addition, if f satisfies the almost product and uniform separation properties and $0 < c_1 < c_2$ then either $\overline{Y}_{\mu,c_1} = \emptyset$ or

$$\begin{aligned} P_{\overline{Y}_{\mu,c_1}}(f, \phi) &= P_{\underline{Y}_{\mu,c_1}}(f, \phi) = P_{Y(\partial B(\mu,c_1))}(f, \phi) = P_{Y(B(\mu,c_1))}(f, \phi) \\ &= P_{Y(\overline{B(\mu,c_1,c_2)})}(f, \phi) = P_{Y(B(\mu,c_1,c_2))}(f, \phi) = P_{\text{top}}(f, \phi) - \inf_{d(\eta,\mu)=c_1} Q(\eta), \end{aligned}$$

where $B(\mu, c_1)$ denotes the ball of radius c_1 around μ and $B(\mu, c_1, c_2)$ denotes the annulus $\{\eta \in \mathcal{M}(X) : c_1 < d(\eta, \mu) < c_2\}$.

Further information can be extracted if one knows the behavior of the rate function Q, in which case one can prove that the topological pressure of the level sets is strictly smaller than the topological pressure $P_{\text{top}}(f, \phi)$. This is the case for repellers, as we now detail.

COROLLARY D. *Suppose $f : M \to M$ is a continuous map admitting a transitive repeller $\Lambda \subset M$, $\phi : M \to \mathbb{R}$ is a continuous potential and there exists a unique equilibrium state μ_ϕ for f with respect to ϕ which is a Gibbs measure under Λ. Then, for all $0 < c_1 < c_2$, either $\overline{Y}_{\mu,c} = \emptyset$ or*

$$P_{\overline{Y}_{\mu,c_1}}(f, \phi) = P_{\underline{Y}_{\mu,c_1}}(f, \phi) = P_{\text{top}}(f, \phi) - \inf_{d(\eta,\mu)=c_1} Q(\eta) < P_{\text{top}}(f, \phi)$$

where $Q(\eta) = P_{top}(f, \phi) - h_\eta(f) + \int \psi \, d\eta$.

The previous result implies that the set of irregular points whose range of values of Birkhoff averages are far from the corresponding value associated with the equilibrium state has topological pressure smaller than $P_{\text{top}}(f, \phi)$. In particular, this shows that in order to build an irregular set of points with large topological pressure, one needs to use some specification property and points whose empirical measures are arbitrarily close to the equilibrium state. In some sense, this means the classical construction of irregular sets with large topological pressure is optimal. Our next results apply for weak Gibbs measures.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

THEOREM D. *Let M be a compact metric space, $f : M \to M$ be a continuous map, $\phi : M \to \mathbb{R}$ be a continuous potential, ν be a (not necessarily invariant) weak Gibbs measure and $\mu_\phi \ll \nu$ be the unique equilibrium state of f with respect to ϕ. For any continuous $\psi : M \to \mathbb{R}$ and closed interval $I \subset \mathbb{R}$, it holds that*

$$P_{\underline{X}_I}(f, \phi) \leq P_{\text{top}}(f, \phi) - L_{I_\delta}$$

for every small δ. If, in addition, $L_{I_\delta} < 0$, then $P_{\underline{X}_I}(f, \phi) < P_{\text{top}}(f, \phi)$.

Estimates for L_{I_δ} will depend on the weak Gibbs property and some can be found in [**36**]. Actually, we can prove a version of the previous results in the non-uniformly expanding setting. Given $\sigma, \delta > 0$, we define $H = H(\sigma, \delta)$ as the set of points in Λ with infinitely many (σ, δ)-hyperbolic times (see e.g. [**37**] for a precise definition). We will say that an f-invariant probability measure μ is *expanding* if $\mu(H(\sigma, \delta)) = 1$ for some positive constants σ, δ. Moreover, given ψ continuous, we will say that we have an *exponential large deviations upper bound* if

$$\limsup_{n \to +\infty} \frac{1}{n} \log \mu_\Phi \left(\left\{ x \in M : \left| \frac{1}{n} S_n \psi(x) - \int \psi \, d\mu_\phi \right| \geq c \right\} \right) < 0 \tag{2.6}$$

for all $c > 0$. A direct consequence of the previous abstract result is as follows.

COROLLARY E. *Let $f : M \to M$ be a $C^{1+\alpha}$-smooth map on a compact manifold M and $\phi : M \to \mathbb{R}$ be a continuous potential so that μ_ϕ is the unique equilibrium state for f with respect to ϕ. If μ_ϕ is an expanding measure and $J_{\mu_\phi} f$ is Hölder continuous and has exponential large deviations upper bound, then for any continuous $\psi : M \to \mathbb{R}$ and $c > 0$,*

$$P_{\underline{X}_c}(f, \phi) \leq P_{\text{top}}(f, \phi) - L_{c-\delta} < P_{\text{top}}(f, \phi)$$

for every small $\delta > 0$.

The key ingredient used in the proof of Corollary E is that hyperbolic times are instants at which the Gibbs property holds provided the Jacobian of the measure has enough regularity to deduce bounded distortion. One should also point out that if a local large deviations principle holds as in equations (2.3) and (2.4) (e.g. [**3**, **31**, **38**]) then it is not hard to see that an upper bound for $P_{\underline{X}_c}(f, \phi)$ can be taken as

$$P_{\text{top}}(f, \phi) - \min \left\{ I_{f,\phi,\psi} \left(\int \psi \, d\mu_\phi + c \right), \; I_{f,\phi,\psi} \left(\int \psi \, d\mu_\phi - c \right) \right\}.$$

There are examples where the right-hand-side term above can also be shown to vary continuously with the data even in the non-uniformly expanding context (see e.g. [**3**]). Since we only estimated the topological pressure of the sets $\underline{X}_c$ in the non-uniformly expanding context, one question that arises naturally is the following: Are there examples of transitive non-uniformly expanding maps under the conditions of the previous theorem where $P_{\overline{X}_c}(f, \phi)$ differs from $P_{\underline{X}_c}(f, \phi)$ and coincides with the topological pressure $P_{\text{top}}(f, \phi)$?

We include some examples in §4 where we give partial answers to this question by proving that these sets may have different upper Carathéodory capacities.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

3. *Proofs of the main results*

3.1. *Proof of Theorem A.* Our purpose is to estimate $P_{\underline{X}_I}(f,\phi)$ and $P_{\overline{X}_I}(f,\phi)$. Consider the sets $X_{I,n}=\{x\in M:(1/n)S_n\psi(x)\in I\}$. Let us first prove a preliminary lemma.

LEMMA 3.1. *Let $I\subset\mathbb{R}$ be a closed set. For any $\delta>0$ there exists $\varepsilon_\delta>0$ and $N=N_\delta\in\mathbb{N}$ so that $B(x,n,\varepsilon)\subset X_{I_\delta,n}$ for all $0<\varepsilon<\varepsilon_\delta$, $n\geq N$ and $x\in X_{I,n}$.*

Proof. Let $\delta>0$ be given. Since ψ is uniformly continuous then there is $\varepsilon=\varepsilon_\delta>0$ and a large $N=N_\delta\in\mathbb{N}$ so that $\gamma_n(\psi,\varepsilon)\leq\delta n$ for all $0<\varepsilon<\varepsilon_\delta$ and $n\geq N$. So, if $n\geq N$, $x\in X_{I,n}$, $y\in B(x,n,\varepsilon)$ and $0<\varepsilon<\varepsilon_\delta$, then

$$\frac{S_n\psi(x)}{n}-\frac{\gamma_n(\psi,\varepsilon)}{n}\leq\frac{S_n\psi(y)}{n}\leq\frac{S_n\psi(x)}{n}+\frac{\gamma_n(\psi,\varepsilon)}{n}$$

and, consequently,

$$\frac{S_n\psi(x)}{n}-\delta\leq\frac{S_n\psi(y)}{n}\leq\frac{S_n\psi(x)}{n}+\delta,$$

meaning that $y\in X_{I_\delta,n}$. This finishes the proof of the lemma. □

Proof of Theorem A. Let $I\subset\mathbb{R}$ be a closed interval and assume $\overline{X}_I$ is non-empty. Let $\delta>0$ be fixed and consider L_{I_δ} as defined in equation (2.2). For any positive integer n consider the set $\mathcal{I}_n\subset M\times\mathbb{N}$ of pairs (x,n) with $x\in M$. Recalling the notion of topological pressure for invariant sets introduced by Pesin and Pitskel (see e.g. [**26**]), in order to prove that $P_{\overline{X}_I}(f,\phi)\leq P_{\text{top}}(f,\phi)-L_{I_\delta}$ it is enough to prove that for all $\alpha>P_{\text{top}}(f,\phi)-L_{I_\delta}$, every $\varepsilon>0$ and $N\in\mathbb{N}$ there exists a subset $\hat{\mathcal{G}}_N\subset\bigcup_{n\geq N}\mathcal{I}_n$ so that

$$\overline{X}_I\subset\bigcup_{(x,n)\in\hat{\mathcal{G}}_N}B(x,n,\varepsilon)\quad\text{and}\quad\sum_{(x,n)\in\hat{\mathcal{G}}_N}e^{-\alpha n+\phi_n(x)}\leq a(\varepsilon)<\infty$$

independently of N.

Let $\alpha>P_{\text{top}}(f,\phi)-L_{I_\delta}$ and $0<\varepsilon<\varepsilon_\delta$ be fixed. Notice that if $x\in\overline{X}_I$ then there exists a sequence of positive integers $(m_j(x))_{j\in\mathbb{N}}$ converging to infinity such that $x\in X_{I_\delta,m_j(x)}$ for all $j\in\mathbb{N}$. Thus, $\overline{X}_I\subset\bigcap_{\ell\geq1}\bigcup_{j\geq\ell}X_{I_\delta,j}$. Given $N\geq1$ and $x\in\overline{X}_I$, pick $m(x)\geq N$ in such a way that $x\in X_{I_{\delta/2},m(x)}$ and consider $\mathcal{G}_N:=\{(x,m(x)):x\in\overline{X}_I\}$. Now, let $\hat{\mathcal{G}}_N\subset\mathcal{G}_N$ be a maximal set with a property of separation, namely, that if (x,l) and (y,l) belong to $\hat{\mathcal{G}}_N$ then $B(x,l,\epsilon/2)\cap B(x,l,\epsilon/2)=\emptyset$. So, for $0<\varepsilon<\delta$ given by Lemma 3.1, using the Gibbs property for ν we deduce that

$$\begin{aligned}\sum_{(x,m(x))\in\hat{\mathcal{G}}_N}e^{-\alpha m(x)+S_{m(x)}\phi(x)}&=\sum_{(x,m(x))\in\hat{\mathcal{G}}_N}e^{(P-\alpha)m(x)}e^{-Pm(x)+S_{m(x)}\phi(x)}\\&\leq\sum_{(x,m(x))\in\hat{\mathcal{G}}_N}e^{(P-\alpha)m(x)}K(\varepsilon)\nu(B(x,m(x),\varepsilon)).\end{aligned}$$

Now, we write $\hat{\mathcal{G}}_N=\cup_{\ell\geq1}\hat{\mathcal{G}}_{\ell,N}$ with the level sets $\hat{\mathcal{G}}_{\ell,N}:=\{(x,\ell)\in\hat{\mathcal{G}}_N\}$ and pick $\zeta>0$ small such that $\alpha>P_{\text{top}}(f,\Phi)-L_{I_\delta}+\zeta$ and $\mu(\{x\in\Lambda:(1/n)S_n\psi(x)\in I_\delta\})\leq$

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

$e^{-(L_{I_\delta}-\zeta)n}$ for all $n \geq N$ large. By Lemma 3.1, each dynamical ball $B(x, \ell, \varepsilon)$ is contained in $X_{I_\delta,\ell}$. Therefore, using that $\nu(B(x, m(x), \varepsilon)) \leq K(\varepsilon)K(\varepsilon/2)\nu(B(x, m(x), \varepsilon/2))$,

$$\begin{aligned}
\sum_{(x,m(x))\in\hat{\mathcal{G}}_N} e^{-\alpha m(x)+S_{m(x)}\phi(x)} &\leq K(\varepsilon) \sum_{(x,m(x))\in\hat{\mathcal{G}}_N} e^{(P-\alpha)(m(x))}\nu(B(x, m(x), \varepsilon)) \\
&= K(\varepsilon) \sum_{\ell\geq N} e^{(P-\alpha)\ell} \sum_{x\in\hat{\mathcal{G}}_{N,\ell}} \nu(B(x, \ell, \varepsilon)) \\
&\leq K(\varepsilon)K\left(\frac{\varepsilon}{2}\right) \sum_{\ell\geq N} e^{(P-\alpha)\ell} \sum_{x\in\hat{\mathcal{G}}_{N,\ell}} \nu(B(x, \ell, \varepsilon/2)) \\
&\leq K(\varepsilon)K\left(\frac{\varepsilon}{2}\right) \sum_{\ell\geq N} e^{(P-\alpha)\ell}\nu(X_{I_\delta,\ell}) \\
&\leq K(\varepsilon)K\left(\frac{\varepsilon}{2}\right) \sum_{\ell\geq N} e^{(P-\alpha-L_{c-\delta}+\zeta)\ell},
\end{aligned}$$

which is finite and independent by the choice of α. This proves that $P_{\overline{X}_I}(f, \phi) \leq P_{\text{top}}(f, \phi) - L_{I_\delta}$. Since $P_{\underline{X}_I}(f, \phi) \leq P_{\overline{X}_I}(f, \phi)$, this finishes the proof of the theorem. □

3.2. *Proof of Theorem B.* Let us assume that both ϕ, ψ satisfy the Bowen condition and ψ is not cohomologous to a constant. Assume without loss of generality that $\int \psi \, d\mu_{f,\phi} = 0$. Our first purpose is to prove

$$P_{\overline{X}_c}(f, \phi) \leq P_{\text{top}}(f, \phi) - \min\left\{I_{f,\phi,\psi}\left(\int \psi \, d\mu_\phi + c\right),\ I_{f,\phi,\psi}\left(\int \psi \, d\mu_\phi - c\right)\right\},$$

where $I_{f,\phi,\psi}$ is the rate function of the large deviations function. Since $f|_\Lambda$ satisfies the specification property and ψ is not cohomologous to a constant, it follows that (see e.g. [**33**])

$$\left\{\alpha \in \mathbb{R} : \exists\, x \in M \text{ such that } \lim_{n\to\infty} \frac{1}{n} S_n\psi(x) = \alpha\right\} = \left\{\int \psi \, d\mu : \mu \text{ is } f\text{-invariant}\right\}$$

is a non-empty compact interval. By the level-1 large deviations principle for uniformly hyperbolic dynamics of Young [**38**], equations (2.3) and (2.4) hold with the rate function $I_{f,\phi,\psi}(s) = \sup\{-P_{\text{top}}(f, \phi) + h_\eta(f) + \int \phi \, d\eta : \int \psi \, d\eta = s\}$. Moreover, it follows from the functional analytic approach using transfer operators and the differentiability of the free energy function that $I_{f,\phi,\psi}$ is the Legendre transform of the free energy. On the one hand, using Theorem A and the previous upper bound,

$$P_{\overline{X}_c}(f, \phi) \leq P_{\text{top}}(f, \phi) - L_{c-\delta} \leq P_{\text{top}}(f, \phi) - \min\{I_{f,\phi,\psi}(c - \delta),\ I_{f,\phi,\psi}(c + \delta)\}$$

for all positive δ. Now, assume for simplicity that $0 < c = c_1 < c_2$ and $c_* = -c$ is defined by equation (2.5) (the other cases are analogous). We claim that it follows from the continuity and convexity of the rate function that if $X_c \neq \emptyset$ then

$$\begin{aligned}
P_{\overline{X}_c}(f, \phi) &= P_{\underline{X}_c}(f, \phi) = P_{X(-c)}(f, \phi) = P_{X([-c_2,-c_1])}(f, \phi) \\
&= P_{X(-c_2,-c_1)}(f, \phi) = P_{\text{top}}(f, \phi) - I_{f,\phi,\psi}(-c_1) \\
&= P_{\text{top}}(f, \phi) - I_{f,\phi,\psi}(c_*).
\end{aligned}$$

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

In fact, using [**33**], the topological pressure of the set $\{x \in M : \lim (1/n) S_n \psi(x) = c\}$ coincides with $\sup\{h_\eta + \int \psi \, d\eta : \eta \text{ is } f\text{-invariant and } \int \psi d\eta = c\}$. Then

$$\begin{aligned} P_{\text{top}}(f, \phi) - I_{f,\phi,\psi}(-c_1) = P_{X(-c_1)}(f, \phi) &\le P_{X(-c_2,-c_1)}(f, \phi) \\ &\le P_{X[-c_2,-c_1]}(f, \phi) \le P_{\underline{X}_{c_1}}(f, \phi) \\ &\le P_{\overline{X}_{c_1}}(f, \phi) \le P_{\text{top}}(f, \phi) - I_{f,\phi,\psi}(c_1) \\ &\le P_{\text{top}}(f, \phi) - I_{f,\phi,\psi}(-c_1). \end{aligned}$$

This proves the first part of the theorem. We proceed to prove the continuity results using that $P_{\underline{X}_{f,\phi,\psi,c}}(f, \phi) = P_{\text{top}}(f, \phi) - \min\{I_{f,\phi,\psi}(c), I_{f,\phi,\psi}(-c)\}$ whenever the set $\underline{X}_{f,\phi,\psi,c}$ is non-empty. On the one hand, it is well known that $\phi \mapsto P_{\text{top}}(f, \phi)$ is continuous in the C^0-topology. On the other hand, if $\Lambda = M$, then f is expanding and $P_{\text{top}}(f, \phi)$ varies continuously with f in the C^1-topology, since it coincides with the logarithm of the spectral radius of the transfer operator $\mathcal{L}_{f,\phi} : C^\alpha(M) \to C^\alpha(M)$ given by

$$\mathcal{L}_{f,\phi} \, g(x) = \sum_{f(y)=x} e^{\phi(y)} \, g(y).$$

In fact, due to the existence of a spectral gap property for $\mathcal{L}_{f,\phi}$, the spectral radius does vary continuously with respect to perturbations of the potential and the Legendre transform varies continuously with respect to the potential. We will provide a sketch of the proof now addressing also the continuity of these objects as a function of the dynamics f and observable ψ.

Given f, ϕ and ψ fixed, the spectral gap property for $\mathcal{L}_{f,\phi}$ implies that the free energy $\mathcal{E}_{f,\phi,\psi}(t) := \limsup_{n\to\infty} (1/n) \log \int e^{t S_n \psi} \, d\mu_{f,\phi}$ is well defined for all $t \in \mathbb{R}$ and, in fact, it verifies $\mathcal{E}_{f,\phi,\psi}(t) = P_{\text{top}}(f, \phi + t\psi) - P_{\text{top}}(f, \phi)$. In particular, if ψ is cohomologous to a constant, then $t \mapsto \mathcal{E}_{f,\phi,\psi}(t)$ is affine and otherwise $t \mapsto \mathcal{E}_{f,\phi,\psi}(t)$ is real analytic, strictly convex. Note also that for every $t \in \mathbb{R}$, the function $(f, \phi, \psi) \mapsto \mathcal{E}_{f,\phi,\psi}(t)$ is differentiable, the function $(\phi, \psi) \mapsto \mathcal{E}_{f,\phi,\psi}(t)$ is analytic and

$$(f, \phi, \psi) \mapsto \mathcal{E}'_{f,\phi,\psi}(t) = \int \psi \, d\mu_{f,\phi+t\psi}$$

is continuous (see e.g. [**3**]). Let $t \in \mathbb{R}$ be fixed. In order to establish the regularity of the rate function $I_{f,\phi,\psi}$ in what follows we assume, without loss of generality, that ψ is not cohomologous to a constant and that $m_{f,\phi} = \int \psi \, d\mu_{f,\phi} = 0$. Using that $\mathbb{R} \ni t \to \mathcal{E}_{f,\phi,\psi}(t)$ is strictly convex, its Legendre transform $I_{f,\phi,\psi}$ is well defined by

$$I_{f,\phi,\psi}(s) = \sup_{t\in\mathbb{R}} \{st - \mathcal{E}_{f,\phi,\psi}(t)\}.$$

This function is non-negative and strictly convex since $\mathcal{E}_{f,\phi,\psi}$ is also strictly convex, and $I_{f,\phi,\psi}(s) = 0$ if and only if $s = m_{f,\phi}$. Moreover, using the differentiability of the free energy function it is not hard to check the variational property

$$I_{f,\phi,\psi}(\mathcal{E}'_{f,\phi,\psi}(t)) = t\mathcal{E}'_{f,\phi,\psi}(t) - \mathcal{E}_{f,\phi,\psi}(t)$$

whenever the expressions make sense and, consequently, the rate function $I_{f,\phi,\psi}$ varies continuously with ϕ and ψ in the C^α-topology.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

Finally, we study the regularity of the function $v \mapsto P_{\overline{X}_c}(f_v, \phi)$ when $V \ni v \mapsto (f_v)_v$ is a continuous family of expanding maps on M and V is a compact metric space. Let $J \subset \mathbb{R}$ be a compact interval in the domain of $I_{f_v,\phi,\psi}$. From the previous variational relation we get that for any $s \in J$ there exists a unique $t = t(s, v)$ such that $s = \mathcal{E}'_{f_v,\phi_v,\psi_v}(t)$ and

$$I_{f_v,\phi,\psi}(s) = s \cdot t(s, v) - \mathcal{E}_{f_v,\phi,\psi}(t(s, v)). \tag{3.1}$$

Now, notice that the skew product

$$\begin{array}{rcl} F : V \times J & \to & V \times \mathbb{R} \\ (v, t) & \mapsto & (v, \mathcal{E}'_{f_v,\phi,\psi}(t)) \end{array}$$

is continuous and injective because it is strictly increasing along the fibers (using the strict convexity of the free energy function). Since $V \times J$ is a compact metric space, F is a homeomorphism onto its image $F(V \times J)$. In particular, this shows that for every $(v, s) \in F(V \times J)$ there exists a unique $t = t(v, s)$ varying continuously with (v, s) such that $F(v, t(v, s)) = (v, s)$ and $s = \mathcal{E}'_{f_v,\phi_v,\psi_v}(t)$. Finally, relation (3.1) above yields that $(s, v) \mapsto I_{f_v,\phi,\psi}(s)$ is continuous on $J \times V$. This finishes the proof of the continuity. □

3.3. *Proof of Corollary C.* Let $f : M \to M$ be a continuous map admitting a mixing repeller $\Lambda \subset M$, $\phi : M \to \mathbb{R}$ be a continuous potential so that μ_ϕ is the unique equilibrium state for f with respect to ϕ and $\mu_\phi \ll \nu$ where ν is a Gibbs measure. Given $c > 0$, consider $\overline{E}_c = \overline{X}_c \cap E_\psi \subset \overline{X_c}$. Taking into account Theorem B and Corollary B, part (1) is immediate and we need only prove the lower bound: if $\overline{E}_c \neq \emptyset$ then $P_{\overline{E}_c}(f, \phi) \geq P_{\overline{X}_c}(f, \phi)$.

Assume $\overline{E}_c \neq \emptyset$ for some $c > 0$. Under our assumptions it is well known that f satisfies the specification property and that for any f-invariant probability measure μ, there exists a sequence of f-invariant ergodic probability measures μ_n so that $\mu_n \to \mu$ in the weak* topology and $h_{\mu_n}(f) \to h_\mu(f)$ as $n \to \infty$ (cf. [**18**, Theorem B]). By Theorem B and the thermodynamical formulation of the large deviations rate function obtained by Young [**38**], we know that

$$P_{\overline{X}_c}(f, \phi) = P_{\text{top}}(f, \phi) - I_{f,\phi,\psi}(c_*) = \sup_\eta \left\{ h_\eta(f) + \int \phi \, d\eta \right\} \tag{3.2}$$

where $|c_*| = |c|$ and the supremum is taken over all f-invariant probability measures η so that $|\int \psi \, d\eta - \int \psi \, d\mu_\phi| \geq c_*$. Assume, for simplicity, that $c_* = c$ (the other case is analogous). Observe that

$$\begin{aligned} &\sup \left\{ h_\eta(f) + \int \phi \, d\eta : \left| \int \psi \, d\eta - \int \psi \, d\mu_\phi \right| \geq c \right\} \\ &\quad = \sup \left\{ h_\eta(f) + \int \phi \, d\eta : \left| \int \psi \, d\eta - \int \psi \, d\mu_\phi \right| > c \right\} \end{aligned}$$

by the continuity of the rate function $c \mapsto I_{f,\phi,\psi}(c)$ (since it coincides with the Legendre transform of the free energy function). Together with the variational relation (3.2), this yields that for any $\gamma > 0$ one can take two f-invariant probability measures η_1, η_2 such that

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

(i) $|\int \psi\, d\eta_i - \int \psi\, d\mu_\phi| > c$;
(ii) $h_{\eta_i}(f) + \int \phi\, d\eta_i \geq P_{\overline{X}_c}(f, \phi) - 2\gamma$; and
(iii) $\int \psi\, d\eta_1 \neq \int \psi\, d\eta_2$

for $i = 1, 2$. Taking the approximation in entropy by f-invariant and ergodic probability measures, there are distinct ergodic probability measures ν_1 and ν_2 satisfying $|\int \psi\, d\nu_i - \int \psi\, d\mu_\phi| > c$, $\int \psi\, d\nu_1 \neq \int \psi\, d\nu_2$ and $h_{\nu_i}(f) + \int \phi\, d\nu_i \geq P_{\overline{X}_c}(f, \phi) - \gamma$ for $i = 1, 2$. Observe that $\overline{X}_c$ is an f-invariant set and the ergodicity together with the first property above implies that $\nu_i(\overline{X}_c) = 1$.

Now the proof follows the same lines as the proof of [**35**, Theorem 2.6]. Consider a strictly decreasing sequence $(\delta_k)_{k\geq 1}$ of positive numbers converging to zero and a strictly increasing sequence of positive integers $(\ell_k)_{k\geq 1}$, such that the sets

$$Y_{2k+i} = \left\{ x \in \overline{X}_c : \left| \frac{1}{n} S_n \psi(x) - \int \psi\, d\nu_i \right| < \delta_k \text{ for every } n \geq \ell_k \right\}$$

satisfy $\nu_i(Y_{2k+i}) > 1 - \gamma$ for every k $(i = 1, 2)$. Consider the fractal set F given *ipsis literis* by the construction in §3.1 with ν_i replacing μ_i, $P_{\overline{X}_c}(f, \phi)$ replacing C and ψ replacing φ. From the construction there is a sequence $(t_k)_{k\geq 1}$ such that

$$\lim_{k\to\infty} \left| \frac{1}{t_{2k+i}} S_{t_{2k+i}} \psi(x) - \int \psi\, d\nu_i \right| = 0 \quad \text{for every } x \in F$$

and $P_F(f, \psi) \geq C - 8\gamma$. In particular, F is contained in the irregular set E_ψ. Since γ was chosen arbitrarily and $F \subset E_\psi$, to complete the proof of the corollary it is enough to prove that $F \subset \overline{X}_c$. This actually follows from item (i) above, since for any $x \in F$,

$$\begin{aligned} \limsup_{k\to\infty} \left| \frac{1}{t_{2k+2}} S_{t_{2k+2}} \psi(x) - \int \psi\, d\mu_\phi \right| &\geq \limsup_{k\to\infty} \left[\left| \int \psi\, d\nu_2 - \int \psi\, d\mu_\phi \right| \right. \\ &\quad \left. - \left| \frac{1}{t_{2k+2}} S_{t_{2k+2}} \psi(x) - \int \psi\, d\mu_\phi \right| \right] \\ &\geq c. \end{aligned}$$

This finishes the proof of the corollary. □

3.4. *Proof of Theorem C.* For the proof of the theorem, we will need the following auxiliary lemma, which will play the same role as Lemma 3.1 in the proof of Theorem A. It is here that we need the metric on $\mathcal{M}_1$ to be translation invariant and affine.

LEMMA 3.2. *Let $c > 0$ be given. For any $\delta > 0$, there exists $\varepsilon_\delta > 0$ and $N = N_\delta \in \mathbb{N}$ such that $B(x, n, \varepsilon) \subset Y_{\mu,c-\delta,n}$ for all $0 < \varepsilon < \varepsilon_\delta$, $n \geq N$ and $x \in Y_{\mu,c,n}$.*

Proof. Since $M \in x \mapsto \delta_x \in \mathcal{M}_1$ is uniformly continuous, given $\delta > 0$ there exists $\varepsilon_\delta > 0$ such that if $d(x, y) < \varepsilon_\delta$ we have $d(\delta_x, \delta_y) < \delta$. Hence, if $x \in Y_{c,n}$ and $y \in B(x, n, \varepsilon)$, we have $d(\delta_{y,n}, \mu) \geq d(\delta_{x,n}, \mu) - d(\delta_{x,n}, \delta_{y,n}) \geq c - (1/n) \sum_{i=0}^{n-1} d(\delta_x, \delta_y) \geq c - \delta$, and thus $y \in Y_{\mu,c,n}$, which proves the lemma. □

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

We proceed with the proof of the theorem assuming that $\mu = \mu_{f,\phi}$ is the unique equilibrium state for the continuous map f with respect to the continuous potential ϕ and also $\overline{Y}_{\mu,c} \neq \emptyset$. In order to prove that $P_{\overline{Y}_{\mu,c}}(f, \phi) \leq P_{\text{top}}(f, \phi) - \inf_{d(\eta,\mu)=c} Q(\eta)$ is strictly smaller than the topological pressure $P_{\text{top}}(f, \phi)$, we proceed to cover $\overline{Y}_{\mu,c}$ with a properly chosen family of dynamical balls. Fix $\delta > 0$ small and $\alpha > P_{\text{top}}(f, \Phi) - \inf_{d(\eta,\mu)\geq c-\delta} Q(\eta)$. Given $\varepsilon > 0$ small and $N \in \mathbb{N}$, for any $x \in \overline{Y}_{\mu,c}$ pick $m(x) \geq N$ in such a way that $x \in Y_{\mu,c-(\delta/2),m(x)}$ and consider $\mathcal{G}_N := \{(x, m(x)) : x \in \overline{Y}_{\mu,c}\}$. Then

$$\overline{Y}_{\mu,c} \subset \bigcup_{(x,n)\in\mathcal{G}_N} B(x, n, \varepsilon)$$

and $B(x, n, \varepsilon) \subset \overline{Y}_{\mu,c-\delta,n}$, for all $x \in Y_{\mu,c-(\delta/2),n}$, $n \geq N$ and ε small (by Lemma 3.2). Therefore, we can proceed as in the proof of Theorem A and extract a subset $\hat{\mathcal{G}}_N \subset \mathcal{G}_N$ in such a way that if (x, l) and (y, l) belong to $\hat{\mathcal{G}}_N$ then $B(x, l, (\varepsilon/2)) \cap B(x, l, (\varepsilon/2))=\emptyset$. If $\zeta = (-P + \alpha + \inf_{d(\eta,\mu)\geq c-\delta} Q(\eta))/2 > 0$, then by the large deviations upper bound, for every closed set U one has $\mu\left(x \in M : \delta_{x,n} \in U\right) \leq \exp(-n[\inf_{\eta\in U} Q(\eta) - \zeta])$ provided that $n \geq N$ is large enough. This, together with the Gibbs property for ν, yields

$$\begin{aligned}
\sum_{(x,n)\in\hat{\mathcal{G}}_N} e^{-\alpha n + S_n\phi(x)} &\leq K(\varepsilon) \sum_{n\geq N} \sum_{x\in\hat{\mathcal{G}}_{N,n}} e^{(P-\alpha)n} \nu(B(x, n, \varepsilon)) \\
&\leq K(\varepsilon) K\left(\frac{\varepsilon}{2}\right) \sum_{n\geq N} e^{(P-\alpha)n} \nu\Bigg(\bigcup_{x\in\hat{\mathcal{G}}_{N,n}} B\left(x, n, \frac{\varepsilon}{2}\right)\Bigg) \\
&\leq K(\varepsilon) K\left(\frac{\varepsilon}{2}\right) \sum_{n\geq N} e^{(P-\alpha)n} \nu(\overline{Y}_{\mu,c-\delta,n}) \\
&\leq K(\varepsilon) K\left(\frac{\varepsilon}{2}\right) \sum_{n\geq N} \exp n\Big(P - \alpha - \inf_{d(\eta,\mu)\geq c-\delta} Q(\eta) + \zeta\Big) \\
&\leq K(\varepsilon) K\left(\frac{\varepsilon}{2}\right) \sum_{n\geq N} e^{-\zeta n},
\end{aligned}$$

which is finite and independent of N. This proves that $P_{\overline{Y}_{\mu,c}}(f, \phi) \leq P_{\text{top}}(f, \phi) - \inf_{d(\eta,\mu)\geq c-\delta} Q(\eta)$. Since Q is lower semicontinuous it follows that

$$P_{\overline{Y}_{\mu,c}}(f, \phi) \leq P_{\text{top}}(f, \phi) - \inf_{d(\eta,\mu)\geq c} Q(\eta).$$

For the proof of the second part of the theorem we make use of the level-2 large deviations principles obtained by Zhou and Chen [**39**] under the assumptions of almost product structure and the uniform separation properties. Let $0 < c_1 < c_2$ be such that $\overline{Y}_{\mu,c_1} \neq \emptyset$. Using [**39**], given a compact connected subset in $\mathcal{M}_1$, the topological pressure of the set $Y(C) := \{x \in M : \lim_{n\to+\infty} \delta_{x,n} \in C\}$ coincides with $\inf\{h_\eta(f) + \int \psi \, d\eta : \eta \text{ is } f\text{-invariant and } \eta \in C\}$. On the other hand, by [**13**], $Q(\eta) = P_{\text{top}}(f, \phi) - h_\eta(f) - \int \phi \, d\eta$. Thus, using that the metric entropy is linear convex and the choice of the metric

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

on $\mathcal{M}_1$, we have

$$\begin{aligned} P_{\text{top}}(f,\phi) - \inf_{d(\eta,\mu)=c_1} Q(\eta) &= P_{Y(\partial B(\mu,c_1))}(f,\phi) \le P_{Y(B(\mu,c_1,c_2))}(f,\phi) \\ &\le P_{Y(\overline{B(\mu,c_1,c_2)})}(f,\phi) \le P_{Y(B(\mu,c_1))}(f,\phi) \\ &\le P_{Y(\overline{B(\mu,c_1)})}(f,\phi) \le P_{\overline{Y}_{\mu,c_1}}(f,\phi) \\ &\le P_{\underline{Y}_{\mu,c_1}}(f,\phi) \le P_{\text{top}}(f,\phi) - \inf_{d(\eta,\mu)\ge c_1} Q(\eta) \\ &\le P_{\text{top}}(f,\phi) - \inf_{d(\eta,\mu)=c_1} Q(\eta), \end{aligned}$$

proving all quantities coincide. This finishes the proof of the theorem. □

3.5. *Proof of Theorem D.* This section is devoted to the proof of Theorem D, which claims that if the set $\underline{X}_c$ is non-empty then it has smaller topological pressure. The strategy for the proof is similar to that used for Theorem A, with the difficulty that in the non-uniformly expanding setting the Gibbs property holds at a sequence of moments that does depend on the point. For that reason we shall give a sketch of the proof with the main ingredients. Since μ is a weak Gibbs measure, there exists $\varepsilon_0 > 0$ such that the following property holds: for every $0 < \varepsilon < \varepsilon_0$ there exists $K(\varepsilon) > 0$ and for μ-almost every x there exists a sequence $n_k(x) \to \infty$ such that

$$K(\varepsilon)^{-1} \le \frac{\mu(B(x, n_k(x), \varepsilon))}{e^{-n_k(x)P + S_{n_k(x)}\phi(x)}} \le K(\varepsilon).$$

Assume the weak Gibbs property holds for *all* points in the invariant set $\Lambda = H$ and in what follows consider $\underline{X}_I := \underline{X}_I \cap \Lambda$.

Let $\delta > 0$ be arbitrary. We proceed to prove that $P_{\underline{X}_I}(f,\phi) \le P_{\text{top}}(f,\phi) - L_{I_\delta}$ is strictly smaller than the topological pressure. Consider $\alpha > P_{\text{top}}(f,\Phi) - L_{I_\delta}$ given and take $\varepsilon > 0$ arbitrarily small and $N \in \mathbb{N}$ arbitrarily large in what follows. One can write

$$\underline{X}_I \subset \bigcup_{\ell\ge 1} \bigcap_{j\ge\ell} X_{I_\delta,j}$$

where, as before, $X_{I,n} = \{x \in M : (1/n)S_n\psi(x) \in I\}$. It is not hard to check that for any $x \in \underline{X}_I$ there exists a sequence of positive integers $(m_j(x))_{j\in\mathbb{N}}$ converging to infinity such that $x \in X_{I_\delta,m_j(x)}$ and $m_j(x)$ is a moment at which the Gibbs property holds. Therefore, one can pick $m(x) \ge N$ in such a way that $x \in X_{I_\delta,m(x)-1}$ and consider $\mathcal{G} := \{(x, m(x)) : x \in \overline{X}_I\}$. Now the proof proceeds with the estimates used in the proof of Theorem A.

Remark 3.3. In the previous proof we did not require the times at which the Gibbs property holds to have positive density at infinity as in usual notions of non-lacunary Gibbs measures. In particular, this gives a wider range of applications.

Remark 3.4. Actually, let us mention that we could not estimate the topological pressure of the larger set $\overline{X}_c$. In fact, for that purpose we would need to guarantee that for each point there would exist a sequence of instants at which, simultaneously, the Gibbs property and the time averages being far from the time average occur. Nevertheless, this can be verified in examples.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

4. *Examples and applications*

4.1. *Hyperbolic diffeomorphisms and flows.* Using the fact that hyperbolic sets admit Markov partitions (cf. [**6**]) then the results of Theorem A also hold in this hyperbolic setting, via semi-conjugation to bilateral shifts and Remark 2.2. Axiom A flows $(Y_t)_t$ are also semi-conjugate to suspension flows over subshifts of finite type, by [**7**]. Recall that given a subshift of finite type $\sigma : \Sigma \to \Sigma$ and a ceiling function $h : \Sigma \to \mathbb{R}^+$ bounded away from zero and infinity, the associated *suspension flow* $(S_t)_t$ is defined in $\Sigma_h = \{(x, t) \in \Sigma \times \mathbb{R}_+ : 0 \le t \le h(x)\}$ with the identification between the pairs $(x, h(x))$ and $(\sigma(x), 0)$. The semiflow defined on Σ_h by $S_t(x, r) = (\sigma^n(x), r + t - \sum_{i=0}^{n-1} h(\sigma^i(x)))$, where $n = n(x, r + t) \in \mathbb{N}_0$, is uniquely defined by $\sum_{i=0}^{n-1} h(\sigma^i(x)) \le r + t < \sum_{i=0}^{n} h(\sigma^i(x))$. It is clear that $(1/(n(x, T + s))) \sum_{i=0}^{n(x,T+s)-1} h(\sigma^i(x)) \le (T + s)/(n(x, T + s)) < 1/(n(x, T + s)) \sum_{i=0}^{n(x,T+s)} h(\sigma^i(x))$ and $n(x, T + s) \to \infty$ as $T \to \infty$. Given $\psi \in C(\Sigma_h, \mathbb{R})$, define $\bar{\psi} \in C(\Sigma, \mathbb{R})$ by $\bar{\psi}(x) = \int_0^{h(x)} \psi(x, t)\, dt$. Let μ and μ_Σ be the unique equilibrium states for $(X_t)_t$ with respect to ϕ and for σ with respect to $\overline{\phi}$, respectively, which are known to satisfy $\mu = \mu_\Sigma \times \mathrm{Leb}_1 / \int h\, d\mu_\Sigma$ (cf. [**7**]). If $\beta_1, \beta_2, \beta_3 > 0$ are small, it is not hard to check that

$$\overline{X}_c = \left\{(x, s) \in \Sigma_h : \limsup_{T\to\infty} \left| \frac{1}{T} \int_0^T \psi(S_t(x, s))\, dt - \int \psi\, d\mu \right| \ge c \right\}$$

is contained in the union of the $(S_t)_t$-invariant sets

$$\begin{aligned} \overline{X}_{c,h} &= \left\{(x, s) \in \Sigma_h : \limsup_{T\to\infty} \left| \frac{n(x, T + s)}{T + s} - \frac{1}{\int h\, d\mu_\Sigma} \right| \ge \beta_2 \right\} \\ &\subseteq \left\{(x, s) \in \Sigma_h : \limsup_{T\to\infty} \left| \frac{1}{n(x, T + s)} \sum_{i=0}^{n(x,T+s)} h(\sigma^i(x)) - \int h\, d\mu_\Sigma \right| \ge \beta_3 \right\} \end{aligned}$$

and

$$\overline{X}_{c,\overline{\psi}} = \left\{(x, s) \in \Sigma_h : \limsup_{T\to\infty} \left| \frac{1}{n(x, T)} \sum_{i=0}^{n(x,T+s)} \overline{\psi}(\sigma^i(x)) - \int \overline{\psi}\, d\mu_\Sigma \right| \ge \beta_1 \right\}.$$

Using $P_{\overline{X}_c}((S_t)_t, \phi) \le \max\{P_{\overline{X}_{c,h}}((S_t)_t, \psi), P_{\overline{X}_{c,\overline{\phi}}}((S_t)_t, \phi)\}$ and Theorem A we deduce by semi-conjugacy that $P_{\overline{X}_c}((X_t)_t, \phi) < P_{\mathrm{top}}((X_t)_t, \phi)$ for any Axiom A flow, any Hölder potential $\phi : M \to \mathbb{R}$, any continuous $\psi : M \to \mathbb{R}$ and $c > 0$.

4.2. *Manneville–Pomeau maps.* If $\alpha \in (0, 1)$, let $f_\alpha : \mathbb{S}^1 \to \mathbb{S}^1$ be the local homeomorphism given by $f_\alpha(x) = x(1 + 2^\alpha x^\alpha)$ for $0 \le x \le \frac{1}{2}$ and by $f_\alpha(x) = 2x - 1$ whenever $\frac{1}{2} < x \le 1$. This map satisfies the specification property since it is topologically conjugate to the double expanding map. Pollicott and Weiss [**30**] established a multifractal formalism for the Lyapunov spectrum associated with this class of transformations and proved precise formulas for the dimension of the level sets of points with the same Lyapunov exponent. Clearly, $(1/n) \log |(f_\alpha^n)'(x)| = (1/n) \sum_{j=0}^{n-1} \psi(f_\alpha^j(x))$ with $\psi(x) = \log |(f_\alpha)'(x)|$. For every $t \in (-\infty, 1)$, there exists a unique equilibrium state μ_t with respect to the Hölder continuous potential $\phi_t = -t \log |(f_\alpha)'(x)|$, and it is well known that there are two equilibrium states for f_α with respect to $-\log |(f_\alpha)'(x)|$, namely, an acip

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

(absolutely continuous invariant probability) μ_1 and the Dirac measure δ_0. Moreover, for every $t \le 1$, the equilibrium state μ_t for f with respect to the potential ϕ_t satisfies a weak Gibbs property: there are constants K_n so that $\limsup_{n\to\infty} (1/n) \log K_n = 0$ and

$$K_n{}^{-1} e^{-nP_t} |(f^n)'(x)|^t \le \mu_t(\mathcal{P}^{(n)}(x)) \le K_n e^{-nP_t} |(f^n)'(x)|^t$$

for all $x \in [0, 1]$ and $n \ge 1$, where $\mathcal{P}$ is the Markov partition for f, $\mathcal{P}^{(n)}(x)$ is the element of the partition $\mathcal{P}^{(n)} = \bigvee_{j=0}^{n-1} f^{-j}\mathcal{P}$ that contains x and $P_t = P_{\text{top}}(f, \phi_t)$. By the Ruelle inequality, all measures μ_t are expanding. Our results do not apply for μ_1 since polynomial upper and lower bounds for Hölder continuous observables have been established in [**24**, **25**, **29**]. If $|t|$ is small, $c > 0$ and ψ is *continuous*, then

$$\limsup_{n\to\infty} \frac{1}{n} \log \mu_t\left(x \in M : \left|\frac{1}{n} S_n\psi(x) - \int \psi \, d\mu_t\right| \ge c\right) < 0$$

(cf. [**36**]). If, in addition, we assume ψ is Hölder continuous then there exists an interval $J \subset \mathbb{R}$ such that the following local large deviations principle holds:

$$\limsup_{n\to\infty} \frac{1}{n} \log \mu_t\left(x \in M : \frac{1}{n} S_n\psi(x) \in [a, b]\right) \le -\inf_{s\in[a,b]} I_{f,\phi_t,\psi}(s)$$

and

$$\liminf_{n\to\infty} \frac{1}{n} \log \mu_t\left(x \in M : \frac{1}{n} S_n\psi(x) \in (a, b)\right) \ge -\inf_{s\in(a,b)} I_{f,\phi_t,\psi}(s)$$

for every $[a, b] \subset J$, and continuity and smoothness of the rate function are also obtained (cf. [**3**]). As a consequence, we deduce from Theorem D that for all $c > 0$ satisfying $[\int \psi \, d\mu_t - c, \int \psi \, d\mu_t + c] \subset J$, either $\underline{X}_c = \emptyset$ or else

$$\begin{aligned} P_{\overline{X}_c}(f, \phi_t) = P_{\underline{X}_c}(f, \phi_t) &= P_{X(c)}(f, \phi_t) = P_{X([c_1,c_2])}(f, \phi_t) \\ &= P_{X(c_1,c_2)}(f, \phi_t) = P_{\text{top}}(f, \phi_t) - I_{f,\phi_t,\psi}(c) \end{aligned}$$

for $c = \max\{|c_1|, |c_2|\}$. Furthermore, the right-hand-side expression varies continuously with c and also varies continuously with f, ϕ_t and ψ. Although the set of irregular points has full topological entropy $\log 2$ (see e.g. [**35**]), the set of Lyapunov irregular points whose Birkhoff averages remain far from $\lambda(\mu_0) := \int \psi \, d\mu_0$ for all large iterates has topological entropy strictly smaller than $\log 2$.

4.3. *Multimodal maps.* Our results also apply to a broad class of transitive multimodal interval maps f with finitely many non-degenerate critical points with negative Schwarzian derivative, considered in [**8**]. If there exist $C > 0$ and $\beta > 2\ell - 1$ such that $|Df^n(c)| \ge n^\beta$ for every critical point c and all $n \ge 1$ (where ℓ denotes the maximal order of the critical points), then it follows from [**8**, Theorem 1] that there exists $t_1 < 1$ such that for all $t \in (t_1, 1)$: (i) there exists a unique equilibrium state μ_t for f with respect to the potential $\varphi_t = -t \log |Df|$; (ii) μ_t has a compatible inducing scheme with exponential tails, and hence it has exponential decay of correlations; and (iii) μ_t has positive Lyapunov exponent almost everywhere. Moreover, there is a conformal probability measure ν_t such that $J_{\nu_t} f(x) = e^{P(t)} |f'(x)|^{-t}$ almost everywhere and $\mu_t \ll \nu_t$, where $P(t) = P_{\text{top}}(f, -t \log |f'|)$. In addition, since μ_t has only positive Lyapunov exponents,

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press




FIGURE 2. Bowen eye attractor.

almost every point has infinitely many hyperbolic times. If n is a hyperbolic time for x, then the Jacobian $J_{\nu_t} f^n$ has bounded distortion and, consequently, ν_t satisfies the weak Gibbs property. Moreover, by property (ii) above with the results in [**1**], one has that μ_t has exponential large deviations and so it satisfies the assumptions of Theorem D.

4.4. *Higher dimensional non-uniformly expanding maps.* Assume that M is a compact metric space where the Besicovitch covering lemma applies and let $f : M \to M$ be a local homeomorphism, such that there exists a bounded function $x \mapsto L(x)$, such that for every $x \in M$ there is a neighborhood U_x of x such that $f_x : U_x \to f(U_x)$ is invertible and $d(f_x^{-1}(y), f_x^{-1}(z)) \leq L(x)\, d(y, z)$ for every $y, z \in f(U_x)$. Assume also that every point has finitely many preimages and that the level sets for the degree $\{x : \#\{f^{-1}(x)\} = k\}$ are closed. Given $x \in M$, set $\deg_x(f) = \#f^{-1}(x)$ and define $h(f) = \liminf_{n\to\infty} (1/n) \log[\min_{x\in M} \deg_x(f^n)]$. Assume that every point in M has at least $e^{h(f)}$ preimages by f and that f is uniformly expanding outside $\mathcal{A}$ and not contracting too much inside $\mathcal{A}$ (see [**37**] for precise statements). A concrete example can be built on the torus by taking a linear expanding map $f_0 : \mathbb{T}^d \to \mathbb{T}^d$, fixing a small open covering $\mathcal{P}$ of $\mathbb{T}^d$ and deforming f_0 on a small neighborhood of a fixed point p inside $P_1 \in \mathcal{P}$ by a pitchfork bifurcation in such a way that p becomes a saddle for the perturbed local homeomorphism f (cf. [**9**, **37**]). Then [**3**, **36**] imply that local large deviation estimates hold for all equilibrium states associated with Hölder continuous potentials with low variation, and Theorem D holds in this context.

4.5. *Bowen-eye like systems and a counterexample.*

4.5.1. *Distinction of $\overline{X}_c$ and $\underline{X}_c$.* We shall present a simple example of a discrete dynamical system f, potential ϕ, observable ψ and constant $c > 0$ such that $\overline{X}_c \neq \underline{X}_c$. The map f corresponds to the time-one map of a flow known as the Bowen eye. The map f has three fixed points p_1, p_2 and p_3 (corresponding to the points on the figure organized from the left in Figure 2) and is such that $\{p_1, p_2, p_3\} = \mathrm{Per}(f) = R(f)$ while the non-wandering set is formed by the fixed point p_2 and the closure D of the two separatrices corresponding to the singularities p_1 and p_3 of the original vector field. Moreover, it is well known that for *every* x in the inner region of the plane determined by D (except p_2), the empirical measures $(1/n) \sum_{j=0}^{n-1} \delta_{f^j(x)}$ have the Dirac measures δ_{p_1}

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

and δ_{p_3} as accumulation points. If $\phi : \mathbb{R}^2 \to \mathbb{R}$ denotes the projection on the x-coordinate then, by the variational principle, $P_{\text{top}}(f, \phi) = \sup\{h_\mu(f) + \int \phi \, d\mu\} = \phi(p_3)$ and δ_{p_3} is the unique equilibrium state for ϕ. On the other hand, for $0 < c < d(p_2, p_3)$ it is clear that $\underline{X}_c = W^s(p_1) \cup \{p_2\}$ and $\overline{X}_c = D \setminus (W^s(p_3) \cup \{p_2\})$. However, in this case one has $P_{\underline{X}_c}(f, \phi) = P_{\overline{X}_c}(f, \phi)$.

4.5.2. *A counterexample.* Despite the fact that the topological pressure of both sets $\underline{X}_c$ and $\overline{X}_c$ do coincide, the previous Bowen-eye construction gives some light on how to construct an example where $\overline{CP}_{\underline{X}_c}(f, \phi) < \overline{CP}_{\overline{X}_c}(f, \phi)$, where $\overline{CP}_\Lambda$ denotes the upper Carathéodory capacity of the set Λ (see e.g. [**26**, §11]). The following can be realized as a non-compact invariant set of a horseshoe. Let $\sigma : \Sigma_A \to \Sigma_A$, with $\Sigma_A \subset \{0, 1, 2, 3\}^{\mathbb{Z}}$, be the subshift of finite type associated with the transition matrix

$$A = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Now, consider the σ-invariant subset $\Sigma \subset \Sigma_A$ that contains the four fixed points for the shift σ (corresponding to the constant sequences) and is such that for any $x = (x_n)_n \in \Sigma \setminus \{\underline{3}\}$ it holds that $\limsup_{n\to\infty} (1/2n)\#\{|j| \le n : x_j \in \{1, 2\}\} = 1$ and $\liminf_{n\to\infty} (1/(2n))\#\{-n \le j \le n : x_j \in \{1, 2\}\} = 0$.

Let ϕ be a continuous potential so that the unique equilibrium state is $\mu_\phi = \delta_{\underline{0}}$ (such a potential can be built to be non-negative following the ideas of Hofbauer [**21**, pp. 226, 239]) and consider the continuous observable $\psi = \chi_{[0]}$. Notice that $\int \psi \, d\mu_\phi = 1$ and for $c > 0$ small enough we get that $\underline{X}_c = \{\underline{3}\}$ while $\overline{X}_c = \Sigma \setminus \{\underline{0}\}$. Since $\phi|_{\underline{X}_c} \equiv 0$ and $\underline{X}_c = \{\underline{3}\}$ then $\overline{CP}_{\underline{X}_c}(f, \phi) = h_{\underline{X}_c}(f) = 0$. On the other hand, since ϕ is non-negative then $\overline{CP}_{\overline{X}_c}(f, \phi) \ge \overline{CP}_{\overline{X}_c}(f, 0)$, which we now claim to be strictly positive. In fact, if $0 < \alpha < \log 2$, we will prove that $m_\alpha(f, \overline{X}_c) = +\infty$ and deduce that $\overline{CP}_{\overline{X}_c}(f, 0) > 0$. Recall that $m_\alpha(f, \overline{X}_c) = \lim_{\text{diam}(\mathcal{U})\to 0} m_\alpha(f, \overline{X}_c, \mathcal{U})$, where $m_\alpha(f, \overline{X}_c, \mathcal{U}) = \lim_{N\to\infty} m_\alpha(f, \overline{X}_c, \mathcal{U}, N)$, and

$$m_\alpha(f, \overline{X}_c, \mathcal{U}, N) = \inf\Big\{ \sum_{U\in\mathcal{G}_N} e^{-\alpha N} : \mathcal{G}_N \text{ is subcover of } \bigvee_{0\le j\le N} \sigma^{-j}\mathcal{U} \Big\}.$$

Let $\varepsilon > 0$ be small and fixed (to be made precise later and depending only on α). For any $\ell \ge 1$, let us consider an open cover $\mathcal{U}_\ell$ of $\overline{X}_c$ formed by cylinders as follows: a $(2n+1)$-cylinder $U = [x_{-n}, \ldots, x_n]$ belongs to $\mathcal{U}_\ell$ if and only if $n \ge \ell$ is the smallest positive integer such that

$$\#\{|j| \le n : x_j \in \{1, 2\}\} \ge (2n + 1)(1 - \varepsilon). \tag{4.1}$$

In fact, given $x = (x_j)_j \in \underline{X}_c$, and defining $n(x) \ge \ell$ to be the first instant such that equation (4.1) holds, it follows that $[x_{-n}, \ldots, x_n]$ belongs to $\mathcal{U}_\ell$ and so $\mathcal{U}_\ell$ covers $\underline{X}_c$. Moreover, by construction, the elements of $\mathcal{U}_\ell$ are all disjoint, every such element contains at least one point of $\underline{X}_c$ and the diameter of $\mathcal{U}_\ell$ goes to zero as $\ell \to \infty$. Thus, $m_\alpha(f, \overline{X}_c) = \lim_{\ell\to\infty} m_\alpha(f, \overline{X}_c, \mathcal{U}_\ell)$. Observe also that $\#\, U_\ell \ge 2^{(2\ell+1)(1-\varepsilon)}$, which corresponds to the

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press




FIGURE 3. Discontinuity of the pressure function.

number of disjoint cylinders of length $(2\ell+1)$ satisfying (4.1). Therefore, given any $N \gg 1$, any subcover $\mathcal{G}_{N,\ell}$ of the space of cylinders $\bigvee_{0\le j\le N} \sigma^{-j}\mathcal{U}_\ell$ that covers $\overline{X}_c$ coincides with the space of all $(N+\ell)$-cylinders. If one writes $N+\ell = (2\ell+1)s + r$ with $s \ge 1$ and $0 \le r \le 2\ell$ then there are at least $2^{(N+\ell-r)(1-\varepsilon)}$ such cylinders (just by considering N-concatenations of $(2\ell+1)$-cylinders that satisfy equation (4.1)). Thus, if $\varepsilon > 0$ is chosen to be small, then it follows that

$$m_\alpha(f, \overline{X}_c, \mathcal{U}_\ell) \ge \limsup_{N\to\infty} \sum_{U\in\mathcal{G}_{N,\ell}} e^{-\alpha N} \ge \limsup_{N\to\infty} e^{-\alpha N}\, 2^{(N-\ell)(1-\varepsilon)} = +\infty.$$

Hence, $\overline{CP}_{\overline{X}_c}(f, 0) \ge \log 2 > 0$, which proves our claim. Let us mention that the question remains of whether it is possible to construct an example where $P_{\underline{X}_c}(f,\phi) < P_{\overline{X}_c}(f,\phi)$.

4.6. *Discontinuity and non-strict monotonicity of the pressure function.*

4.6.1. *Porcupine-like horseshoes.* To present an example where there is discontinuity and non-strict monotonicity of the pressure function $c \mapsto P_{\overline{X}_c}(f,\phi)$, we use the class of local diffeomorphisms f studied by Díaz, Gelfert and Rams that exhibit porcupine-like horseshoes. In fact, it follows from the analysis of the Lyapunov spectrum in the central direction (see [**16**, Remark 5.4] and [**17**]) that there are constants $\lambda < 0 < \tilde{\beta} < \beta$ such that

$$\liminf_{n\to\infty} \frac{1}{n} \log \|Df^n|_{E^c}(x)\|, \quad \limsup_{n\to\infty} \frac{1}{n} \log \|Df^n|_{E^c}(x)\| \in [\log\lambda, \log\tilde{\beta}] \cup \{\log\beta\}$$

and also that there exists a unique point Q (indeed, it is a fixed point, by f) so that the central Lyapunov exponent is $\log\beta > 0$. Let us consider the Hölder continuous potential $\phi_t = -t \log \|Df|_{E^c}\|$ for a large value of negative t and the observable $\psi = \log\|Df|_{E^c}\|$. It follows from [**16**, Proposition 5.6] that for all $t \ll 0$ the Dirac measure δ_Q is the unique equilibrium state for f with respect to ϕ_t, and, consequently, $P_{\text{top}}(f,\phi_t) = -t\log\|Df(Q)|_{E^c}\| = -t\log\beta$. On the other hand, if $c \ne \log\beta$ then, taking all invariant measures with central Lyapunov exponent equal to c, it follows that $\sup\{h_\eta(f) + \int -t\log\|Df(x)|_{E^c}\|\, d\eta : \lambda(\eta) = c\} \le h_{\text{top}}(f) - t\log\tilde{\beta}$ is strictly smaller than $P_{\text{top}}(f,\phi_t)$. This shows the discontinuity of the pressure function $c \mapsto P_{\overline{X}_c}(f,\phi_t)$ where $\overline{X}_c$ is associated with the observable ψ (see Figure 3). Actually, the same argument

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

leads to the proof that

$$\begin{aligned}&\sup\left\{h_\eta(f)+\int -t\log\|Df(x)|_{E^c}\|\,d\eta : \lambda(\eta)\in[\log\lambda,\log\tilde{\beta}]\right\}\\&\quad=\sup\left\{h_\eta(f)+\int -t\log\|Df(x)|_{E^c}\|\,d\eta : \lambda(\eta)\in[\log\lambda,\log\beta)\right\}\\&\quad<P_{\mathrm{top}}(f,\phi_t),\end{aligned}$$

and so there exists an interval of constancy for this pressure function.

Acknowledgements. This work was partially supported by CNPq and is part of the first author's PhD thesis at Federal University of Bahia. The authors are deeply grateful to V. Climenhaga, D. Kwietniak and M. Todd for useful comments and to the anonymous referee for many suggestions that helped to improve the manuscript.

References

[**1**] J. F. Alves, S. Luzzatto, J. Freitas and S. Vaienti. From rates of mixing to recurrence times via large deviations. *Adv. Math.* **228** (2011), 1203–1236.
[**2**] L. Barreira, Y. Pesin and J. Schmeling. Multifractal spectra and multifractal rigidity for horseshoes. *J. Dyn. Control Syst.* **3**(1) (1997), 33–49.
[**3**] T. Bomfim, A. Castro and P. Varandas. Differentiability of thermodynamical quantities in non-uniformly expanding dynamics. *Preprint*, 2012, arXiv:1205.5361.
[**4**] T. Bomfim and P. Varandas. Multifractal analysis of the irregular set for almost-additive sequences via large deviations. *Preprint*, 2014, arXiv:1410.2220.
[**5**] R. Bowen. Some systems with unique equilibrium states. *Math. Syst. Theory* **8** (1974), 193–202.
[**6**] R. Bowen. *Equilibrium States and the Ergodic Theory of Anosov Diffeomorphisms (Lecture Notes in Mathematics, 470)*. Springer, Berlin, 1975.
[**7**] R. Bowen and D. Ruelle. The ergodic theory of Axiom A flows. *Invent. Math.* **29**(3) (1975), 181–202.
[**8**] H. Bruin and M. Todd. Equilibrium states for interval maps: the potential $-t\log|Df|$. *Ann. Sci. Éc. Norm. Supér.* (4) **42** (2009), 559–600.
[**9**] A. Castro and P. Varandas. Equilibrium states for non-uniformly expanding maps: decay of correlations and strong stability. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **30**(2) (2013), 225–249.
[**10**] A. Castro and T. Nascimento. Statistical properties of equilibrium states for partially hyperbolic attractors. *Preprint*, UFBA, 2014.
[**11**] V. Climenhaga. Multifractal formalism derived from thermodynamics for general dynamical systems. *Electron. Res. Announc. Math. Sci.* **17** (2010), 1–11.
[**12**] V. Climenhaga. Topological pressure of simultaneous level sets. *Nonlinearity* **26** (2013), 241–268.
[**13**] V. Climenhaga, D. Thompson and K. Yamamoto. Large deviations for systems with non-uniform structure. *Preprint*, 2013, arXiv:1304.5497.
[**14**] H. Comman and J. Rivera-Letelier. Large deviations principles for non-uniformly hyperbolic rational maps. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **15** (1998), 539–579.
[**15**] M. Denker and M. Kesseböhmer. Thermodynamical formalism, large deviation and multifractals. *Stochastic Climate Models (Progress in Probability, 49)*. Birkhäuser, Basel, 2001, pp. 159–169.
[**16**] L. Díaz and K. Gelfert. Porcupine-like horseshoes: transitivity, Lyapunov spectrum, and phase transitions. *Fund. Math.* **216**(1) (2012), 55–100.
[**17**] L. Díaz, K. Gelfert and M. Rams. Rich phase transitions in step skew products. *Nonlinearity* **24**(12) (2011), 3391–3412.
[**18**] A. Eizenberg, Y. Kifer and B. Weiss. Large deviations for $\mathbb{Z}^d$ actions. *Comm. Math. Phys.* **164** (1994), 433–454.
[**19**] D. Feng and W. Huang. Lyapunov spectrum of asymptotically sub-additive potentials. *Comm. Math. Phys.* **297** (2010), 1–43.
[**20**] K. Gelfert and M. Rams. The Lyapunov spectrum of some parabolic systems. *Ergod. Th. & Dynam. Sys.* **29**(3) (2009), 919–940.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press

[**21**] F. Hofbauer. Examples for the nonuniqueness of the equilibrium state. *Trans. Amer. Math. Soc.* **228** (1977), 223–241.
[**22**] T. Jordan and M. Rams. Multifractal analysis of weak Gibbs measures for non-uniformly expanding C^1 maps. *Ergod. Th. & Dynam. Sys.* **31**(1) (2011), 143–164.
[**23**] V. Kleptsyn, D. Ryzhov and S. Minkov. Special ergodic theorems and dynamical large deviations. *Nonlinearity* **25** (2012), 3189–3196.
[**24**] I. Melbourne. Large and moderate deviations for slowly mixing dynamical systems. *Proc. Amer. Math. Soc.* **137** (2009), 1735–1741.
[**25**] I. Melbourne and M. Nicol. Large deviations for nonuniformly hyperbolic systems. *Trans. Amer. Math. Soc.* **360** (2008), 6661–6676.
[**26**] Y. Pesin. *Dimension Theory in Dynamical Systems (Contemporary Views and Applications)*. University of Chicago Press, Chicago, IL, 1997.
[**27**] Y. Pesin and H. Weiss. The multifractal analysis of Gibbs measures: motivation, mathematical foundation, and examples. *Chaos* **7**(1) (1997), 89–106.
[**28**] C. Pfister and W. Sullivan. Large deviations estimates for dynamical systems without the specification property. Applications to the β-shifts. *Nonlinearity* **18** (2005), 237–261.
[**29**] M. Pollicott and R. Sharp. Large deviations for intermittent maps. *Nonlinearity* **22** (2009), 2079–2092.
[**30**] M. Pollicott and H. Weiss. Multifractal analysis of Lyapunov exponent for continued fraction and Manneville–Pomeau transformations and applications to Diophantine approximation. *Comm. Math. Phys.* **207** (1999), 145–171.
[**31**] L. Rey-Bellet and L.-S. Young. Large deviations in non-uniformly hyperbolic dynamical systems. *Ergod. Th. & Dynam. Sys.* **28** (2008), 587–612.
[**32**] F. Takens and E. Verbitski. Multifractal analysis of local entropies for expansive homeomorphisms with specification. *Comm. Math. Phys.* **203** (1999), 593–612.
[**33**] D. Thompson. A variational principle for topological pressure for certain non-compact sets. *J. Lond. Math. Soc.* **80**(3) (2009), 585–602.
[**34**] M. Todd. Multifractal analysis for multimodal maps. *Preprint*, 2008.
[**35**] D. Thompson. The irregular set for maps with the specification property has full topological pressure. *Dyn. Syst.* **25**(1) (2010), 25–51.
[**36**] P. Varandas. Non-uniform specification and large deviations for weak Gibbs measures. *J. Stat. Phys.* **146** (2012), 330–358.
[**37**] P. Varandas and M. Viana. Existence, uniqueness and stability of equilibrium states for non-uniformly expanding maps. *Ann. Inst. H. Poincaré Anal. Non Linéaire* **27** (2010), 555–593.
[**38**] L.-S. Young. Some large deviations for dynamical systems. *Trans. Amer. Math. Soc.* **318** (1990), 525–543.
[**39**] X. Zhou and E. Chen. Multifractal analysis for the historic set in topological dynamical systems. *Nonlinearity* **26**(7) (2013), 1975–1997.

https://doi.org/10.1017/etds.2015.46 Published online by Cambridge University Press