Combinatorics, Probability and Computing (1997) 6, 263–271. Printed in the United Kingdom
© 1997 Cambridge University Press

Dependent Sets of Constant Weight Binary Vectors

NEIL J. CALKIN

School of Mathematics, Georgia Institute of Technology,
Atlanta, GA 30332-0160, USA

Received 11 April 1995; revised 20 February 1996

We determine lower bounds for the number of random binary vectors, chosen uniformly from vectors of weight k, needed to obtain a dependent set.

1. Introduction

In this paper we determine lower bounds for the number of random binary vectors of weight k needed to obtain a dependent set of vectors with probability 1.

We denote by $S_{n,k}$ the set of binary vectors having k 1's. If we choose a random sequence $\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_m$ uniformly from $S_{n,k}$, how large must m be for these vectors to be dependent (over GF(2)) with probability 1?

In the case $k = 1$ this is exactly the birthday problem: given a set of n elements, how long must a sequence chosen (with replacement) be before an element occurs at least twice with probability close to 1? It is a standard combinatorics exercise to show that so long as $m/\sqrt{n} \to \infty$, a sequence of length m will almost surely contain a repetition as $n \to \infty$.

In the case $k = 2$, we can view the vectors of weight two as being edges in a graph on $\{1, 2, \ldots, n\}$: here a dependent set of vectors corresponds exactly to a set of edges which contain a cycle. There are two distinct modes of behaviour here: first, if the edges are chosen without replacement, and if the number of edges is cn, then the probability that there is a cycle is strictly less than 1 as $n \to \infty$ if $c < 1/2$ and tends to 1 if $c \geqslant 1/2$ [2]. If the edges are chosen with replacement, then if we choose cn edges, there is a positive probability that we get a repeated edge. Hence the probability increases up to $c = 1/2$, at which point we almost surely get a cycle.

In what follows, we will assume that k is a fixed integer greater than or equal to 3. Denote by $p_{n,k}(m)$ the probability that $\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_m$ are linearly dependent. We will prove the following:

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

Theorem 1.1. *For each k there is a constant β_k so that if $\beta < \beta_k$ then*

$$\lim_{n\to\infty} p_{n,k}(\beta n) = 0.$$

Furthermore, $\beta_k \sim 1 - e^{-k}/\log 2$ as $k \to \infty$.

We obtain this theorem as a corollary of the following: let r be the rank of the set $\{\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_m\}$, and let $s = m - r$ (equivalently, the dimension of the kernel of the matrix having columns $\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_m$).

Theorem 1.2. (a) *If $\beta < \beta_k$ and $m = m(n) < \beta n$ then $E(2^s) \to 1$ as $n \to \infty$.*
(b) *If $\beta > \beta_k$ and $m = m(n) > \beta n$ then $E(2^s) \to \infty$ as $n \to \infty$.*

Similar results have been obtained for different models by Balakin, Kolchin and Khokhlov [1, 3]: their methods are completely different.

Our approach is the following: we consider a Markov chain derived from a suitable random walk on the hypercube 2^n; using this we will determine an exact expression for $E(2^s)$. We then estimate $E(2^s)$ to determine β_k.

2. A random walk on the hypercube, and an associated Markov chain

We define a random walk on the hypercube 2^n as follows: let $\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_m, \ldots$ be vectors chosen uniformly at random from $S_{n,k}$. Define

$$\underline{x}_0 = \underline{0}, \quad \text{and} \quad \underline{x}_i = \underline{x}_{i-1} + \underline{u}_i$$

(so the steps in the walk correspond to flipping k random bits).

We associate with this random walk the following Markov chain: we define y_i to be the weight of $\underline{x}_1$. Then $y_0, y_1, \ldots, y_m$, is a Markov chain with states $\{0, 1, \ldots, n\}$. The transition matrix A for this chain, with $A = \{a_{pq}\}$, where a_{pq} is the probability of moving from state q to state p is given by

$$a_{pq} = \binom{q}{\frac{k-p+q}{2}} \binom{n-q}{\frac{k+p-q}{2}} \Big/ \binom{n}{k}$$

where the binomial coefficients are interpreted to be 0 if $k + p + q$ is odd.

Theorem 2.1. *The eigenvalues λ_i and corresponding eigenvectors $\underline{e}_i$ for A, $i = 0, 1, \ldots, n$, are given by*

$$\lambda_i = \sum_{t=0}^{k} (-1)^t \frac{\binom{i}{t}\binom{n-i}{k-t}}{\binom{n}{k}} \tag{1}$$

and the jth component of $\underline{e}_i$ is given by

$$\underline{e}_i[j] = \sum_{t=0}^{j} (-1)^t \binom{i}{t}\binom{n-i}{j-t}.$$

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

Proof. We first show that $\underline{e}_i$ is an eigenvector for A with eigenvalue λ_i: indeed the jth coefficient of $A\underline{e}_i$ is

$$\sum_{l=0}^{n} \frac{\binom{l}{\frac{k-j+l}{2}}\binom{n-l}{\frac{k+j-l}{2}}}{\binom{n}{k}} \sum_{t=0}^{l}(-1)^t \binom{i}{t}\binom{n-i}{l-t}$$

and the jth coefficient of $\lambda_i \underline{e}_i$ is

$$\sum_{s=0}^{k}(-1)^s \frac{\binom{i}{s}\binom{n-i}{k-s}}{\binom{n}{k}} \sum_{t=0}^{j}(-1)^t \binom{i}{t}\binom{n-i}{j-t}.$$

Observe now that

$$\sum_{t=0}^{j}(-1)^t \binom{i}{t}\binom{n-i}{j-t} = \sum_{t=0}^{i}(-1)^{i+t}2^t \binom{i}{t}\binom{n-t}{j},$$

since each is the coefficient of x^j in

$$(1-x)^i(1+x)^{n-i} = \left(1-\frac{2}{1+x}\right)^i (1+x)^n.$$

Hence it is sufficient to show that

$$\sum_{l=0}^{n}\binom{l}{\frac{k-j+l}{2}}\binom{n-l}{\frac{k+j-l}{2}} \sum_{t=0}^{j}(-1)^{t+i}2^t \binom{i}{t}\binom{n-t}{j}$$

$$= \sum_{s=0}^{k}(-1)^s \binom{i}{s}\binom{n-i}{k-s} \sum_{t=0}^{j}(-1)^t \binom{i}{t}\binom{n-i}{j-t}.$$

We show this by multiplying both sides by $x^j y^k$ and summing over j and k. Writing $j = l - 2r + k$, the left hand side becomes

$$\begin{aligned}
&\sum_{l,k,r,t}\binom{l}{r}\binom{n-l}{k-r}\binom{i}{t}\binom{n-t}{l}(-1)^{i+t}2^t x^{l+k-2r}y^k \\
&\quad = \sum_{l,r,t}\binom{l}{r}\binom{i}{t}\binom{n-t}{l}(-1)^{i+t}2^t x^{l-r}(1+xy)^{n-l}y^r \\
&\quad = \sum_{l,t}\binom{i}{t}\binom{n-t}{l}(-1)^{i+t}2^t(1+xy)^{n-l}(x+y)^l \\
&\quad = \sum_{t}(-1)^{i+t}\binom{i}{t}2^t(1+xy)^t(1+x)^{n-t}(1+y)^{n-t} \\
&\quad = (1+x)^n(1+y)^n\left(\frac{2(1+xy)}{(1+x)(1+y)}-1\right)^i \\
&\quad = (1-x)^i(1+x)^{n-i}(1-y)^i(1+y)^{n-i}.
\end{aligned}$$

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

Similarly, the right-hand side becomes

$$\begin{aligned}
&\sum_{j,k,s,t} \binom{i}{s}\binom{n-i}{k-s}\binom{i}{t}\binom{n-i}{j-t}(-1)^{s+t}x^j y^k \\
&\quad = \sum_{s,t} \binom{i}{s}\binom{i}{t}(-1)^{s+t}x^t y^s (1+x)^{n-i}(1+y)^{n-i} \\
&\quad = (1-x)^i(1+x)^{n-i}(1-y)^i(1+y)^{n-i}
\end{aligned}$$

as required. Hence $\underline{e}_i$ is an eigenvector with eigenvalue λ_i for each i. □

Moreover, we see that the $\underline{e}_i$'s are linearly independent (as vectors over Q): indeed, we have the following lemma.

Lemma 2.1. *Let U be the matrix whose columns are $\underline{e}_0, \underline{e}_1, \ldots, \underline{e}_n$. Then $U^2 = 2^n I$, and if Λ is the diagonal matrix of eigenvalues, then $A = 1/2^n U\Lambda U$.*

Proof. The ijth entry of U^2 is

$$\sum_{l=0}^{n} \underline{e}_l[i]\underline{e}_j[l] = \sum_{l,s,t}(-1)^s\binom{l}{s}\binom{n-l}{i-s}(-1)^t\binom{j}{t}\binom{n-j}{l-t}.$$

Multiplying by x^i and summing over i we obtain

$$\begin{aligned}
&\sum_{i,l,s,t}(-1)^{s+t}\binom{l}{s}\binom{n-l}{i-s}\binom{j}{t}\binom{n-j}{l-t}x^i \\
&\quad = \sum_{l,s,t}(-1)^{s+t}\binom{l}{s}\binom{j}{t}\binom{n-j}{l-t}(1+x)^{n-l}x^s \\
&\quad = \sum_{l,t}(-1)^t\binom{j}{t}\binom{n-j}{l-t}(1+x)^{n-l}(1-x)^l \\
&\quad = \sum_{t}(-1)^t\binom{j}{t}(1+x)^j 2^{n-j}\left(\frac{1-x}{1+x}\right)^t \\
&\quad = 2^n x^j,
\end{aligned}$$

from which we see that $U^2 = 2^n I$. Hence the eigenvectors are linearly independent as claimed. □

Observation. The eigenvectors do not depend upon k: hence the matrices A and A' corresponding to distinct values of k commute. This corresponds roughly to the idea that when walking around the hypercube it doesn't matter if you take a step of size l then a step of size k, or a step of size k then a step of size l.

We can now compute the probability that $\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_t$ sum to $\underline{0}$: indeed, this is exactly

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

the 00th coefficient in A^t, which is equal to

$$\sum_{i=0}^{n} \frac{1}{2^n} \lambda_i^t \binom{n}{i}$$

(since $A = 1/2^n U \Lambda U$).

Hence if $\underline{u}_1, \underline{u}_2, \ldots, \underline{u}_m$ are vectors with k 1's chosen independently at random, then the expected number of subsequences $\underline{u}_{a_1}, \underline{u}_{a_2}, \ldots, \underline{u}_{a_t}$ which sum to $\underline{0}$ is exactly

$$E(2^s) = \sum_{t=0}^{m} \binom{m}{t} \sum_{i=0}^{n} \frac{1}{2^n} \lambda_i^t \binom{n}{i} = \sum_{i=0}^{n} \frac{1}{2^n} \binom{n}{i} (1 + \lambda_i)^m .$$

3. Asymptotics of λ_i

To estimate the size of $E(2^s)$, we require asymptotics for the value of λ_i.

Lemma 3.1. (a) $|\lambda_i| < 1$ *for all* $0 \leqslant i \leqslant n$.
(b) *If* $i > \frac{n}{2}$ *then* $\lambda_i = (-1)^k \lambda_{n-i}$.
(c) *Let* $0 < c < \frac{1}{2}$. *If* $i = cn$ *then*

$$\lambda_i = \left(1 - \frac{2i}{n}\right)^k - \frac{4\binom{k}{2}}{n} \left(1 - \frac{2i}{n}\right)^{k-2} \frac{i}{n} \left(1 - \frac{i}{n}\right) + O\left(\frac{k^3}{c^2 n^2}\right)$$

Proof. Parts (a) and (b) are immediate from the definition of λ_i. To prove part (c), since k is fixed, we have

$$\binom{n}{k} = \frac{n^k}{k!} \left(1 - \frac{\binom{k}{2}}{n} + O\left(\frac{k^3}{n^2}\right)\right)$$

$$\binom{i}{t} = \frac{i^t}{t!} \left(1 - \frac{\binom{t}{2}}{i} + O\left(\frac{t^3}{i^2}\right)\right)$$

$$\binom{n-i}{k-t} = \frac{(n-i)^{k-t}}{(k-t)!} \left(1 - \frac{\binom{k-t}{2}}{n-i} + O\left(\frac{(k-t)^3}{(n-i)^2}\right)\right).$$

Hence

$$\frac{\binom{i}{t}\binom{n-i}{k-t}}{\binom{n}{k}} = \left(\frac{i}{n}\right)^t \left(1 - \frac{i}{n}\right)^{k-t} \binom{k}{t} \left(1 + \frac{\binom{k}{2}}{n} - \frac{\binom{t}{2}}{i} - \frac{\binom{k-t}{2}}{n-i} + O\left(\frac{k^3}{c^2 n^2}\right)\right)$$

and

$$\begin{aligned}
\lambda_i &= \sum_{t=0}^{k} (-1)^t \left(\frac{i}{n}\right)^t \left(1 - \frac{i}{n}\right)^{k-t} \binom{k}{t} \left(1 + \frac{\binom{k}{2}}{n} - \frac{\binom{t}{2}}{i} - \frac{\binom{k-t}{2}}{n-i} + O\left(\frac{k^3}{c^2 n^2}\right)\right) \\
&= \left(1 - \frac{2i}{n}\right)^k + \frac{\binom{k}{2}}{n} \left(1 - \frac{2i}{n}\right)^k - \frac{\binom{k}{2}}{i} \left(\frac{i}{n}\right)^2 \left(1 - \frac{2i}{n}\right)^{k-2} \\
&\quad - \frac{\binom{k}{2}}{n-i} \left(\frac{n-i}{n}\right)^2 \left(1 - \frac{2i}{n}\right)^{k-2} + O\left(\frac{k^3}{c^2 n^2}\right)
\end{aligned}$$

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

$$= \left(1-\frac{2i}{n}\right)^k + \frac{\binom{k}{2}}{n}\left(1-\frac{2i}{n}\right)^{k-2}\left(\left(1-\frac{2i}{n}\right)^2 - \frac{i}{n} - \frac{n-i}{n}\right) + O\left(\frac{k^3}{c^2n^2}\right)$$
$$= \left(1-\frac{2i}{n}\right)^k - \frac{4\binom{k}{2}}{n}\left(1-\frac{2i}{n}\right)^{k-2}\left(-\frac{4i}{n} + 4\frac{i^2}{n}\right) + O\left(\frac{k^3}{c^2n^2}\right)$$
$$= \left(1-\frac{2i}{n}\right)^k - \frac{4\binom{k}{2}}{n}\left(1-\frac{2i}{n}\right)^{k-2}\left(\frac{i}{n}\right)\left(1-\frac{i}{n}\right) + O\left(\frac{k^3}{c^2n^2}\right)$$

as claimed. □

Observe that, since we are assuming that $k \geqslant 3$ throughout, when i is close to $\frac{n}{2}$, say $\frac{n}{2} - i = \frac{n^\theta}{2}$, we have

$$\lambda_i = \left(\frac{1}{n^{1-\theta}}\right)^k - \frac{4\binom{k}{2}}{n}\left(\frac{1}{n^{1-\theta}}\right)^{k-2} + O\left(\frac{k^3}{n^2}\right)$$

Then, provided that $\theta < 1 - \frac{1}{k}$, we see that if $\frac{n}{2} - i = \frac{n^\theta}{2}$, then $\lambda_i n \to 0$ as $n \to \infty$. In the estimation of $E(2^s)$ we will use this to show that the middle part of the sum is asymptotic to 1.

4. Asymptotics of $E(2^s)$

Define

$$f(\alpha,\beta) = -\log 2 - \alpha\log(\alpha) - (1-\alpha)\log(1-\alpha) + \beta\log(1+(1-2\alpha)^k)$$

and let (α_k, β_k) be the root of

$$f(\alpha,\beta) = 0$$
$$\frac{\partial f(\alpha,\beta)}{\partial \alpha} = 0.$$

We shall show:

Lemma 4.1. *If $\beta < \beta_k$ and $m < \beta n$ then $\sum_i 2^{-n}\binom{n}{i}(1+\lambda_i)^m \to 1$ as $n \to \infty$, and if $\beta > \beta_k$ and $m > \beta n$ then $\sum_i 2^{-n}\binom{n}{i}(1+\lambda_i)^m \to \infty$ as $n \to \infty$.*

Proof. We proceed as follows: since our goal is to show that the behaviour of $E(2^s)$ changes when m goes from below $\beta_k n$ to above $\beta_k n$, and since our value β_k is less than 1, we may assume that $\frac{m}{n} < 1 - \delta$ for some $\delta > 0$. We shall show:

1. The extreme tails of the sum for $E(2^s)$ are small.
2. The middle range of the sum contributes 1 to the sum.
3. The rest of the sum is small if $\frac{m}{n} < \beta < \beta_k$.
4. The rest of the sum is large if $\frac{m}{n} > \beta > \beta_k$.

1. There is an $\epsilon > 0$ so that

$$\sum_{i=0}^{\epsilon n} 2^{-n}\binom{n}{i}(1+\lambda_i)^m \to 0 \text{ as } n \to \infty$$

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

Indeed,

$$\sum_{i=0}^{\epsilon n} 2^{-n}\binom{n}{i}(1+\lambda_i)^m < \sum_{i=0}^{\epsilon n} 2^{m-n}\binom{n}{i} < n\epsilon 2^{m-n}\binom{n}{\epsilon n}$$

and provided ϵ is sufficiently small, this tends to 0 (indeed, if $-\delta\log 2 - \epsilon\log\epsilon + \epsilon < 0$ then the sum tends to 0).

Similarly,

$$\sum_{i=(1-\epsilon)n}^{n} 2^{-n}\binom{n}{i}(1+\lambda_i)^m \to 0 \text{ as } n\to\infty.$$

Hence, if $E(2^s)\to\infty$ for some $m < (1-\delta)n$, we must have the major contribution from

$$\sum_{i=\epsilon n}^{(1-\epsilon)n} 2^{-n}\binom{n}{i}(1+\lambda_i)^m.$$

2. We now show that the middle range of the sum contributes 1 to $E(2^s)$. Indeed, in the range $\frac{n}{2} - n^{4/7} < i < \frac{n}{2} + n^{4/7}$

$$(1+\lambda_i)^m = \left(1 + o\left(\frac{1}{n}\right)\right)^m = 1 + o(1)$$

we have

$$\sum_{i=\frac{n}{2}-n^{4/7}}^{\frac{n}{2}+n^{4/7}} 2^{-n}\binom{n}{i}(1+\lambda_i)^m \sim \sum_{1=\frac{n}{2}-n^{4/7}}^{\frac{n}{2}+n^{4/7}} 2^{-n}\binom{n}{i} \to 1 \text{ as } n\to\infty.$$

3. We now show that we can widen the interval about the middle:

$$\sum_{i=\frac{n}{2}(1-\epsilon)}^{\frac{n}{2}(1+\epsilon)} 2^{-n}\binom{n}{i}(1+\lambda_i)^m \to 1.$$

Since $\lambda_{n-i} = (-1)^k\lambda_i$, it suffices to show that

$$\sum_{i=\frac{n}{2}(1-\epsilon)}^{\frac{n}{2}-n^{4/7}} 2^{-n}\binom{n}{i}(1+\lambda_i)^m \to 0.$$

In this range,

$$\lambda_i < \epsilon^k - \frac{\binom{k}{2}}{n}\epsilon^{k-2} + O\left(\frac{k^3}{n^2}\right).$$

Hence

$$(1+\lambda_i)^m < e^{n\epsilon^k} e^{-\binom{k}{2}\epsilon^{k-2}},$$

and since $k \geqslant 3$, the $n\epsilon^k$ term in the exponent is dominated by the $-n\epsilon^2$ term from the binomial coefficient, provided that ϵ is sufficiently small.

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

4. We now consider the remainder of the sum (or rather, the part in $(0, \frac{n}{2})$: if k is even, the remaining part follows by symmetry, and if k is odd, then $(1+\lambda_i)^m < 1$ for $i > n/2$, and the remaining part tends to 0).

Define

$$f(\alpha,\beta) = -\log 2 - \alpha\log\alpha - (1-\alpha)\log(1-\alpha) + \beta\log(1+(1-2\alpha)^k).$$

Then if $f(\frac{i}{n},\frac{m}{n}) < \gamma < 0$ the corresponding term of the sum is exponentially small, and if $f(\frac{i}{n},\frac{m}{n}) > \gamma > 0$ the corresponding term of the sum is exponentially large. Thus, if $f(\alpha,\frac{m}{n}) < \gamma < 0$ for all α in $(\epsilon, 1-\epsilon)$, we have

$$\sum_{i=\epsilon n}^{\frac{n}{2}(1-\epsilon)} 2^{-n}\binom{n}{i}(1+\lambda_i)^m < ne^{\gamma n+o(n)} \to 0,$$

and if $f(\alpha,\frac{m}{n}) > \gamma > 0$ for some α in $(\epsilon, 1-\epsilon)$, then

$$\sum_{i=\epsilon n}^{\frac{n}{2}(1-\epsilon)} 2^{-n}\binom{n}{i}(1+\lambda_i)^m > \binom{n}{\alpha n}(1+\lambda_{\alpha n})^m 2^{-n}e^{\gamma n+o(n)} \to \infty.$$

Now let β_k be so that if $\beta < \beta_k$ then $f(\alpha,\beta) < 0$ for all α in $(\epsilon, 1-\epsilon)$, and if $\beta > \beta_k$ then there is an alpha in $(\epsilon, 1-\epsilon)$ so that $f(\alpha,\beta) > 0$. Thus we wish to find α_k, β_k so that

$$f(\alpha_k,\beta_k) = 0 \text{ and } \frac{\partial}{\partial\alpha}f(\alpha,\beta) = 0.$$

As k goes to ∞, the value of β_k is asymptotic to

$$1 - \frac{e^{-k}}{\log 2} - \frac{1}{2\log 2}\left(k^2 - 2k + \frac{2k}{\log 2} - 1\right)e^{-2k} + O(k^4)e^{-3k}.$$

To see this, we observe first that

$$\alpha_k = e^{-k}$$

and

$$\beta_k = 1 - \frac{e^{-k}}{\log 2}$$

are close to a root (by considering a small constant times the error term, and expanding out $f(\alpha,\beta)$, we see that both $f(\alpha,\beta)$ and $\frac{\partial}{\partial\alpha}f(\alpha,\beta)$ change sign as the constant changes). Furthermore, there are no other roots α in (0,1/2) and β in (0,1): indeed, we note

1 That for any root of the two equations, indeed, for any root of $f(\alpha,\beta)$ with $\alpha \in (0, \frac{1}{2})$, either $\alpha \to 0$ as $k \to \infty$ or $\alpha \to \frac{1}{2}$ as $k \to \infty$.
2 If α is close to $\frac{1}{2}$, then there it is not part of a root (by expanding out $f(\alpha,\beta)$ in terms of α and observing that the term involving β is of order α^k, and since $k \geqslant 3$ and $\beta \leqslant 1$, we cannot have a root.
3 By considering the expansion in α, we observe that $k\alpha \to 0$ as $k \to \infty$.
4 Now, by expanding out both $f(\alpha,\beta)$ and $\frac{\partial}{\partial\alpha}f(\alpha,\beta)$, we see that the root is as claimed. Using a symbolic algebra package (in our case Maple), it is easy now to see that β_k

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press

has an asymptotic expansion

$$\beta_k \sim 1 - \frac{e^{-k}}{\log 2} - \frac{1}{2\log 2}(k^2 - 2k + \frac{2k}{\log 2} - 1)e^{-2k} + O(k^4)e^{-3k}$$

as k goes to infinity.

□

Now, since $E(2^s) = \sum_i 2^{-n}\binom{n}{i}(1+\lambda_i)^m$ this completes the proof of Theorem 1.2, and Theorem 1.1 follows by the simple observation that since s is integer valued, the probability that $2^s > 1$ is less than $E(2^s) - 1$.

References

[1] Balakin, G. V., Kolchin, V. F. and Khokhlov, V. I. (1991) Hypercycles in a random hypergraph. *Diskretnaya Matematika* **3**(3) 1-2–108 (in Russian).

[2] Erdős, P. and Rényi, A. (1960) On the evolution of random graphs. *Publ. Math. Inst. Hungar. Acad. Sci.* **5**17–61.

[3] Kolchin, V. F. and Khokhlov, V. I. (1990) On the number of cycles in a non-equiprobably random graph. *Diskretnaya Matematika* **2**(3) 137–145 (in Russian).

https://doi.org/10.1017/S0963548397003040 Published online by Cambridge University Press