Ergod. Th. & Dynam. Sys. (2015), **35**, 910–914 doi:10.1017/etds.2013.73
© *Cambridge University Press, 2013*

Bohr density of simple linear group orbits

ROGER HOWE† and FRANÇOIS ZIEGLER‡

† *Department of Mathematics, Yale University, New Haven, CT 06520-8283, USA*
(e-mail: howe@math.yale.edu)

‡ *Department of Mathematical Sciences, Georgia Southern University, Statesboro, GA 30460-8093, USA*
(e-mail: fziegler@georgiasouthern.edu)

(*Received* 15 *November* 2012 *and accepted in revised form* 26 *July* 2013)

Abstract. We show that any non-zero orbit under a non-compact, simple, irreducible linear group is dense in the Bohr compactification of the ambient space.

1. *Introduction*

Let V be a locally compact abelian group, V^* its Pontryagin dual and bV its Bohr compactification, that is, bV is the dual of the discretized group V^*. On identifying V with its double dual we have a dense embedding $V \hookrightarrow bV$, namely,

$$\{\text{continuous characters of } V^*\} \hookrightarrow \{\text{all characters of } V^*\}.$$

The relative topology of V in bV is known as the *Bohr topology* of V. Among its many intriguing properties (surveyed in [**G07**]) is the observation due to Katznelson [**K73a**] (see also [**G79**, §7.6]) that very 'thin' subsets of V can be Bohr dense in very large ones.

While Katznelson was concerned with the case $V = \mathbf{Z}$ (the integers), we shall illustrate this phenomenon in the setting where V is the additive group of a real vector space, and the subsets of interest are the orbits of a Lie group acting linearly on V. Indeed our aim is to establish the following result, which was conjectured in [**Z96**, p. 45].

THEOREM 1. *Let G be a non-compact, simple real Lie group and V a non-trivial, irreducible, finite-dimensional real G-module. Then every non-zero G-orbit in V is dense in bV.*

We prove this in §3 on the basis of four lemmas found in §2. Before that, let us record a similar property of *nilpotent* groups. In that case, orbits typically lie in proper affine subspaces, so we cannot hope for Bohr density in the whole space; but we have the following theorem.

THEOREM 2. *Let G be a connected nilpotent Lie group and V a finite-dimensional G-module of unipotent type. Then every G-orbit in V is Bohr dense in its affine hull.*

https://doi.org/10.1017/etds.2013.73 Published online by Cambridge University Press

Proof. Recall that *unipotent type* means that the Lie algebra $\mathfrak{g}$ of G acts by nilpotent operators. So $Z \mapsto \exp(Z)v$ is a polynomial map of $\mathfrak{g}$ onto the orbit of $v \in V$, and the claim follows immediately from [**Z93**, Theorem]. □

2. *Four lemmas*

Our first lemma gives several characterizations of Bohr density—each of which can also be regarded as providing a corollary of Theorem 1.

LEMMA 1. *Let $\mathcal{O}$ be a subset of the locally compact abelian group V. Then the following are equivalent:*

(1) *$\mathcal{O}$ is dense in bV;*
(2) *$\alpha(\mathcal{O})$ is dense in $\alpha(V)$ whenever α is a continuous morphism from V to a compact topological group;*
(3) *every almost periodic function on V is determined by its restriction to $\mathcal{O}$;*
(4) *Haar measure η on bV is the weak$*$ limit of probability measures μ_T concentrated on $\mathcal{O}$.*

Proof. (1) ⇔ (2): Clearly (2) implies (1) as the special case where α is the natural inclusion $\iota : V \hookrightarrow bV$. Conversely, suppose (1) holds and $\alpha : V \to X$ is a continuous morphism to a compact group. By the universal property of bV [**D82**, Theorem 16.1.1], $\alpha = \beta \circ \iota$ for a continuous morphism $\beta : bV \to X$. Now continuity of β implies $\beta(\overline{\iota(\mathcal{O})}) \subset \overline{\beta(\iota(\mathcal{O}))}$, which is to say that $\beta(bV) \subset \overline{\alpha(\mathcal{O})}$ and hence $\alpha(V) \subset \overline{\alpha(\mathcal{O})}$, as claimed.

(1) ⇔ (3): Recall that a function on V is *almost periodic* if and only if it is the pull-back of a continuous $f : bV \to \mathbf{C}$ by the inclusion $V \hookrightarrow bV$. If two such functions coincide on $\mathcal{O}$ and $\mathcal{O}$ is dense in bV, then clearly they coincide everywhere. Conversely, suppose that $\mathcal{O}$ is not dense in bV. Then by complete regularity [**H63**, Theorem 8.4] there is a non-zero continuous $f : bV \to \mathbf{C}$ which is zero on the closure of $\mathcal{O}$ in bV. Now clearly this f is not determined by its restriction to $\mathcal{O}$.

(1) ⇔ (4) [**K73a**]: Suppose that η is the weak$*$ limit of probability measures μ_T concentrated on $\mathcal{O}$. So we have $\mu_T(f) \to \eta(f)$ for every continuous f, and the complement of $\mathcal{O}$ in bV is μ_T-null [**B04**, Definition V.5.7.4 and Proposition IV.5.2.5]. If f vanishes on the closure of $\mathcal{O}$ in bV then so do all $\mu_T(|f|)$ and hence also $\eta(|f|)$, which forces f to vanish everywhere. So $\mathcal{O}$ is dense in bV. Conversely, suppose that $\mathcal{O}$ is dense in bV. We have to show that given continuous functions $f_1, \ldots, f_n$ on bV and $\varepsilon > 0$, there is a probability measure μ concentrated on $\mathcal{O}$ such that $|\eta(f_j) - \mu(f_j)| < \varepsilon$ for all j. Writing

$$F = (f_1, \ldots, f_n) \quad \text{and} \quad \eta(F) = (\eta(f_1), \ldots, \eta(f_n)),$$

we see that this amounts to $\|\eta(F) - \mu(F)\| < \varepsilon$, where the norm is the sup norm in $\mathbf{C}^n$. Now by [**B04**, Corollary V.6.1] $\eta(F)$ lies in the convex hull of $F(bV)$ (which is compact by Carathéodory's theorem [**B87**, Corollary 11.1.8.7]). So $\eta(F)$ is a convex combination $\sum_{i=1}^{N} \lambda_i F(\omega_i)$ of elements of $F(bV)$. But $F(\mathcal{O})$ is dense in $F(bV)$, so we can find $w_i \in \mathcal{O}$ such that $\|F(\omega_i) - F(w_i)\| < \varepsilon$. Putting $\mu = \sum_{i=1}^{N} \lambda_i \delta_{w_i}$, where δ_{w_i} is Dirac measure at w_i, we obtain the desired probability measure μ. □

https://doi.org/10.1017/etds.2013.73 Published online by Cambridge University Press

Remark 1. One might wonder if condition (2) is equivalent to the following *a priori* weaker but already interesting property:

(2′) $\mathcal{O}$ has dense image in any compact quotient group of V.

Here is an example showing that (2′) *does not* imply (2). Let $V = \mathbf{R}$ and $\mathcal{O} = \mathbf{Z} \cup 2\pi\mathbf{Z}$. Then clearly $\mathcal{O}$ has dense image in every compact quotient $\mathbf{R}/a\mathbf{Z}$. On the other hand, considering the irrational winding $\alpha : \mathbf{R} \to \mathbf{T}^2$ defined by $\alpha(v) = (e^{iv}, e^{2\pi i v})$, one can check without difficulty that $\overline{\alpha(\mathcal{O})} = \mathbf{T} \times \{1\} \cup \{1\} \times \mathbf{T}$, which is strictly smaller than $\overline{\alpha(V)} = \mathbf{T}^2$.

Remark 2. A net of probability measures μ_T converging to Haar measure on bV as in (4) has been called a *generalized summing sequence* by Blum and Eisenberg [**B74**]. They observed, among others, the following characterization.

LEMMA 2. *The following conditions are equivalent:*

(1) *μ_T is a generalized summing sequence;*

(2) *the Fourier transforms $\hat{\mu}_T(u) = \int_{bV} \omega(u)\, d\mu_T(\omega)$ converge pointwise to the characteristic function of $\{0\} \subset V^*$.*

Proof. This characteristic function is the Fourier transform of Haar measure η on bV. Thus, condition (2) says that $\mu_T(f) \to \eta(f)$ for every continuous character $f(\omega) = \omega(u)$ of bV, whereas condition (1) says that $\mu_T(f) \to \eta(f)$ holds for every continuous function f on bV. Since linear combinations of continuous characters are uniformly dense in the continuous functions on bV (Stone–Weierstrass), the two conditions imply each other. □

For our third lemma, let G be a group, V a finite-dimensional G-module, and write V^* for the dual module wherein G acts contragrediently: $\langle gu, v\rangle = \langle u, g^{-1}v\rangle$.

LEMMA 3. *Suppose that V is irreducible and $\phi(g) = \langle u, gv\rangle$ is a non-zero matrix coefficient of V. Then every other matrix coefficient $\psi(g) = \langle x, gy\rangle$ is a linear combination of left and right translates of ϕ.*

Proof. Irreducibility of V and (therefore) V^* ensures that u and v are cyclic, that is, their G-orbits span V^* and V. So we can write $x = \sum_i \alpha_i g_i u$ and $y = \sum_j \beta_j g_j v$, whence $\psi(g) = \sum_{i,j} \alpha_i \beta_j \phi(g_i^{-1} g g_j)$. □

Our fourth and final preliminary result is the following famous lemma.

LEMMA 4. (Van der Corput) *Suppose that $F : [a, b] \to \mathbf{R}$ is differentiable, its derivative F' is monotone, and $|F'| \geqslant 1$ on (a, b). Then $|\int_a^b e^{iF(t)}\, dt| \leqslant 3$.*

Proof. See [**S93**, p. 332], or [**R05**, Lemma 3] which actually gives the sharp bound 2. □

3. *Proof of Theorem 1*

By Lemma 1, it is enough to show that Haar measure on bV is the weak$*$ limit of probability measures μ_T concentrated on the orbit under consideration; or equivalently (Lemma 2), that the Fourier transforms of the μ_T tend pointwise to the characteristic function of $\{0\} \subset V^*$. (Here we identify the Pontryagin dual with the dual vector space or module.)

https://doi.org/10.1017/etds.2013.73 Published online by Cambridge University Press

To construct such μ_T, we assume without loss of generality that the action of G on V is effective, so that we may regard $G \subset \mathrm{GL}(V)$. Let $K \subset G$ be a maximal compact subgroup, $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ a Cartan decomposition, $\mathfrak{a} \subset \mathfrak{p}$ a maximal abelian subalgebra, $C \subset \mathfrak{a}^*$ a Weyl chamber, $P \subset \mathfrak{a}$ the dual positive cone, and H an interior point of P; thus we have that $\langle \nu, H \rangle$ is positive for all non-zero $\nu \in C$. (For all this structure see, for example, [**K73b**].) We fix a non-zero $v \in V$, and for each positive $T \in \mathbf{R}$ we let μ_T denote the image of the product measure Haar $\times$ (Lebesgue$/T$) $\times$ Haar under the composed map

$$\begin{aligned} K \times [0, T] \times K &\longrightarrow Gv \longrightarrow bV \\ (k, t, k') &\longmapsto k \exp(tH)k'v \\ w &\longmapsto e^{i\langle \cdot, w \rangle}. \end{aligned}$$

Here $\exp : \mathfrak{a} \to A$ is the usual matrix exponential with inverse $\log : A \to \mathfrak{a}$, and the brackets $\langle \cdot, \cdot \rangle$ denote both pairings, $\mathfrak{a}^* \times \mathfrak{a} \to \mathbf{R}$ and $V^* \times V \to \mathbf{R}$. By construction the μ_T are concentrated on the subset Gv of bV [**B04**, Corollary V.6.2.3]. It remains to show that as $T \to \infty$ we have, for every non-zero $u \in V^*$,

$$\int_{K \times K} dk\, dk' \, \frac{1}{T} \int_0^T e^{i\langle u, k \exp(tH)k'v \rangle}\, dt \to 0. \qquad (*)$$

To this end, let

$$F_{kk'}(t) = \langle u, k \exp(tH)k'v \rangle$$

denote the exponent in $(*)$. We will show that Lemma 4 applies to almost every $F_{kk'}$. In fact, it is well known (see, for example, [**K73b**, Proposition 2.4 and proof of Proposition 3.4]) that $\mathfrak{a}$ acts diagonalizably (over $\mathbf{R}$) on V. Thus, letting E_ν be the projector of V onto the weight ν eigenspace of $\mathfrak{a}$, we can write

$$F_{kk'}(t) = \sum_{\nu \in \mathfrak{a}^*} \langle u, kE_\nu k'v \rangle e^{\langle \nu, H \rangle t}.$$

Now we claim that there are non-zero ν such that the coefficient $f_\nu(k, k') = \langle u, kE_\nu k'v \rangle$ is not identically zero on $K \times K$. (Then f_ν, being analytic, will be non-zero *almost everywhere*.) Indeed, suppose otherwise. Then, writing any $g \in G$ in the form kak' (KAK decomposition [**K02**]), we would have

$$\langle u, gv \rangle = \sum_{\nu \in \mathfrak{a}^*} \langle u, kE_\nu k'v \rangle e^{\langle \nu, \log(a) \rangle} = \langle u, kE_0 k'v \rangle.$$

In particular, the matrix coefficient $\langle u, gv \rangle$ would be bounded. Hence so would all matrix coefficients, since they are linear combinations of translates of this one (Lemma 3); and this would contradict the non-compactness of $G \subset \mathrm{GL}(V)$.

So the set $N = \{\nu \in \mathfrak{a}^* : \nu \neq 0,\ f_\nu \neq 0\}$ is not empty. It is also Weyl group invariant, hence contains weights $\nu \in C$ for which we know that $\langle \nu, H \rangle$ is positive. Therefore, maximizing $\langle \nu, H \rangle$ over N produces a positive number $\langle \nu_0, H \rangle$, in terms of which our exponent and its derivatives can be written

$$\frac{d^n}{dt^n} F_{kk'}(t) = e^{\langle \nu_0, H \rangle t} \sum_{\nu \in \mathfrak{a}^*} f_\nu(k, k') \langle \nu, H \rangle^n e^{\langle \nu - \nu_0, H \rangle t},$$

where $\langle \nu - \nu_0, H \rangle < 0$ in all non-zero terms except the one indexed by ν_0. (Here we assume, as we may, that H was initially chosen outside the kernels of all pairwise

https://doi.org/10.1017/etds.2013.73 Published online by Cambridge University Press

differences of weights of V.) From this it is clear that for almost all (k, k') there is a T_0 beyond which the first two derivatives of $F_{kk'}$ are greater than 1 in absolute value. So Lemma 4 applies and gives

$$\left| \int_{T_0}^{T} e^{iF_{kk'}(t)}\, dt \right| \leqslant 3 \quad \text{for all } T.$$

Therefore, $\lim_{T\to\infty}(1/T)\int_0^T e^{iF_{kk'}(t)}\, dt = 0$ for almost all (k, k'), whence the conclusion $(*)$ by dominated convergence. This completes the proof.

4. *Outlook*

Theorem 1 says that the G-action on $V\backslash\{0\}$ is *minimal* [**P83**] in the Bohr topology. It would be interesting to determine if it is still minimal, and/or *uniquely ergodic*, on $bV\backslash\{0\}$.

It is also natural to speculate whether our theorems have a common extension to more general group representations. Here we shall content ourselves with noting two obstructions. First, Theorem 1 clearly fails for *semi*simple groups with compact factors. Secondly, Theorem 2 fails for V not of unipotent type, as one sees by observing that the orbits of $\mathbf{R}$ acting on $\mathbf{R}^2$ by $\exp \left(\begin{smallmatrix} t & 0 \\ 0 & -t \end{smallmatrix}\right)$ (i.e., hyperbolas) already have non-dense images in $\mathbf{R}^2/\mathbf{Z}^2$.

Acknowledgement. We thank Francis Jordan, who found the example in Remark 1.

References

[B87] M. Berger. *Geometry. I*. Springer-Verlag, Berlin, 1987.

[B74] J. Blum and B. Eisenberg. Generalized summing sequences and the mean ergodic theorem. *Proc. Amer. Math. Soc.* **42** (1974), 423–429.

[B04] N. Bourbaki. *Integration. I*. Springer, Berlin, 2004, Chs 1–6.

[D82] J. Dixmier. *C^*-algebras*. North-Holland, Amsterdam, 1982.

[G07] J. Galindo, S. Hernández and T.-S. Wu. Recent results and open questions relating Chu duality and Bohr compactifications of locally compact groups. *Open Problems in Topology. II*. Ed. E. Pearl. Elsevier, Amsterdam, 2007, pp. 407–422.

[G79] C. C. Graham and O. Carruth McGehee. *Essays in Commutative Harmonic Analysis*. Springer, New York, 1979.

[H63] E. Hewitt and K. A. Ross. *Abstract Harmonic Analysis*. Vol. 1 Springer, Berlin, 1963.

[K73a] Y. Katznelson. Sequences of integers dense in the Bohr group. *Proc. Roy. Inst. Tech. (Stockholm) (June 1973)* 79–86, available at http://math.stanford.edu/~katznel/.

[K02] A. W. Knapp. *Lie Groups beyond an Introduction*. Birkhäuser, Boston, 2002.

[K73b] B. Kostant. On convexity, the Weyl group and the Iwasawa decomposition. *Ann. Sci. Éc. Norm. Sup.* (4) **6** (1973), 413–455.

[P83] K. Petersen. *Ergodic Theory*. Cambridge University Press, Cambridge, 1983.

[R05] K. M. Rogers. Sharp van der Corput estimates and minimal divided differences. *Proc. Amer. Math. Soc.* **133** (2005), 3543–3550.

[S93] E. M. Stein. *Harmonic Analysis: Real-variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton University Press, Princeton, NJ, 1993.

[Z93] F. Ziegler. Subsets of R^n which become dense in any compact group. *J. Algebraic Geom.* **2** (1993), 385–387.

[Z96] F. Ziegler. Méthode des orbites et représentations quantiques. *PhD Thesis*, Université de Provence, Marseille, 1996, arXiv:1011.5056.

https://doi.org/10.1017/etds.2013.73 Published online by Cambridge University Press