Ergod. Th. & Dynam. Sys. (2010), **30**, 441–456
doi:10.1017/S0143385709000169
© *2009 Cambridge University Press*
Printed in the United Kingdom

Diffeomorphisms Hölder conjugate to Anosov diffeomorphisms

ANDREY GOGOLEV
Mathematics Department, Penn State University, University Park, State College, PA 16802, USA
(e-mail: agogolev@math.psu.edu)

(*Received* 8 *September* 2008 *and accepted in revised form* 9 *February* 2009)

Abstract. We show by means of a counterexample that a $C^{1+\mathrm{Lip}}$ diffeomorphism Hölder conjugate to an Anosov diffeomorphism is not necessarily Anosov. Also we include a result from the 2006 PhD thesis of Fisher: a $C^{1+\mathrm{Lip}}$ diffeomorphism Hölder conjugate to an Anosov diffeomorphism is Anosov itself provided that Hölder exponents of the conjugacy and its inverse are sufficiently large.

1. *Introduction*

Consider Anosov diffeomorphisms f and g of a compact smooth manifold M that are conjugate by a homeomorphism h:

$$h \circ f = g \circ h.$$

It is well known and easy to show that h is in fact Hölder continuous. When we say that the conjugacy is Hölder or that two diffeomorphisms are Hölder conjugate we mean that the conjugacy and its inverse are Hölder continuous.

It is natural to ask the following converse question.

Question. Is every diffeomorphism that is Hölder conjugate to an Anosov diffeomorphism itself Anosov?

This question was asked by Katok. His motivation came from differentiable rigidity of higher-rank Anosov actions. For example, a popular object of study is a $\mathbb{Z}^k$-action which contains Anosov elements and which is conjugate to an algebraic action for which Anosov elements are dense. If the answer to the question above were positive then we would immediately obtain that Anosov elements are dense in the original action. Moreover, the Weyl chamber picture in $\mathbb{R}^k$ for non-algebraic action would be the same as that for the algebraic action. Normally this information is unavailable or only available through difficult means otherwise. See the upcoming book [**KN08**] for an introduction to the rigidity of Anosov actions.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

Unfortunately the answer is negative. We provide a concrete counterexample of a $C^{1+\mathrm{Lip}}$ diffeomorphism of the 2-torus $\mathbb{T}^2$ Hölder conjugate to Anosov but not Anosov itself. In fact, the counterexample can be constructed to be C^r for any $r \in (1, 3)$ (see Remark).

The basic method to produce a non-Anosov diffeomorphism that is topologically conjugate to an Anosov diffeomorphism is to start with an Anosov diffeomorphism and isotope it pushing stable eigenvalues at a fixed point to the unit circle. This can be done so that stable and unstable foliations persist. They remain mutually transverse everywhere but not uniformly contracting and expanding. The new system is topologically conjugate to the original Anosov map. See [**K79**] for a detailed construction and the proof. Another similar example was considered in [**L80**]. It has an additional feature: stable and unstable manifolds at the fixed point are tangent.

Looking at the behavior of orbits approaching a fixed point along the stable manifold we have an exponentially fast approach for the Anosov map conjugated to a much slower sub-exponential approach. A Hölder continuous conjugacy would necessarily preserve the exponential speed, only changing the exponent. This shows that these diffeomorphisms are far from being Hölder conjugate. Note that in the meantime the conjugacy or its inverse may even turn out to be Lipschitz.

Another way to produce such a diffeomorphism is to start with an Anosov diffeomorphism and 'bend' the unstable manifold of a heteroclinic point R until stable and unstable manifolds at R become tangent. This isotopy can be performed locally in the neighborhood of R. The result is a diffeomorphism with stable and unstable foliations being transverse everywhere but along the orbit of R. Along this orbit stable and unstable manifolds exhibit a tangency. If we isotope inside of $\mathrm{Diff}^{\infty}(M)$, then the tangency is at least cubic: it cannot be quadratic since the stable and unstable foliations are topologically transverse. This bifurcation at the boundary of Anosov systems was independently studied by Enrich [**E98**] and Bonatti *et al* [**BDV98**]. It was shown in [**E98**] that the new system is conjugate to the original Anosov map. All periodic points remain hyperbolic. Hence, unlike in the previous situation, there is a hope that the topological conjugacy is in fact Hölder continuous.

We look at the simplest bifurcation of the type described above (Figure 1). Consider the arc f_t, $t \in [0, 1]$, that starts with a linear hyperbolic automorphism $L = f_0$ and ends with diffeomorphism $f = f_1$ with tangency at a heteroclinic point R. The difference is that instead of creating a cubic tangency we create a transverse quadratic tangency. The price we pay is that f_1 is only $C^{1+\mathrm{Lip}}$ smooth. Higher-order derivatives do not exist at the point $L^{-1}(R)$.

THEOREM 1. *Suppose that L and f are as in the previous paragraph. The conjugacy between L and f and its inverse are Hölder continuous with exponents equal to $1/2 - \delta$ and $1/4 - \delta$. The number δ can be made arbitrarily small by an appropriate choice of L and f.*

Remark. The number $1/4$ is not a sharp bound for the exponent. It clearly can be improved. For a diffeomorphism with a heteroclinic tangency of order $1 + \alpha$, $0 < \alpha < 2$, our arguments imply that the conjugacy and its inverse are Hölder continuous with

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press




FIGURE 1. Heteroclinic tangency.

exponents $1/(1+\alpha)-\delta$ and $(2-\alpha)/(2(1+\alpha))-\delta$. Clearly such a diffeomorphism is only $C^{1+\alpha-\varepsilon}$. We stick to the case $\alpha=1$ mainly to avoid cumbersome notation. Note that if α is close to zero, then both exponents are close to 1. In the smoothness class C^3 and higher our arguments fail.

In the next section we point out that this construction also provides an example of a system for which the Mather spectrum differs from the periodic spectrum.

We heavily rely on the results in [**E98**] as well as [**BDV98**] and [**C98**]. Thus, in §3 we formulate results that are relevant to our goal. In §4 we prove that the conjugacy is Hölder continuous.

Finally in the last section we present a very short proof of a positive result from the 2006 PhD thesis of Travis Fisher that complements our own.

THEOREM 2. [**F06**] *A $C^{1+\mathrm{Lip}}$ diffeomorphism that is conjugate to an Anosov diffeomorphism via a Hölder conjugacy h is Anosov itself provided that the product of Hölder exponents for h and h^{-1} is greater than $1/2$.*

Remark. This result also holds for any hyperbolic set; the proof is the same. Also, we remark that we have removed an unnecessary condition that was present in the formulation of the result in [**F06**].

2. *Periodic spectrum versus Mather spectrum*

Recall the definition of the Mather spectrum. Denote by $\Gamma(TM)$ the set of continuous vector fields with supremum norm. Given a diffeomorphism $f\colon M\to M$ define $f_*\colon \Gamma(TM)\to\Gamma(TM)$

$$f_*v(\cdot)=Df(v(f^{-1}(\cdot))).$$

The spectrum Q_f of the complexification of f_* is called the Mather spectrum of f.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

THEOREM 3. (Mather [**Math68**]) *If non-periodic points of f are dense, then any connected component of Q_f is an annulus centered at 0. A diffeomorphism f is Anosov if and only if $1 \notin Q_f$.*

Define the periodic spectrum of a diffeomorphism. Given a periodic point x of period p denote by $\{\lambda_1(x)^p, \dots \lambda_d(x)^p\}$ the set of absolute values of eigenvalues of $Df^p(x)$. Then

$$P_f \stackrel{\text{def}}{=} \overline{\bigcup_{x \in \text{Per}(f)} \{\lambda_1(x), \dots, \lambda_d(x)\}}.$$

The following is easy to prove.

PROPOSITION 4. *Let f be an Anosov diffeomorphism of $\mathbb{T}^2$. Then $P_f = Q_f \cap \mathbb{R}_+$.*

In contrast to the above, Theorems 1 and 3 imply the following result.

COROLLARY 5. *Diffeomorphism f from Theorem 1 provide an example of a diffeomorphism with dense set of periodic points such that $P_f \neq Q_f \cap \mathbb{R}_+$.*

3. *First heteroclinic tangency at the boundary of Anosov systems*

Here we describe some results of [**E98**], [**BDV98**] and [**C98**] that we need.

Let L be hyperbolic automorphism of $\mathbb{T}^2$. Denote by e_u and e_s the eigenvectors of L and by $\lambda > 1$ the unstable eigenvalue, $Le_u = \lambda e_u$. Let P and Q be two different fixed points of L and let R be an intersection of the stable manifold of P and the unstable manifold of Q. We may assume that distances to R from P and Q are equal. Also we assume that the size of a ball containing $\{P, Q, R\}$ is much smaller than the size of $\mathbb{T}^2$. Let (x, y) be coordinates in the neighborhood of R that make stable foliation horizontal and unstable foliation vertical. Let B be a small ball of radius r centered at R.

Define $f_t = \theta_t \circ L$, $t \in [0, 1]$, where $\theta_t : \mathbb{T}^2 \to \mathbb{T}^2$ is the identity outside of B and given by the following formula on B:

$$\theta_t(x, y) = \begin{pmatrix} \cos(t\gamma(\rho)) & \sin(t\gamma(\rho)) \\ -\sin(t\gamma(\rho)) & \cos(t\gamma(\rho)) \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$$

where $\rho = \sqrt{x^2 + y^2}$ and $\gamma : [0, \infty) \to [0, \pi/2]$ is a C^∞ map satisfying $\gamma(0) = \pi/2$; $\gamma(\rho) = 0$ for $\rho \geq r$; γ is strictly decreasing on $[0, r]$. Thus, on every circle centered at R, θ_t is a rotation by an angle no greater than $\pi/2$. Value $\pi/2$ is achieved for $\rho = 0$ and $t = 1$. Let $v \in T_R\mathbb{T}^2$ be the unit vertical vector. From the definition of $f = f_1$ we have

$$\lim_{n \to \pm\infty} (Df^n)v = 0.$$

Hence, f is not Anosov.

Denote by $\mathcal{U}$ and $\mathcal{V}$ neighborhoods of segments PR and $Qf^{-1}(R)$ that contain B and $f^{-1}(B)$, respectively.

THEOREM 6. (Enrich [**E98**]; Bonatti *et al* [**BDV98**]; Carvalho [**C98**]) *There exist r small enough and a function γ such that the corresponding arc f_t as defined above satisfies the following.*

(T1) *Diffeomorphisms f_t are Anosov for $t < 1$.*

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

(T2) *A diffeomorphism f possesses invariant contracting and expanding foliations W^s and W^u. The leaves of W^s and W^u are C^1 immersed curves. Denote by E^s and E^u the distributions tangent to these foliations.*

(T3) *Foliations W^s and W^u are transverse everywhere but along the orbit of R. At the point R they have cubic tangency. Namely, there is $\tau > 1$ such that for $S(x, y) \in B$*

$$\tan \measuredangle(E^u(S), e_s) \geq \tau(x^2 + y^2). \tag{1}$$

Analogous inequality holds for the distribution E^s.

(T4) *For any $S \notin \mathcal{U}$*

$$\tan \measuredangle(E^u(S), e_u) < \varepsilon.$$

The number ε can be made arbitrarily small by the choice of L, f and $\mathcal{U}$. If we let

$$B_\infty = \{x \in \mathbb{T}^2 : \exists i > 0 \text{ such that } f^{-i}(x) \in B, \{f^{-i}(x), f^{-i+1}(x), \ldots x\} \subset \mathcal{U}\},$$

then, for any $S \notin B_\infty$,

$$\tan \measuredangle(E^u(S), e_u) < 1.$$

Analogous statement holds for E^s and $\mathcal{V}$.

(T5) *Diffeomorphism f is conjugate to L by a homeomorphism h, $h \circ f = L \circ h$.*

Remark. Technical statements (T3) and (T4) are not stated explicitly in the papers quoted but they follow from the cone constructions that are carried out there.

It may seem that since the size of B is small E^u is almost vertical inside of B_∞ and almost horizontal outside. In fact, the transition through the boundary of B_∞ is continuous. The parameter τ increases when r goes to zero.

We are working with exactly the same construction, but θ must be chosen differently. The function γ can be chosen differently with $\gamma^{(r)}(0) = 0$ for $r < 2$ and $\gamma''(0) < 0$. Then $\theta \in C^{1+\mathrm{Lip}}$ and the tangency is quadratic. In this way, instead of (T3), we have

$$\tan \measuredangle(E^u(S), e_s) \geq \tau\sqrt{x^2 + y^2}, \quad \tau > 1. \tag{2}$$

We provide an outline of the proofs of (T4) and (2) at the end of this section.

Remark. For conservative systems the theorem above was established in [**C98**]. The original proof [**E98**] required that the product of the eigenvalues at P is greater than one while the product of the eigenvalues at Q is less than one. The assumption on the eigenvalues at P and Q in [**BDV98**] is even more restrictive. The main motivation of [**C98**] was to extend the example to systems with homoclinic tangency. Our proof works for homoclinic intersection as well. We work with a heteroclinic intersection only for convenience. Also we would like to remark that our proof of Hölder continuity works for the original construction in [**E98**]. One needs to start the isotopy with a C^1 small perturbation of L that satisfies the above assumptions on eigenvalues at P and Q instead of starting with L.

Let us recall the proof of (T5) from [**E98**] since this is the statement that we strengthen.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press




FIGURE 2. The hatched set is B_∞. The distance $|QR|$ is much larger than $f^N(B) = f^2(B)$. Hence, it is possible to fatten $\mathcal{U}$ to $\tilde{\mathcal{U}}$ so that the unstable distribution outside $\tilde{B}_\infty$ is ε-close to the horizontal vector e_u.

Proof. The main tool here is the following result of Walters.

THEOREM 7. (Walters [**W70**]) *Let $g\colon M \to M$ be an Anosov diffeomorphism. Then there exists an $\varepsilon_0 > 0$ such that for each $0 < \varepsilon < \varepsilon_0$ there exists $\delta > 0$ such that if $\tilde{g}\colon M \to M$ is a homeomorphism and if $d(g, \tilde{g}) < \delta$, then there exists a unique continuous map h of M onto M with $h \circ \tilde{g} = g \circ h$ and $d(h, id) < \varepsilon$.*

Apply the theorem for $g = L$ and $\tilde{g} = f$ to obtain semiconjugacy h with $d_{C^0}(h, id) < \varepsilon$. Note that $d_{C^0}(L, f) \to 0$ as $r \to 0$. We have to take r small enough so that ε is smaller than a constant associated with the local product structure of W^s and W^u. This guarantees that h is injective. Indeed if h glues together some points on, say, unstable manifolds, then iterating forward we obtain that h glues together some points a fixed distance apart. This is impossible since h is close to identity.

Hence, by the invariance of domain theorem, h is a homeomorphism. □

Sketch of Proof of (T4). Cone constructions in [**E98**] and [**C98**] imply that for all $S \notin B_\infty$

$$\tan \measuredangle(E^u(S), e_u) < 1.$$

Then given a small number ε there exists N such that for all $S \notin \bigcup_{i=0}^{N} f^i(B_\infty)$

$$\tan \measuredangle(E^u(S), e_u) < \varepsilon.$$

It is possible to fatten $\mathcal{U}$ so that the set $\tilde{B}_\infty$ that corresponds to the new fattened neighborhood $\tilde{\mathcal{U}}$ of PR contains $\bigcup_{i=0}^{N} f^i(B_\infty)$ as shown in Figure 2.

For that we need to make sure that $f^N(B)$ is small compared with the distance $|QR|$. This can be achieved by an appropriate choice of automorphisms L, P and Q.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

We fix a hyperbolic matrix L that induces an automorphism of $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$ with d fixed points. We fix the size r of the ball B and the map $\theta|_B$. The trick now is to choose the torus $\mathbb{T}^2$ to be 'large' when compared with the eigenvalue λ of L and r.

The linear map L induces a hyperbolic automorphism of $\mathbb{T}^2 = \mathbb{R}^2/k\mathbb{Z}^2$, where k is a large integer. This automorphism is a finite cover of the automorphism of $\mathbb{T}^2 = \mathbb{R}^2/\mathbb{Z}^2$. It also has d fixed points. Obviously the distances between those fixed points are now large. Hence, P and Q can be chosen so that $|QR|$ is large. □

Remark. We use the fact that $|QR|$ can be chosen large independently of r and λ several times in the course of the proof of Hölder continuity.

PROPOSITION 8. *Given a point $S \in \mathcal{U}$. Denote by $d(S)$ the distance to PR. Then*

$$\tan \measuredangle(E^s(S), e_s) \le \kappa d(S)^2. \tag{3}$$

where κ is a number that depends on ε from (T4).

Proof of Proposition. Let N be the smallest positive integer such that $f^N(S) \notin \mathcal{U}$. Then $\lambda^n d(S) \approx 1$ and $\tan \measuredangle(E^s(f^N(S)), e_s) \le \varepsilon$ by (T4) since $f^N(S) \notin \mathcal{V}$ as well:

$$\tan \measuredangle(E^s(S), e_s) = \lambda^{-2n} \tan \measuredangle(E^s(f^N(S)), e_s) \le C\varepsilon d(S)^2. \qquad \square$$

Sketch of proof of (2). Denote by A and $\tilde{A}$ points of intersection of the line QR and boundary ∂B as shown in Figure 3. It follows from the definition of f that the straight segment $[Q, A]$ is inside $W^u(Q)$ while the straight segment $[P, R]$ is inside $W^s(P)$. The shape of $W^u(Q)$ between A and $\tilde{A}$ is completely determined by θ. Namely, it is the image of the segment $[A, \tilde{A}]$ under θ. We remark that on the other hand the shape of $W^u(Q)$ between A and R determines θ since the map θ is a rigid rotation on every circle around R.

First let us establish the linear variation of the angle (2) along $\theta[A, R]$. It follows from the choice of θ. Let $C \in [A, R]$ be the point such that $\measuredangle(E^u(\theta(C)), e_s) = \pi/2$. Then for any $S \in \theta[C, R]$ we have

$$\tan \measuredangle(E^u(S), e_s) \ge \tau\sqrt{x^2 + y^2}$$

since the tangency is quadratic. For $S \in \theta[A, C]$

$$\tan \measuredangle(E^u(S), e_s) \ge c$$

where c is a large constant, $c \gg 1$.

Fix a circle $\mathcal{C}$ of radius ρ centered at R. Let $D = \mathcal{C} \cap \theta[A, R]$. We know that the estimate (2) holds for D and we would like to establish it for other points on $\mathcal{C}$.

Consider the distribution $\tilde{E}^u \subset T_{\mathcal{C}}\mathbb{T}^2$ given by the formula $\tilde{E}^u = DL\, E^u$. Then E^u on $\mathcal{C}$ is given by $E^u = Df\, E^u = D\theta \tilde{E}^u$.

If we denote by v and u the normal and the tangent vector fields to $\mathcal{C}$, then $D\theta$ with respect to bases $(v(\cdot), u(\cdot))$ and $(v(\theta(\cdot)), u(\theta(\cdot)))$ is given by the shear matrix

$$D\theta = \begin{pmatrix} 1 & \alpha \\ 0 & 1 \end{pmatrix}, \quad \alpha = \alpha(\rho) > 0. \tag{4}$$

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press




FIGURE 3. Establishing linear variation of the angle on the circle $\mathcal{C}$.

According to Proposition 8

$$\tan \measuredangle(\tilde{E}^u(S), e_u) \leq \kappa \rho^2, \quad S \in \mathcal{C}.$$

Meanwhile

$$\tan \measuredangle(E^u(D), e_s) \geq \tau \rho.$$

These inequalities together with the formula for $D\theta$ above imply linear variation of the angle (2) for all $S \in \mathcal{C}$. □

4. *Topological conjugacy is Hölder*

Here we prove that the conjugacy h and its inverse are Hölder continuous.

We mimic the standard proof of Hölder continuity of conjugacy between two Anosov systems (e.g. [**KH95**, §19.1]). The conjugacy maps stable and unstable foliations of f, W^s and W^u, into stable and unstable foliations of L, W^s_L and W^u_L, respectively.

Step 1. A restriction of h to a leaf of W^u is Hölder continuous with the exponent equal to $1 - \delta$. A restriction of h^{-1} to a leaf of W^u_L is Hölder continuous with the exponent equal to $1/4 - \delta$. The number δ depends on the choice of L and r and can be made arbitrarily small.

Remark. The distances with respect to which we show Hölder continuity are induced Riemannian distances on the leaves of W^u and W^u_L.

Analogously, h and h^{-1} are Hölder continuous when restricted to stable leaves with exponents $1 - \delta$ and $1/4 - \delta$, respectively. We immediately obtain the following result.

PROPOSITION 9. *The homeomorphism h^{-1} is Hölder continuous with exponent $1/4 - \delta$.*

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

This follows from a standard argument that utilizes uniform transversality of W^s_L and W^u_L.

To conclude that h is Hölder as well we need W^s and W^u to be uniformly transverse. In our situation W^s and W^u are uniformly transverse only outside a neighborhood of the orbit of R. This leads to further loss of the exponent by a factor of $1/2$.

Step 2. Conjugacy h is Hölder continuous with exponent equal to $1/2 - \delta$.

Heuristically it is clear that the loss of the exponent at the second step is inevitable. In the second step we 'straighten out' the quadratic tangency. Thus, the exponent is no greater than $1/2$.

Together with (2), Proposition 8 gives us control on the angle between E^s and E^u in the neighborhood of the orbit of R which is crucial for carrying out estimates in Step 2.

Throughout the proof we denote by d^u, d^s, d^u_L and d^s_L the induced Riemannian distances along the leaves of the corresponding foliations.

Proof of Step 1. Uniform continuity of h implies that there exists $C_1 > 0$ such that

$$\frac{1}{C_1} d^u_L(h(a), h(b)) \le d^u(a, b) \le C_1 d^u_L(h(a), h(b)) \quad \text{whenever } d^u(a, b) \ge \frac{r}{10}.$$

To prove Hölder estimates for nearby points a and $b \in W^u(a)$ we need to have exponential estimates on expansion along E^u. Given a point $a \in \mathbb{T}^2$ let

$$D^u(a) = \|Df(a)v^u\|, \quad v^u \in E^u(a), \|v^u\| = 1.$$

If $a \notin L^{-1}(B)$, then, obviously, $\lambda^{-1} \le D^u a \le \lambda$. If $a \in L^{-1}(B)$, then $D^u a$ can be larger. Still there exists D such that $D^u a \le D$ for $a \in L^{-1}(B)$. In fact, using estimates on α from (4) one can show that $D = 2\lambda$ works but we do not use it. Keeping r and θ fixed choose L (e.g. pass to a finite cover as in the proof of (T4)), P and Q so that $|PR|$ is large and hence the first return time m to $L^{-1}(B)$ is large. Thus, for all $a \in \mathbb{T}^2$

$$\prod_{i=1}^{m-1} D^u f^i(a) \le \lambda^{m-1} D \le (\lambda^{1+\delta})^m, \tag{5}$$

where $\delta = \delta(m, D)$ is a small number. It follows that there exists C_2 such that for all $n > 0$

$$\|Df^n v^u\| \le C_2(\lambda^{1+\delta})^n \|v^u\|, \quad v^u \in E^u. \tag{6}$$

Now we use a standard argument to prove Hölder continuity. Take $a \in W^u(b)$ close to b. Let N be the smallest number such that $d^u(f^N(a), f^N(b)) \ge r/10$. Then

$$\begin{aligned} d^u_L(h(a), h(b)) &= \frac{1}{\lambda^N} d^u_L(L^N(h(a)), L^N(h(b))) \le \frac{C_1}{\lambda^N} d^u(f^N(a), f^N(b)) \\ &\le \frac{C_1 C_3}{\lambda^N} d^u(f^N(a), f^N(b))^{1-\delta} \\ &\le \frac{C_1 C_2^{1-\delta} C_3}{\lambda^N} (\lambda^{1+\delta})^{N(1-\delta)} d^u(a, b)^{1-\delta} \le C d^u(a, b)^{1-\delta}. \end{aligned}$$

To show that h^{-1} is Hölder along W^u_L we need an estimate on the product in (5) from below. According to (T4) if $a \notin \mathcal{U} \cup L^{-1}(B)$, then $D^u(a) \ge \mu$, where $\mu = \mu(\varepsilon)$ and $\mu \nearrow \lambda$

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

when $\varepsilon \to 0$ (we remark that the choices we make to enure that ε is small do not affect λ). If $a \in \mathcal{U} \setminus B_\infty$, then another inequality from (T4) provides an estimate on expansion.

If $a \in B_\infty$, then, obviously, $D^u(a) \geq \lambda^{-1}$ but we need to have a better control. Split B_∞ into its connected components

$$B_\infty = \bigcup_{i \geq 0} B_i$$

(see Figure 2). Let (x^i, y^i) be coordinates in B_i obtained by parallel transport of (x, y) from R to $f^i(R)$. Consider rectangles

$$\bar{B}_i = \{(x^i, y^i) : |x^i| \leq r\lambda^{-i},\ |y^i| \leq r\lambda^{-3i}\}. \tag{7}$$

Also let

$$\bar{B} = \bigcup_{i \geq 0} \bar{B}_i. \tag{8}$$

Let $\xi = \tau r$ where τ is from (2).

LEMMA 10. *Consider a point $S(x^n, y^n) \in B_n$, $S \notin \bar{B}_n$. Then*

$$\tan \measuredangle(E^u(S), e_s) \geq \xi.$$

Proof. Note that this follows from (2) if $n = 0$.

$$\tan \measuredangle(E^u(S), e_s) = \lambda^{2n} \tan \measuredangle(E^u(f^{-n}(S)), e_s).$$

Clearly, $(\lambda^{-n}x^n, \lambda^n y^n)$ are (x, y) coordinates of $f^{-n}(S)$. If $|x^n| \geq r\lambda^{-n}$, then

$$\lambda^{2n} \tan \measuredangle(E^u(f^{-n}(S)), e_s) \geq \lambda^{2n}\tau(\lambda^{-n}x^n) \geq \tau r.$$

If $|y^n| \geq r\lambda^{-3n}$, then

$$\lambda^{2n} \tan \measuredangle(E^u(f^{-n}(S)), e_s) \geq \lambda^{2n}\tau(\lambda^n y^n) \geq \tau r. \qquad \square$$

Take an arbitrary point $a \notin \bar{B}$. Then $\tan \measuredangle(E^u(a), e^s) \geq \xi$ and hence there exists $m > 0$ such that $D^u f^m(a) > \mu$. Therefore, contraction along the unstable distribuition mostly happens only inside $\bar{B}$. After the point leaves $\bar{B}$ its unstable direction 'recovers' after m steps even before the point leaves $\mathcal{U}$.

Now we estimate from below the expansion along E^u by analyzing the itinerary of a point.

As before (in the sketch of the proof of (T4)), we make sure by the choice of L that after a point leaves $\mathcal{U}$ it spends at least time $4m + 4$ before it enters $\mathcal{V}$.

Take a point $a(x, y) \in B$ and assume that $f^i(a) \in \bar{B}_i$, $i = 0, \ldots n$, $f^{n+1}(a) \notin \bar{B}_{n+1}$. By the lemma, $|x| \leq r\lambda^{-2n}$, $|y| \leq r\lambda^{-2n}$. Hence, $f^{-1}(a), \ldots f^{-2n}(a) \in \mathcal{V}$, $D^u f^{-i}(a) \geq \mu$, $i = 2, \ldots, 2n$. A simple calculation with $D\theta$ shows that $D^u f^{-1}(a) \geq 1$. Start with $f^{-2n}(a)$ and wait time $4n + 4m + 4$. We know for sure that during that time the orbit has not entered $\mathcal{V}$ again. Thus,

$$\begin{aligned}\prod_{i=-2n}^{2n+4m+4} D^u f^i(a) &= \prod_{i=-2n}^{-1} D^u f^i(a) \prod_{i=0}^{n+m} D^u f^i(a) \prod_{i=n+m+1}^{2n+4m+4} D^u f^i(a) \\ &\geq \mu^{2n-1}\lambda^{-n-m-1}\mu^{n+3m+4} = \lambda^{-n-m-1}\mu^{3n+3m+3}.\end{aligned} \tag{9}$$

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

This estimate is good enough to obtain the exponent $1/2 - \delta$. The only problem is that it holds only along specific orbit segments as described above. We call them 'cycles'.

As before take $a \in W^u(b)$ close to b and let N be the smallest number such that $d^u(f^N(a),\ f^N(b)) \geq r/10$.

Let us first explain the idea informally. We have to study how the length of $d^u(f^i(a),\ f^i(b))$ changes as $i = 0, \ldots, N$. We decompose this time segment into 'cycles' as in (9). These 'cycles' do not overlap. There might be some 'gaps' between the 'cycles' that only improve the estimate since the time spent in the 'bad' set $\bar{B}_\infty$ is inside the 'cycles'. The difficulty that we have to deal with is that at the beginning 'cycle' might be 'incomplete'. The 'worst' situation is when a is close to R. The same problem occurs at the end: the last 'cycle' might happen to be 'cut' at the end.

Note that in the description above we ignore returns to $\mathcal{U} \setminus B$. The expansion rate inside this set might be less than μ but still is greater than one according to (T4). Therefore, these returns are much easier to take care of. Hence, we consider only 'cycles' that correspond to returns to B.

The problem of the 'cut' at the end is easy to deal with. Denote by $t_j,\ j = 1, \ldots, l$, the lengths of 'cycles'. We count incomplete 'cycles' as well. We also consider numbers $n_j,\ j = 1, \ldots, l$. In the notation of (9) $t_j = 4n_j + 4m + 4$. Assume that the last 'cycle' number l is incomplete. During the last 'cycle' the segment enters B, spends time n_l inside $\bar{B}_\infty$ and then it recovers to the size of $r/10$ during the time less than $n_l + 4m + 4$. When the segment leaves $\bar{B}_{n_l}$ it has length of order λ^{-3n_l} since that is the vertical size of $\bar{B}_{n_l}$ and the unstable foliation is roughly vertical in $\bar{B}_{n_l}$. Clearly the number n_l is not large since otherwise the segment cannot recover to the size $r/10$. We can control N independently of n_l by choosing a and b extremely close to each other. This may result in large Hölder constant but the exponent will not be affected. In fact, since n_l is bounded this difficulty at the end of the 'time window' $i = 0, \ldots, N$ can be taken care of by the Hölder constant.

Now assume that the first 'cycle' is incomplete as well. In contrast to above n_1 might happen to be large when compared with N since the size of the segment is small at the beginning. Clearly we need to examine the 'worst' situation when $a \in B$. As before we can argue that after time n_1 the size of the segment is of the order λ^{-3n_1}. The number $N - n_1$ is greater than $3n_1$ since during this time the segment grows up to size $r/10$. Hence, using (9) we obtain

$$\begin{aligned} d^u(f^N(a),\ f^N(b)) &\approx \lambda^{-n_1}\lambda^{n_1}\lambda^{\frac{3}{4}(N-2n_1)}\lambda^{-\frac{1}{4}(N-2n_1)}d^u(a,b) \\ &= \lambda^{-\frac{1}{2}(N-2n_1)}d^u(a,b) = \lambda^{-\frac{1}{2}(\frac{1}{2}N+\frac{1}{2}N-2n_1)}d^u(a,b) \geq \lambda^{\frac{1}{4}N}d^u(a,b). \end{aligned} \tag{10}$$

This clearly good enough to obtain the exponent $\frac{1}{4} - \delta$.

From now on we are providing details to the scheme described above. Still, we stay away from a completely rigorous technical discussion. The technical details are plentiful while the way the argument works is fairly transparent.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

To obtain the estimate $N \geq 4n_1$ we need to redefine the set $\bar{B}$ slightly, numbers n_i and t_i accordingly. First fix μ close to λ, $\mu < \lambda$.

$$\bar{B}_i = \{(x^i, y^i) : |x^i| \leq \tilde{r}\lambda^{-i}, |y^i| \leq \tilde{r}\lambda^{-3i}\},$$
$$\bar{B} = \bigcup_{i \geq 0} \bar{B}_i,$$

where $\tilde{r} = \tilde{r}(\mu) < r/20$ is chosen so that $D^u(x) \geq \mu^{-1}$ for any $x \in B \setminus \bar{B}_0$ and by the arguments of Lemma 10 $D^u(x) \geq \mu^{-1}$ for any $x \notin \bar{B}_\infty$.

We study lengths of the segment during the time N defined above. We consider the 'worst' case when a and b are close to R. Let n_1 be the smallest integer such that

$$\frac{d^u(f^{n_1+1}(a),\, f^{n_1+1}(b))}{d^u(f^{n_1}(a),\, f^{n_1}(b))} \geq \mu^{-1}.$$

Then it is easy to see that after some fixed number of iterates $m = m(\mu)$ which is independent of n_1 we have

$$\frac{d^u(f^{n_1+m+1}(a),\, f^{n_1+m+1}(b))}{d^u(f^{n_1+m}(a),\, f^{n_1+m}(b))} \geq \mu.$$

Remark. Note that any iterate of the segment does not cross more than one connected component of $\bar{B}_\infty$. Indeed, the vertical distance between $\bar{B}_n$ and $\bar{B}_{n+1}$ is of the order of λ^{-n}, horizontal sizes of $\bar{B}_n$ and $\bar{B}_{n+1}$ are of order λ^{-3n} while in the 'gap' between B_n and B_{n+1} unstable foliation is in the horizontal cone field according to (T4). It follows that local unstable leaves do not intersect $\bar{B}_n$ and $\bar{B}_{n+1}$.

Analogously define numbers n_j, $j = 1, \ldots, l$. Then define $t_j = n_j + 4m + 4$ and the corresponding 'cycles' as before. Clearly we have an analog of (9)

$$d^u(f^{s_j+t_j}(a),\, f^{s_j+t_j}(b)) \geq \lambda^{-n-m-1}\mu^{3n+3m+3} d^u(f^{s_j}(a),\, f^{s_j}(b)), \tag{11}$$

where s_j is the starting time of a full 'cycle'. If the itinerary of the segment has complete first 'cycle', then in the same way as in the proof of Hölder continuity of h using (11) we obtain

$$d^u(h^{-1}(a), h^{-1}(b)) \leq C d^u(a, b)^{1/2-\delta}$$

with $\delta = \delta(\mu) \searrow 0$ as $\mu \nearrow \lambda$.

Now we go back to the 'worst' case when the first 'cycle' starts near R. Definitions of $\bar{B}_\infty$ and n_1 guarantee that at least half of the segment $[f^{n_1-1}(a),\, f^{n_1-1}(b)] \subset W^u(f^{n_1-1}(a))$ lies inside of $\bar{B}_{n_1-1}$. Recall that unstable foliation is almost vertical inside $\bar{B}_\infty$. Hence,

$$d^u(f^{n_1-1}(a),\, f^{n_1-1}(b)) \leq \frac{r}{10}\lambda^{-3(n_1-1)}.$$

Recall that $D^u(x) \leq \lambda$ for any $x \notin L^{-1}(B)$ and $d^u(f^N(a),\, f^N(b)) \geq r/10$. It follows that $N - n_1 \geq \Gamma n_1$, where Γ depends on the frequency of visits to $L^{-1}(B)$ and can be made arbitrarily close to three. Now the preliminary estimate (10) transforms into the following:

$$\begin{aligned} d^u(f^N(a),\, f^N(b)) &\geq \lambda^{-n_1-m}\mu^{n_1+3m+4}\mu^{\frac{3}{4}(N-2n_1)}\lambda^{-\frac{1}{4}(N-2n_1)} \\ &\geq \lambda^{-n_1-m}\mu^{n_1+3m+4}(\mu/\lambda)^{\frac{1}{4}(N-2n_1)}\mu^{\frac{1}{2}(N-2n_1)} \\ &= \lambda^{-n_1-m}\mu^{n_1+3m+4}(\mu/\lambda)^{\frac{1}{4}(N-2n_1)}\mu^{\frac{1}{2}(1-\Delta)(N-2n_1)}\mu^{\frac{1}{2}\Delta(N-2n_1)} \\ &\geq \mu^{\frac{1}{2}\Delta(N-2n_1)} \geq \mu^{\frac{1}{4}(N-\delta)}, \end{aligned}$$

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

where $\delta = \delta(\Gamma, \Delta)$ is small and $\Delta = \Delta(\lambda, \mu)$ is chosen so that $\mu^{\frac{1}{2}(1-\Delta)(N-2n_1)}$ compensates the factors in front of it. The number $\Delta \nearrow 1$ as $\mu \nearrow \lambda$. It follows that δ can be arbitrarily small. □

Proof of Step 2. First of all let us note that outside of $\mathcal{U}$ foliations W^s and W^u are uniformly transverse. Then, by the standard argument, Hölder continuity along W^s and W^u implies Hölder continuity of h with exponent $1 - \delta$ outside of $\mathcal{U}$.

It follows from (2) and Proposition 8 that inside B the angle $\measuredangle(E^s, E^u)$ varies linearly with distance to R. This allows us to show that h is Hölder continuous with exponent $1/2 - \delta$ inside B. More work is required to establish Hölder continuity in the rest of $\mathcal{U}$. We start with an observation that allows us to reduce our task to establishing Hölder inequality for points inside of a single B_n, $n \geq 0$.

Introduce vertical and horizontal cones

$$\mathcal{C}_v(x) = \{v \in T_x\mathbb{T}^2 \mid \tan \measuredangle(v, e_s) < \varepsilon\},$$
$$\mathcal{C}_h(x) = \{v \in T_x\mathbb{T}^2 \mid \tan \measuredangle(v, e_s) > \xi\}$$

with ε as in (T4) and ξ as in Step 1. These cones have disjoint interiors. Moreover, $E^s(x) \in \mathcal{C}_v(x)$ for any $x \in \mathcal{U}$ by (T4) and $E^u(x) \in \mathcal{C}_h(x)$ for any $x \in \mathcal{U} \setminus \bar{B}_\infty$ by Lemma 10. Thus, $\mathcal{C}_v$ and $\mathcal{C}_h$ provide good control of E^s and E^u in $\mathcal{U} \setminus \bar{B}_\infty$.

Take a and b nearby inside $\mathcal{U}$. Let e be the intersection of local unstable manifold $W^u(a, r/10)$ and local stable manifold $W^s(b, r/10)$.

LEMMA 11. *Assume that some fixed proportion with respect to the length of local unstable manifold connecting a and e lies inside of $\mathcal{C}_h$, meaning that the tangent vector is in the cone. Then*

$$d(h(a), h(b)) \leq C d(a, b)^{1-\delta}.$$

Recall that we know that the local stable manifold is in $\mathcal{C}_v$. Then the proof of the lemma is a straightforward adjustment of the standard proof when the whole local unstable manifold lies inside of $\mathcal{C}_h$ as well.

We have remarked in the course of the proof of Step 1 that local unstable leaves do not meet different connected components of $\bar{B}_\infty$. Together with Lemma 11 this implies that we are only left to deal with points a and b such that local unstable manifold connecting a and e lies almost entirely in $\bar{B}_n$ for some $n \geq 0$.

Denote by (x, y) the coordinate system centered at $f^n(R)$ with x-axis being horizontal. Observe further that it is enough to consider points a and b that have the same y-coordinate. Let $a = a(x_1, y_1)$, $b = b(x_2, y_1)$ and $e = e(x_3, y_3)$. We are aiming at proving the estimate

$$C|x_1 - x_2|^{\frac{1}{2}} \geq |y_1 - y_3|. \tag{12}$$

Together with Step 1 this would imply that h is Hölder with exponent $\frac{1}{2} - \delta$.

Given a point $S(x, y) \in B_n$ we have

$$\tan \measuredangle(E^u(S), e_s) = \lambda^{2n} \measuredangle(E^u(f^{-n}(S)), e_s) \geq \lambda^{2n} \tau \sqrt{(\lambda^{-n}x)^2 + (\lambda^n y)^2}$$
$$\geq \lambda^{2n} \tau \lambda^{-n} \sqrt{x^2 + y^2} \geq \sqrt{x^2 + y^2}$$

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press




FIGURE 4. Beaks.

and by Proposition 8

$$\tan \measuredangle(E^s(S), e_s) \le \kappa d(S)^2 \le \frac{x^2}{2}.$$

These inequalities provide control on the angle needed to carry out the estimate (12).

Denote by $\mathcal{B}$ the 'beak' formed by the unstable manifold connecting a and e and the stable manifold connecting b and e. We consider two representative cases as illustrated in Figure 4.

Case A. Assume that $D \stackrel{\text{def}}{=} \operatorname{dist}(\mathcal{B}, f^n(R)) \ge |y_1 - y_3|$. In this case according to the estimates above we have that $[a, e]$ is tilted by at least D while $[b, e]$ is tilted by at most D^2. Hence,

$$|x_1 - x_2| \ge |y_1 - y_3|(D - D^2) \ge \tfrac{1}{2}|y_1 - y_3|D \ge \tfrac{1}{2}|y_1 - y_3|^2.$$

Case B. We allow $|y_1 - y_3|$ to be greater than $\operatorname{dist}(\mathcal{B}, f^n(R))$. Also we make a simplifying assumption that $x_3 = 0$ and $y_3 > 0$. Then

$$\begin{aligned} |x_1 - x_3| &\ge \int_{y_3}^{y_1} \sqrt{x^2 + y^2}\, d \text{ length} - \int_{y_3}^{y_1} \frac{x^2}{2} d \text{ length} \\ &\ge \int_{y_3}^{y_1} (y - y^2/2) d \text{ length} \ge \frac{1}{3}|y_1 - y_3|^2. \end{aligned}$$

When $x_3 \neq 0$ the estimate is similar but first one needs to 'cut' the 'tip of the beak' where $x^2 \le y^2$ does not hold. When $y_3 < 0$ while $y_1 > 0$ or *vice versa*, modification in the same spirit is required. □

5. *A positive result*

Here we prove Theorem 2.

Let M be a Riemannian manifold and $d(\cdot, \cdot)$ the distance function induced by the Riemannian metric. Let $f\colon M \to M$ be an Anosov diffeomorphism and let $g\colon M \to M$ be a diffeomorphism Hölder conjugate to $\tilde{f}$:

$$h \circ \tilde{f} = f \circ h.$$

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

Let α be the Hölder exponent of h and β the Hölder exponents of h^{-1}. Recall that we assume that $\alpha\beta > 1/2$.

Definition 12. A sequence of points $\{y_i \in M;\ i \in \mathbb{Z}\}$ is called a ε *pseudo-orbit* for $g\colon M \to M$ if $d(g(y_i),\ y_{i+1}) < \varepsilon$, $i \in \mathbb{Z}$.

Definition 13. We say that a real orbit $\{f^i(x);\ i \in \mathbb{Z}\}$ δ-*shadows* a ε pseudo-orbit $\{y_i;\ i \in \mathbb{Z}\}$ if $d(f^i(x),\ y_i) < \delta$.

Definition 14. A diffeomorphism $g\colon M \to M$ is *quasi-Anosov* if for all non-zero $v \in TM$ the sequence $\{\|Dg^i v\|;\ i \in \mathbb{Z}\}$ is unbounded.

We need the following characterization of Anosov systems.

THEOREM 15. (Mather [**M77**]) *A diffeomorphism $g\colon M \to M$ is Anosov if and only if g is quasi-Anosov and all dimensions of stable manifolds at periodic points are the same.*

Dimensions of stable manifolds at periodic points of $\tilde{f}$ are the same since $\tilde{f}$ is topologically conjugate to f. Hence, we only need to show that $\tilde{f}$ is quasi-Anosov.

Assume that $\tilde{f}$ is not quasi-Anosov. Then there exist $v \in T_{\tilde{x}}M$, $\|v\| = 1$, such that $\|D\tilde{f}^n v\| \le 1$ for all $n \in \mathbb{Z}$. Define $v_n = D\tilde{f}^n v$, $n \in \mathbb{Z}$. For any sufficiently small $\varepsilon > 0$ consider a sequence $\{\tilde{x}_n = \exp(\varepsilon v_n);\ n \in \mathbb{Z}\}$. A sequence $\{\varepsilon v_n\}$ is ε-small while diffeomorphism $\tilde{f}$ is $C^{1+\mathrm{Lip}}$. This implies that there exists a constant $\tilde{c}$ that depends on $\tilde{f}$ only such that $\{\tilde{x}_n\}$ is a $\tilde{c}\varepsilon^2$ pseudo-orbit which is obviously δ-shadowed by the orbit of $\tilde{x}$ with $\varepsilon/2 < \delta < 2\varepsilon$.

Let $x = h(\tilde{x})$ and $x_n = h(\tilde{x}_n)$, $n \in \mathbb{Z}$. Applying Hölder inequalities we obtain that $\{x_n\}$ is a $c_1\varepsilon^{2\alpha}$ pseudo-orbit for f that is δ-shadowed by $\{f^n(x)\}$ with $\delta > c_2\varepsilon^{1/\beta}$. Constants c_1 and c_2 do not depend on ε. To make it more transparent let $\xi = c_1\varepsilon^{2\alpha}$.

For arbitrarily small $\xi > 0$ we have constructed a ξ pseudo-orbit and a true orbit that δ-shadows the pseudo-orbit with $\delta > c\xi^{\kappa} \stackrel{\text{def}}{=} \xi^{1/2\alpha\beta}$ where $\kappa < 1$ by the assumption. Meanwhile it is a well-known simple fact that δ can be estimated from above $\delta < C\xi$ where C depends only on f. The proof is straightforward and exploits local product structure of stable and unstable foliation of f. For ξ small enough these bounds on δ contradict each other. Hence, we have arrived at a contradiction.

Acknowledgements. Anatole Katok suggested to the author to look at the system with a heteroclinic tangency since this is the simplest situation when one can hope to obtain a counterexample. The author would like to thank A. Katok for discussions, encouragement and useful comments on the text itself. He also would like to thank M. Guysinsky for explaining the idea of the proof of Theorem 2.

REFERENCES

[BDV98] Ch. Bonatti, L. Diaz and F. Vuillemin. Topologically transverse nonhyperbolic diffeomorphisms at the boundary of the stable ones. *Bol. Soc. Bras. de Mat.* **29** (1998), 99–144.

[C98] M. Carvalho. First homoclinic tangencies in the boundary of Anosov diffeomorphisms. *Discrete Continuous Dynam. Syst. A* **4** (1998), 765–782.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press

[E98] H. Enrich. A heteroclinic bifurction of Anosov diffeomorphisms. *Ergod. Th. & Dynam. Sys.* **18** (1998), 567–608.

[F06] T. Fisher. *PhD Thesis*. Available online at http://www.etda.libraries.psu.edu/theses/approved/WorldWideFiles/ETD-1174/fisher-thesis-draft.pdf. PennState, 2006.

[K79] A. Katok. Bernoulli diffeomorphisms on surfaces. *Ann. of Math.* (2) **110**(3) (1979), 529–547.

[KH95] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge, 1995.

[KN08] A. Katok and V. Nitica. Differentiable rigidity of higher rank abelian group actions. (2008) in preparation.

[L80] J. Lewowicz. Lyapunov functions and topological stability. *J. Differential Equations* **38** (1980), 192–209.

[M77] R. Mañé. Quasi-Anosov diffeomorphisms and hyperbolic manifolds. *Trans. Amer. Math. Soc.* **229** (1977), 351–370.

[Math68] J. Mather. Characterization of Anosov diffeomorphisms. *Nederl. Akad. Wetensch. Proc. Ser. A* **30** (1968), 479–483.

[W70] P. Walters. Anosov diffeomorphisms are topologically stable. *Topology* **9** (1970), 71–78.

https://doi.org/10.1017/S0143385709000169 Published online by Cambridge University Press