Canad. J. Math. 2026, pp. 1–41
http://dx.doi.org/10.4153/S0008414X2510179X
© The Author(s), 2025. Published by Cambridge University Press on behalf of Canadian Mathematical Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

Computability of Brjuno-like functions

Michael Yampolsky and Ivan O. Shevchenko

Abstract. In his seminal paper from 1936, Alan Turing introduced the concept of non-computable real numbers and presented examples based on the algorithmically unsolvable Halting problem. We describe a different, analytically natural mechanism for the appearance of non-computability. Namely, we show that additive sampling of orbits of certain skew products over expanding dynamics produces Turing non-computable reals. We apply this framework to Brjuno-type functions to demonstrate that they realize bijections between computable and lower-computable numbers, generalizing previous results of M. Braverman and the second author for the Yoccoz–Brjuno function to a wide class of examples, including Wilton's functions and generalized Brjuno functions.

1 Introduction

In 1936, Alan Turing published a seminal paper [Tur36] which is rightly considered foundational for modern computer science. The main subject of his paper, as seen from the title, was the concept of a computable real number. Informally, such numbers can be approximated to an arbitrary desired precision using some algorithm. Turing formalized the latter as a Turing machine (TM), which is now the commonly accepted theoretical model of computation. Appearing before actual computers, TMs can be somewhat cumbersome to describe, but their computational power is equivalent to those of programs in a modern programming language, such as Python, for instance. Let

$$\mathbb{D} = \{p2^q,\ p, q \in \mathbb{Z}\}$$

denote the set of dyadic rationals. A number $x \in \mathbb{R}$ is *computable* if there exists a TM M, which has a single input $n \in \mathbb{N}$ and which outputs $d_n \in \mathbb{D}$ such that

$$|x - d_n| < 2^{-n}.$$

The dyadic notation here is purely to support the intuition that modern computers operate in binary; replacing 2^{-n} with any other constructive bound, such as, for instance, n^{-1}, would result in an equivalent definition.

Since there are only countably many programs in Python, there are only countably many computable numbers. Yet, it is surprisingly nontrivial to present an example of one. To this end, Turing introduced an algorithmically unsolvable problem, now

Received by the editors March 26, 2025; revised October 27, 2025; accepted November 3, 2025.
Published online on Cambridge Core November 12, 2025.
I.O.S. was partially supported by NSERC USRA fellowship. M.Y. was partially supported by NSERC Discovery grant.
AMS subject classification: **03D78**, **37E99**.
Keywords: Computability, Brjuno function, Wilton function.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

known as the Halting Problem: determine algorithmically whether a given TM halts or runs forever. Turing showed that there does not exist a program M whose input is another program M_1 and whose output is 1 if M_1 halts, and 0 if M_1 does not halt.

From this, a non-computable real is constructed as follows. Let us enumerate all programs

$$M_1, M_2, \dots, M_n, \dots$$

in some explicit way. For instance, all possible finite combinations of symbols in the Python alphabet can be listed in the lexicographic order. Some of these programs will not run at all (and thus halt by definition) but some will run without ever halting. Let the *halting predicate* $p(n)$ be equal to 1 if M_n halts and 0 otherwise.

Now, set

$$\alpha = \sum_{n \geq 1} p(n) 3^{-n}.$$

This number is clearly non-computable – an algorithm computing it could be used to determine the value of the halting predicate for every given n, and thus cannot exist.

It is worth noting a further property of α. Let us say that $x \in \mathbb{R}$ is *left-computable* if there exists a TM M which outputs an increasing sequence

$$a_n \nearrow x.$$

There are, again, only countably many left-computable numbers, and it is trivial to see that computable numbers form their subset (a proper subset, as seen below). Right computability is defined in the same way with decreasing sequences, and it is a nice exercise to show that being simultaneously left- and right-computable is equivalent to being computable.

The non-computable number α is left-computable, as it is the limit of

$$\alpha_n = \sum_{j \leq n \mid M_j \text{ halts in at most } n \text{ steps}} 3^{-j},$$

which can be generated by a program which emulates $M_1, \dots, M_k$ for k steps to produce α_k.

Since Turing, examples of non-computable reals were typically constructed along similar lines. In the 2000s, working on problems of computability in dynamics, M. Braverman and the second author discovered that non-computability may be produced via an analytic expression [BY09]. This expression came in the form of the *Brjuno function* $\mathcal{B}(\theta)$ which was introduced by Yoccoz [Yoc96] to study linearization problems of irrationally indifferent dynamics. We will discuss $\mathcal{B}(\theta)$ and its various cousins in detail below, but let us give an explicit formula here. For $\theta \in (0,1) \setminus \mathbb{Q}$, we have

$$\mathcal{B}(\theta) = \sum_{n \geq 0} \theta_{-1} \theta_0 \theta_1 \dots \theta_{n-1} \log \frac{1}{\theta_n}.$$

Here, $\theta_{-1} = 1$, $\theta_0 \equiv \theta$, and θ_{i+1} is obtained from θ_i by applying the Gauss map $x \mapsto \{1/x\}$. Of course, for rational values of θ, the summand will eventually turn infinite.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

The sum may also diverge for an irrational θ, and yet can be shown to converge to a finite value for almost all values in $(0,1)$.

It is evident that if θ is computable, then $\mathcal{B}(\theta)$ is left-computable. Very surprisingly, this can be reversed in the following way. Setting

$$y_* \equiv \inf\{\mathcal{B}(\theta),\ \theta \in (0,1)\},$$

for each left-computable $y \in [y_*, \infty)$, there exists a computable $\theta \in (0,1)$ with $\mathcal{B}(\theta) = y$. The Brjuno function can be seen as a "machine" mapping computable values in $(0,1)$ surjectively onto left-computable values in $[y_*, \infty)$.

Moreover, there exists an *explicit algorithm* $\hat{M}$ which, given a sequence $a_n \nearrow y$, computes θ such that $\mathcal{B}(\theta) = y$. Taking Turing's non-computable α, as we saw above, there is an explicit program to produce a sequence of rationals $\alpha_n \nearrow \alpha$. "Feeding" the sequence α_n to $\hat{M}$, we obtain an explicitly computable θ_* for which $\mathcal{B}(\theta_*) = \alpha$. This example demonstrates that $\mathcal{B}(\theta)$ is a natural analytic mechanism for producing non-computable reals.

The purpose of this article is as follows. We distill the proof of the above result from [BY09], where it is somewhat hidden in the considerations of complex dynamics and Julia sets. Moreover, we generalize the result to cover other Brjuno-type functions which have previously appeared in the mathematical literature, some of them 60 years before Yoccoz's work and in a completely different context. Our generalization describes the phenomenon in the language of dynamical systems. As we will see, ergodic sampling of a particular type of dynamics with suitable weights leads to non-computability.

Finally, for an even broader natural class of functions, which have also previously been studied, and which do not quite fit the above framework, we prove a more general, albeit weaker, non-computability result.

2 Preliminaries

2.1 Computable functions

The "modern" definition of a computable function requires the concept of an *oracle*. Loosely speaking, an oracle for a real number x, for example, is a user who *knows* x and, when queried by a TM, can input its value with any desired precision. In the world of TMs (as well as Python programs), an oracle may be conceived as an infinite tape on which an infinite string of dyadic rationals is written (encoding, for instance, a Cauchy sequence for $x \in \mathbb{R}$) and which the program is able to read at will. Of course, only a finite amount of information can be read off this tape each time. As well as a real number, an oracle can be used to encode anything else which could be written on an infinite tape, for instance, the magically obtained solution to the Halting Problem.

Formally, an oracle is a function $\phi : \mathbb{N} \to \mathbb{D}$. An *oracle for* $x \in \mathbb{R}$ satisfies

$$|\phi(n) - x| < 2^{-n} \text{ for all } n \in \mathbb{N}.$$

We say a TM M^ϕ is an *oracle Turing Machine* if at any step of the computation, M^ϕ can query the value $\phi(n)$ for any n. We treat an oracle TM as a function of the oracle; that is, we think of ϕ in M^ϕ as a placeholder for any oracle, and the TM performs its

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

computational steps depending on the particular oracle it is given. We will talk about "querying the oracle," "being given the access to an oracle for x," or just "given x."

We need oracle TMs to define computable functions on the reals. For $S \subset \mathbb{R}$, we say that *a function* $f : S \to \mathbb{R}$ *is computable* if there exists an oracle TM M^ϕ with a single input n such that for any $x \in S$, the following is true. If ϕ is an oracle for x, then upon input n, the machine M^ϕ outputs $d_n \in \mathbb{D}$ such that

$$|f(x) - d_n| < 2^{-n}.$$

In other words, there is an algorithm which can output the value of $f(x)$ with any desired precision if it is allowed to query the value of x with an arbitrary finite precision.

The domain of the real-valued function plays an important role in the above definition. The definition states that there is a single algorithm which, given x, works for every $x \in S$. We will abbreviate this by saying that f is *uniformly computable on* S.

In the case when S is a singleton, $S = \{x_0\}$, we will say that the function $f(x)$ is *computable at the point* x_0. Evidently, the weakest computability result and the strongest non-computability result one can obtain in regards to real-valued functions is when the domain is restricted to a single point.

It is worth making note of the following easy fact, whose proof we leave as an exercise.

Proposition 2.1 *If f is uniformly computable on S, then f is continuous on S.*

Uniform left- or right-computability of functions is defined in a completely analogous way.

2.2 Brjuno function and friends

Every irrational number θ in the unit interval admits a unique (simple) continued fraction expansion:

$$[a_1, a_2, a_3, \ldots] \equiv \cfrac{1}{a_1 + \cfrac{1}{a_2 + \cfrac{1}{a_3 + \ldots}}} \in (0,1)\backslash\mathbb{Q},$$

where $a_i \in \mathbb{N}$. An important related concept is that of the *Gauss map* $G : (0,1] \to [0,1]$ given by

$$G(x) = \left\{\frac{1}{x}\right\};$$

it has the property

$$G([a_1, a_2, a_3, \ldots]) = [a_2, a_3, a_4, \ldots].$$

In what follows, for a function F, we denote F^n its n-th iterate. For ease of notation, for each $j \in \mathbb{N}$, let us define the function $\eta_j : (0,1)\backslash\mathbb{Q} \to (0,1)\backslash\mathbb{Q}$ as $\eta_j(x) = G^{j-1}(x)$,

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

so that

$$\eta_j([a_1, a_2, a_3, \ldots]) = [a_j, a_{j+1}, a_{j+2}, \ldots].$$

We define *Yoccoz's Brjuno function*, or for brevity, just the *Brjuno function* [Yoc96] by

$$\mathcal{B}(x) = \sum_{i=1}^{\infty} \eta_0(x)\eta_1(x)\ldots\eta_{i-1}(x)\cdot(-\log(\eta_i(x))), \tag{2.1}$$

where we set $\eta_0(x) = 1$ for all x. Irrationals in $(0,1)$ for which $\mathcal{B}(x) < +\infty$ are known as *Brjuno numbers*; they form a full measure subest of $(0,1)$. The original work of Brjuno [Brj71] characterized Brjuno numbers using a different infinite series, whose convergence is equivalent to that of (2.1).

The Brjuno condition has been introduced in the study of the linearization of neutral fixed points. The function $\mathcal{B}$ has an important geometric meaning in this context as an estimate on the size of the domain of definition of a linearizing coordinate. It has a number of remarkable properties and has been studied extensively (see, for instance, [MMY97]).

Intuitively, the condition $\mathcal{B}(x) < +\infty$ is a Diophantine-type condition; if x is a Diophantine number, then it can be shown that the series (2.1) is majorized by a geometric series. As we have learned from a talk by Marmi [Mar22], similar expressions have appeared much earlier in the theory of Diophantine approximation. Notably, in 1933, Wilton [Wil33] defined the sums

$$\mathcal{W}_1(x) = \sum_{i=1}^{\infty} \eta_0(x)\eta_1(x)\ldots\eta_{i-1}(x)\cdot(-\log^2(\eta_i(x))), \tag{2.2}$$

$$\mathcal{W}_2(x) = \sum_{i=1}^{\infty} (-1)^{i+1}\cdot\eta_0(x)\eta_1(x)\ldots\eta_{i-1}(x)\cdot(-\log(\eta_i(x))), \tag{2.3}$$

which we will call *the first and second Wilton functions*, respectively.

To illustrate how different the application of Wilton's functions is from the Brjuno function, let us quote Wilton's results. If we denote $d(n)$ to be the number of divisors of a positive integer n, Wilton showed that

$$\sum_{n=1}^{\infty} \frac{d(n)}{n}\cos 2\pi nx < \infty \text{ if and only if } \mathcal{W}_1(x) < \infty, \text{ and}$$
$$\sum_{n=1}^{\infty} \frac{d(n)}{n}\sin 2\pi nx < \infty \text{ if and only if } \mathcal{W}_2(x) < \infty.$$

One important generalization of the simple continued fraction expansion of an irrational number is the *α-continued fraction expansion* for $\alpha \in [1/2, 1]$. Let $A_\alpha : [0,\alpha] \to [0,\alpha]$ be the map

$$A_\alpha(0) = 0, A_\alpha(s) = \left|\frac{1}{x} - \left\lfloor \frac{1}{x} - \alpha + 1 \right\rfloor\right|, x \neq 0.$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

By iterating this mapping, we define the infinite α-continued fraction expansion for any $x \in (0,\alpha) - \mathbb{Q}$ as follows. For $n \geq 0$, we let

$$x_0 = |x - \lfloor x - \alpha + 1 \rfloor|, a_0 = \lfloor x - \alpha + 1 \rfloor, \varepsilon_0 = 1,$$
$$x_{n+1} = A_\alpha(x_n) = A_\alpha^{n+1}(x), a_{n+1} = \left\lfloor \frac{1}{x_n} - \alpha + 1 \right\rfloor \geq 1, \varepsilon_{n+1} = \operatorname{sgn}(x_n).$$

Then, we can write

$$x = [(a_1,\varepsilon_1),(a_2,\varepsilon_2),\dots,(a_n,\varepsilon_n),\dots] := \cfrac{\varepsilon_0}{a_1 + \cfrac{\varepsilon_1}{a_2 + \cfrac{\varepsilon_2}{a_3 + \dots}}} \in (0,\alpha) - \mathbb{Q}.$$

Note that when $\alpha = 1$, we recover the standard continued fraction expansion.

Generalizations of the Brjuno function based on the above expansions have been studied, for example, in [LMNN10]. There, the authors considered the properties of the function

$$\mathcal{B}_{\alpha,u}(x) = \sum_{i=1}^{\infty} \eta_{\alpha,0}(x)\eta_{\alpha,1}(x)\dots\eta_{\alpha,i-1}(x)\cdot u(\eta_{\alpha,i}(x)), \tag{2.4}$$

where $\eta_{\alpha,0}(x) = 1$, $\eta_{\alpha,j}(x) = A_\alpha^{j-1}(x)$, $\alpha \in [1/2,1]$, $j \geq 1$ is a generalization of η_j to alpha-continued fraction maps, and $u : (0,1) \to \mathbb{R}^+$ is a $\mathcal{C}^1$ function such that

$$\lim_{x\to 0^+} u(x) = \infty, \lim_{x\to 0^+} x\cdot u(x) < \infty, \lim_{x\to 0^+} x^2\cdot u'(x) < \infty.$$

A further generalized class $\{\mathcal{B}_{\alpha,u,v}\}$ of Brjuno functions is discussed in [BCM24], where the last two conditions above are dropped and the term $\eta_{\alpha,1}(x)\eta_{\alpha,2}(x)\dots\eta_{\alpha,i-1}(x)$ is raised to some power $v \in \mathbb{Z}^+$:

$$\mathcal{B}_{\alpha,u,v}(x) = \sum_{i=1}^{\infty} (\eta_{\alpha,0}(x)\eta_{\alpha,1}(x)\dots\eta_{\alpha,i-1}(x))^v \cdot u(\eta_{\alpha,i}(x)). \tag{2.5}$$

As shown in [BCM24]:

Proposition 2.2 *For all $\alpha \in \mathbb{Q}\cap[1/2,1]$, such functions $\mathcal{B}_{\alpha,u,v}$ are lower semi-continuous and thus attain their global minima.*

Note that if we take $\alpha = 1$, $v = 1$, and $u = -\log$, we recover Yoccoz's Brjuno function discussed above; similarly, taking $\alpha = 1$, $v = 1$, and $u = -\log^2$ recovers the first Wilton function $\mathcal{W}_1$. Note, however, that we cannot obtain $\mathcal{W}_2$ as a special case of $\mathcal{B}_{\alpha,u,v}$.

3 Statements of the results

3.1 A general framework

We refer the readers to the survey [MMY06] which discusses a cohomological interpretation of the Brjuno function and lays the groundwork for its generalizations. Our discussion will be much less technical, yet essentially equivalent in the cases we consider. It will yield a generalization which is (a) broad enough to include the relevant examples we have quoted and (b) captures the essence of the non-computability

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

phenomenon discovered in [BY09]. This is achieved via the following framework. Suppose $G(x)$ is a piecewise-defined expanding mapping whose domain is an infinite collection of subintervals of $(0,1)$, each of which is mapped surjectively over all of $(0,1)$. Let $\psi : (0,1) \to \mathbb{R}$ (in our case, $\psi(x) = x^\nu$), and consider the skew product dynamics given by

$$F : \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} G(x) \\ \psi(x)y \end{pmatrix}. \tag{3.1}$$

The class of functions $\mathcal{F}$ for which non-computability arises is produced by additive (ergodic) sampling of orbits (x_n, y_n) of F using a suitable weight function $u(x,y) \equiv u(x)$ with positive values:

$$\mathcal{F}(x) = \sum_{i=1}^{\infty} y_n \cdot u(x_n). \tag{3.2}$$

It is worth noting that such a function is a formal solution of the twisted cohomological equation

$$\mathcal{F} - \psi \cdot \mathcal{F} \circ G = u.$$

As noted above, in our case, we set

$$\psi(x) \equiv x^\nu.$$

We will now need to make somewhat technical but straightforward general assumptions on G and u.

Let $G : (0,1) \to (0,1)$ be a function which is $\mathcal{C}^1$ on a set $S \subseteq (0,1)$, with $s_0 := \inf S$, $s_1 := \sup S$. Suppose that for a countable collection of disjoint open intervals $J_i = (\ell_i, r_i)$ with $\ell_1 > \ell_2 > \cdots$, we have $S = J_1 \sqcup J_2 \sqcup J_3 \sqcup \cdots$. Below, we will denote $G_i := G|_{J_i}$, so that $G_i^{-1} : (s_0, s_1) \to J_i$ is the unique branch of G^{-1} mapping into J_i. We assume that G satisfies the following criteria:

(i) $G(J_i) = (s_0, s_1)$ for each i.

(ii) $|G'| > 1$, and additionally there exist $\tau > 1$, $\sigma > 1$, and $\kappa \in \mathbb{Z}$ such that if we let

$$\tau_{i,1} := \inf_{x \in J_i} |G'(x)|, \tau_{i,\kappa} := \inf_{x \in J_i} |(G^\kappa)'(x)|,$$

then we have both $\tau_{i,1}^{-1} < \ell_i \cdot \sigma$ and $\tau_{i,\kappa}^{-1} < \ell_i \cdot \tau^{-1}$ for all i. In particular, since $\ell_i \le s_1 < 1$, this means that $|(G^\kappa)'(x)| > \tau$ for all $x \in (s_0, s_1)$.

(iii) G_1 is decreasing on J_1.

(iv) Let φ be the unique fixed point of G_1, and let $\delta_G(N) := G_N^{-1}(\varphi) - G_{N+1}^{-1}(\varphi) > 0$. Then, there is some constant $D > 0$ such that for all $N \in \mathbb{Z}^+$,

$$\frac{r_{N+1}}{\ell_{N+1}} \cdot \frac{r_N - \ell_{N+1}}{\delta_G(N)} < D.$$

(v) We have $\dfrac{r_i - \ell_i}{\ell_i^2} < D$ for some constant $D > 0$ independent of i.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

(vi) G is computable on its domain.

(vii) $g(i) := \dfrac{r_i}{\ell_{i+1}} - 1 \to 0$ as $i \to \infty$. Note that since g is positive and bounded from above, there is some constant $m_g > 0$ for which $0 < g(i) \le m_g$.

It is easy to show that properties (i) and (ii) imply that G restricted to the set $\Lambda = \bigcap_{j=0}^{\infty} G^{-j}((s_0, s_1))$ is topologically conjugate to the full shift over the alphabet of positive integers $\mathbb{Z}^+$. As such, we can write any $x \in \Lambda$ as its symbolic representation $x = [a_1, a_2, \ldots]$, where $G^{j-1}(x) \in J_{a_j}$. In this notation, φ above can be written as $[1, 1, 1, \ldots]$. By (iii), we have $\varphi < s_1$, from which it is straightforward to check that we indeed have $\delta_G(N) > 0$ in (iv). For convenience, for $j \in \mathbb{Z}^+$, we will denote $\eta_j : \Lambda \to \Lambda$, $\eta_j = G^{j-1}$, so that

$$\eta_j([a_1, a_2, \ldots]) = [a_j, a_{j+1}, \ldots].$$

Henceforth, we will assume that G is restricted to Λ.

Now, let $u : (s_0, s_1) \to \mathbb{R}^+$ be a $\mathcal{C}^1$ function satisfying the following:

(i) $\lim_{x \to s_0^+} u(x) = \infty$.

(ii) If $s_1 = 1$, then $\liminf_{N \to \infty} \inf_{z, w \in (s_0, s_1)} \dfrac{u \circ G_1^{-1} \circ G_N^{-1}(z)}{u \circ G_1^{-1} \circ G_N^{-1}(w)} > 0$.

(iii) There is some $C > 0$ such that $|u'(x)| < \dfrac{C}{(x - s_0)^2}$ for all $x \in (s_0, s_1)$.

(iv) u is computable on numbers $[a_1, a_2, \ldots]$ such that $a_l = 1$ for all $l > l_0$ for some integer l_0.

(v) u is left-computable on all of Λ.

In what follows, for any $v > 0$, we define the *generalized Brjuno function* to be

$$\Phi(x) := \sum_{i=1}^{\infty} (\eta_0(x) \ldots \eta_{i-1}(x))^v \cdot u(\eta_i(x)),$$

where $x = [a_1, a_2, \ldots] \in \Lambda$.

We note:

Theorem 3.1 *The following functions restricted to their corresponding sets Λ fall under the definition of a generalized Brjuno function given above:*

- *Yoccoz's Brjuno function $\mathcal{B}$ (2.1).*
- *The first Wilton function $\mathcal{W}_1$ (2.2).*
- *The functions $\mathcal{B}_{\alpha,u,v}$ (2.5) under the additional assumptions of (ii)–(v) on u. For example, taking $G = A_\alpha$ for $\alpha \in [1/2, 1]$ and $u(x)$ to be any of $\log^n(1/x)$ for $n \in \mathbb{Z}$ or x^{-1} yields a generalized Brjuno function.*

Let us postpone the proof to Section A.1 in the Appendix, and proceed to formulating the results.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

3.2 Main results

Theorem 3.2 *Let x_* be a computable real number in Λ with the property $y_* = \Phi(x_*) < +\infty$. There exists an oracle TM M^ϕ with a single input $n \in \mathbb{N}$ such that the following holds. Suppose $y \in [y_*, +\infty)$ and*

$$\phi(n) = y_n \nearrow y.$$

Then, M^ϕ outputs d_n such that

$$|d_n - x| < 2^{-n} \text{ for } x \in \Lambda \text{ such that } \Phi(x) = y.$$

As a corollary:

Corollary 3.3 *If $y \in [y_*, +\infty)$ is left computable, then there exists a computable $x \in \Lambda$ with*

$$\Phi(x) = y.$$

In fact, it is clear from the proof of Theorem 3.2 that countably many such $x \in \Lambda$ exist.

As was shown in [BM12], the Brjuno function $\mathcal{B}$ attains its global maximum at

$$w_* = \frac{\sqrt{5}-1}{2} = [1, 1, 1, \ldots].$$

Therefore:

Corollary 3.4 *If $y \in [\mathcal{B}(w_*), +\infty)$ is left computable, then there exists a computable $x \in (0,1)$ with*

$$\mathcal{B}(x) = y.$$

4 Proof of Theorem 3.2

4.1 Three main lemmas

It will be helpful to first outline the general strategy for the proof of Theorem 3.2. We are given a computable sequence $\{y_n\}$ converging upward to some left-computable y, and we must find a computable $x \in (s_0, s_1)$ for which $y = \Phi(x)$. This is done by starting with some $\gamma_0 \in (s_0, s_1)$ and iteratively modifying the symbolic representation of γ_k to "squeeze" $\Phi(\gamma_k)$ to be in the interval $(\Phi(\gamma_{s+k}) - 2^{-k}\varepsilon, \Phi(\gamma_{s+k}) + 2^{-k}\varepsilon)$ for some positive integer s. Passing to the limit, we obtain $x = \gamma_\infty$ for which $\Phi(x) = y$ as needed.

To ensure this strategy works, we need ways of carefully controlling the value of $\Phi(\gamma)$ from the symbolic representation of γ. This role is played by Lemmas 4.1–4.3 below, which are analogous to Lemmas 5.18–5.20, respectively, in [BY09] and whose proofs are in the Section A.2 in the Appendix.

Lemma 4.1 *For any initial segment $I = [a_1, a_2, \ldots, a_n]$, write $\omega = [a_1, a_2, \ldots, a_n, 1, 1, 1, \ldots]$. Then, for any $\varepsilon > 0$, there is an $m > 0$ and an integer*

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

N such that if we write $\beta^N = [a_1, a_2, \ldots, a_n, 1, 1, \ldots, 1, N, 1, 1, \ldots]$, where the N is located in the $(n+m)$-th position, then

$$\Phi(\omega) + \varepsilon < \Phi(\beta^N) < \Phi(\omega) + 2\varepsilon.$$

In the Appendix, this lemma will be proven under the slightly weaker assumptions used in Section 5. Also, it is clear from the proof that m can be taken arbitrarily large.

Lemma 4.2 *With ω as above, for any $\varepsilon > 0$, there is an $m_0 > 0$, which can be computed from $(a_0, a_1, \ldots, a_n)$ and ε, such that for any $m \geq m_0$ and any tail $I = [a_{n+m}, a_{n+m+1}, \ldots]$, we have*

$$\Phi(\beta^I) > \Phi(\omega) - \varepsilon,$$

where

$$\beta^I = [a_1, a_2, \ldots, a_n, 1, 1, \ldots, 1, a_{n+m}, a_{n+m+1}, \ldots].$$

Lemma 4.3 *Let $\omega = [a_1, a_2, \ldots]$ be such that $\Phi(\omega) < \infty$. Write $\omega_k = [a_1, a_2, \ldots, a_k, 1, 1, \ldots]$. Then, for every $\varepsilon > 0$, there is an m such that for all $k \geq m$,*

$$\Phi(\omega_k) < \Phi(\omega) + \varepsilon.$$

We proceed with proving Theorem 3.2.

4.2 The proof

We first require a few preliminary results.

Proposition 4.4 *Let $m_0 > 0$ be an integer. Then, there exists an oracle TM which, given access to a number $x = [a_1, a_2, \ldots] \in \Lambda$ such that $a_m = 1$ for $m > m_0$, computes $\Phi(x)$.*

Proof Let $\varphi = [1, 1, 1, \ldots]$. For any x, we have

$$\begin{aligned}\Phi(x) &= \sum_{i=1}^{m_0} (\eta_0(x) \ldots \eta_{i-1}(x))^\nu \cdot u(\eta_i(x)) + \sum_{i=m_0+1}^{\infty} (\varphi^\nu)^{n-1} \cdot u(\varphi) \\ &= \sum_{i=1}^{m_0} (\eta_0(x) \ldots \eta_{i-1}(x))^\nu \cdot u(\eta_i(x)) + \frac{\varphi^{\nu \cdot m_0}}{1 - \varphi^\nu} \cdot u(\varphi).\end{aligned}$$

Since u is computable on each number $\eta_i(x)$ whose symbolic representation ends in all ones by assumption (iv) on u, and each η_j is computable on all of Λ by assumption (vi) on G, there is an oracle TM which, given access to x, computes the sum on the left to an arbitrary precision. The construction of this oracle TM is independent of x and depends only on m_0. Additionally, since φ also ends in all ones, there is a TM which computes the value of the term on the right to an arbitrary precision, also depending only on m_0. Combining these TMs in the obvious way, we obtain an oracle TM which, given access to x, computes $\Phi(x)$ to an arbitrary precision. ∎

Lemma 4.5 *Given an initial segment $I = [a_0, a_1, \ldots, a_n]$ and $m_0 > 0$, write $\omega = [a_0, a_1, \ldots, a_n, 1, 1, \ldots]$. Then, for all $\varepsilon > 0$, we can uniformly compute $m > m_0$ and $t, N \in \mathbb{Z}^+$ such that if we write $\beta = [a_0, a_1, \ldots, a_n, 1, 1, \ldots, 1, N, 1, 1, \ldots]$, where N is in*

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

the $(n+m)$*-th position, we have*

$$\Phi(\omega) + \varepsilon < \Phi(\beta) < \Phi(\omega) + 2\varepsilon, \tag{4.1}$$

and for any $\gamma = [a_0, a_1, \ldots, a_n, 1, 1, \ldots, 1, N, 1, \ldots, 1, c_{n+m+t+1}, c_{n+m+t+2}, \ldots]$*, we have*

$$\Phi(\gamma) > \Phi(\omega) - 2^{-n}. \tag{4.2}$$

Proof We will first show that such m, N exist, then give an algorithm to compute them. Let $\varepsilon > 0$ be given. By Lemma 4.1, there exists m (and we can make $m > m_0$) and $\exists N \in \mathbb{Z}^+$ such that

$$\Phi(\omega) + \varepsilon < \Phi(\beta) < \Phi(\omega) + 2\varepsilon.$$

Taking $I' = [a_0, a_1, \ldots, a_n, 1, 1, \ldots, 1, N]$ and $\omega' = \beta$, applying Lemma 4.2 with $\varepsilon' = 2^{-n}$, we get $t_0 > 0$, which can be computed from $(a_0, a_1, \ldots, a_n, 1, 1, \ldots, 1, N)$ such that for all $t \geq t_0$ and any tail $I = [c_{n+m+t+1}, c_{n+m+t+2}, \ldots]$, we have

$$\Phi(\gamma) > \Phi(\omega') - 2^{-n} = \Phi(\beta) - 2^{-n} > \Phi(\omega) - 2^{-n}$$

as needed.

Since ω, β have symbolic representations ending in all ones, for any specific m, N, we can compute $\Phi(\omega)$ and $\Phi(\beta)$ by Proposition 4.4. So, we can find the required m, N by enumerating all pairs (m, N) and exhaustively checking equations (4.1) and (4.2) for each of them. We know that we will eventually find a pair for which these equations hold. Once we have m and N, we can use Lemma 4.2 to compute t. ∎

Lemma 4.6 *The infimum* $\Phi(x_*)$ *of* $\Phi(x)$ *over all* $x \in \Lambda$ *is equal to the infimum over the numbers whose symbolic representations have only finitely many terms that are not* 1*:*

$$\Phi(x_*) = \inf_{x=[a_1, a_2, \ldots, a_k, 1, 1, \ldots]} \Phi(x).$$

Proof Let $\varepsilon > 0$ be given. By definition of infimum, there exists $x = [a_1, a_2, \ldots]$ such that

$$\Phi(x) < \Phi(x_*) + \frac{\varepsilon}{2}.$$

Write $x_k = [a_1, a_2, \ldots, a_k, 1, 1, \ldots]$. By Lemma 4.3, there exists m such that for $k \geq m$,

$$\Phi(x_k) < \Phi(x) + \frac{\varepsilon}{2}.$$

Thus, $\Phi(x_k) < \Phi(x_*) + \varepsilon$, so we can make $\Phi(x_k)$ as close to $\Phi(x_*)$ as we need. ∎

Now, we are given

$$y_n \nearrow y, \qquad y \in [y_*, +\infty).$$

The case of $y = y_*$ is trivial, so we suppose $y > y_*$. Then, there is an s and an $\varepsilon > 0$ such that

$$y_s > \Phi(x_*) + 2\varepsilon.$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

By Lemma 4.6, there exists $\gamma_0 = [a_1, a_2, \ldots, a_n, 1, 1, \ldots]$ such that

$$y_s - \varepsilon < \Phi(\gamma_0) < y_s - \frac{\varepsilon}{2}.$$

We will now give an algorithm for computing a number $x \in \Lambda$ for which $\Phi(x) = \lim \nearrow y_n = y$, which would complete the proof of Theorem 3.2. The algorithm works as follows. At state k, it produces a finite initial segment $I_k = [a_0, \ldots, a_k]$ such that the following properties hold:

(1) $I_0 = [a_1, a_2, \ldots, a_n]$.

(2) I_k has at least k terms, i.e., $m_k \geq k$.

(3) For each k, I_{k+1} is an extension of I_k.

(4) For each k, define $\gamma_k = [I_k, 1, 1, \ldots]$. Then,

$$y_{s+k} - 2^{-k}\varepsilon < \Phi(\gamma_k) < y_{s+k} - 2^{-(k+1)}\varepsilon.$$

(5) For each k, $\Phi(\gamma_k) > \Phi(\gamma_{k+1})$.

(6) For each k and any extension $\beta = [I_k, b_{m_k+1}, b_{m_k+2}, \ldots]$, we have

$$\Phi(\beta) > \Phi(\gamma_k) - 2^{-k}.$$

The first three properties are easy to verify. The last three are checked using Lemma 4.5. By this lemma, we can increase $\Phi(\gamma_{k-1})$ by any given amount, possibly in more than one step, by extending I_{k-1} to I_k. Thus, if we have

$$y_{s+k-1} - 2^{-(k-1)}\varepsilon < \Phi(\gamma_{k-1}) < y_{s+k-1} - 2^{-k}\varepsilon,$$

by virtue of $\{a_{s+k}\}_{k=1}^{\infty}$ being non-decreasing, we have both

$$y_{s+k-1} - 2^{-(k-1)}\varepsilon < y_{s+k} - 2^{-k}\varepsilon \quad \text{and} \quad y_{s+k-1} - 2^{-k}\varepsilon < y_{s+k} - 2^{-(k+1)}\varepsilon.$$

So, we can increase $\Phi(\gamma_{k-1})$ by such a fine amount that

$$y_{s+k} - 2^{-k}\varepsilon < \Phi(\gamma_k) < y_{s+k} - 2^{-(k+1)}\varepsilon,$$

satisfying the fourth and fifth properties. In performing this fine increase, we have used the fact that the y_{s+k}'s are computable. The last property is satisfied by Lemma 4.5 (4.2).

Denote

$$x = \lim_{k \to \infty} \gamma_k.$$

The symbolic representation of x is the limit of the initial segments I_k. This algorithm gives us at least one term of the symbolic representation of x per iteration, and hence we would need at most $O(n)$ iterations to compute x with precision 2^{-n}. The initial segment of γ_0 can also be computed as in the proof of Lemma 4.5. It remains to show that x is the number we are after.

Lemma 4.7 *We have* $\Phi(x) = y$.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Proof Taking limits on all sides of (4), we get

$$\lim_{k\to\infty} \Phi(\gamma_k) = \lim_{k\to\infty} y_k = y.$$

It remains to show $\lim_{k\to\infty} \Phi(\gamma_k) = \Phi(x)$. As in Proposition 4.4, denote $x = [a_1, a_2, \ldots]$ and let $\alpha_k = [a_k, a_{k+1}, \ldots]$, $\alpha_0 = 1$. Let $\varphi = [1, 1, 1, \ldots]$, and additionally for any number $\xi = [b_1, b_2, \ldots] \in \Lambda$ denote $(\xi)_k = [b_k, b_{k+1}, \ldots]$. We have

$$\begin{aligned}\Phi(x) &= \lim_{k\to\infty} \left(\sum_{n=1}^{m_k} \alpha_0\alpha_1 \ldots \alpha_{n-1} \cdot u(\alpha_n) \right) \\ &\le \lim_{k\to\infty} \left(\sum_{n=1}^{m_k} \alpha_0\alpha_1 \ldots \alpha_{n-1} \cdot u(\alpha_n) + \sum_{n=m_k+1}^{\infty} \varphi^{n-1} \cdot u(\varphi) \right) \\ &= \lim_{k\to\infty} \Phi(\gamma_k).\end{aligned}$$

Additionally, taking limits on both sides of (6) with $\beta = x$ yields

$$\lim_{k\to\infty} \Phi(\gamma_k) \le \Phi(x),$$

therefore $\lim_{k\to\infty} \Phi(\gamma_k) = \Phi(x)$. ∎

This concludes the proof of Theorem 3.2.

5 Generalized non-computability result

5.1 Modified assumptions

We will now prove a non-computability result about a slightly more broad class of generalized Brjuno functions than the one considered in Section 5. For this result, we require all the assumptions on G except (v) and (vi), and on u we require only assumption (i) together with a slightly weaker variation of assumption (iii):

(iii′) There is some $C > 0$ such that $u'(x) < \dfrac{C}{(x - s_0)^2}$ for all $x \in (s_0, s_1)$.

We can additionally allow for more flexibility in the definition of the generalized Brjuno function by adding a "sign" term:

$$\Phi(x) := \sum_{i=1}^{\infty} s(i) \cdot (\eta_0(x) \ldots \eta_{i-1}(x))^{\nu} \cdot u(\eta_i(x)),$$

where $\nu > 0$ as before and $s(i) \in \{-1, 1\}$.

Theorem 5.1 *The following functions restricted to their corresponding sets Λ fall under the definition of a generalized Brjuno function given above:*

- *Yoccoz's Brjuno function $\mathcal{B}$ (2.1).*
- *The first Wilton function $\mathcal{W}_1$ (2.2) and the second Wilton function $\mathcal{W}_2$ (2.3).*
- *The functions $\mathcal{B}_{\alpha,u,\nu}$ (2.5) under the additional assumption of (iii′) on u. As before, taking $G = A_\alpha$ for $\alpha \in [1/2, 1]$ and $u(x)$ to be any of $\log^n(1/x)$ for $n \in \mathbb{Z}$ or x^{-1} yields a generalized Brjuno function.*

The proof is completely analogous to the proof of Theorem 3.1 and will be omitted.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

5.2 Noncomputability result

We proceed with the main result of this section, which concerns the computability of the function Φ as opposed to the computability of real numbers.

Theorem 5.2 *There exists a number $x \in \Lambda$ for which $\Phi(\cdot)$ as defined above is not computable at x.*

Without loss of generality, we will assume in what follows that the sign term $s(i) = 1$ infinitely often (otherwise, just replace Φ with $-\Phi$).

As before, it will be instructive to first go over the strategy of the proof. This outline is rough and is not fully logically sound, however, it captures the main idea of the argument. To prove a function is non-computable at a single point x, it suffices to enumerate all oracle TMs M_i^ϕ, $i \in \mathbb{N}$ (recall that there are countably many oracle TMs), and show that if ϕ is any oracle of x, then M_i^ϕ does not approximate $\Phi(x)$ arbitrarily well.

We start with $x_0 = [1, 1, 1, \ldots]$ and the first TM $M_{n_1}^\phi$ in our enumeration which computes x_0. If any of the digits a_j in the symbolic representation of x_0 are changed to some $N \in \mathbb{Z}^+$, as $N \to \infty$, the series $\Phi(x)$ diverges. However, if we change some digit a_j far enough in the representation of x_0, for any N, the new value of x_0 changes by at most some fixed small amount ε_j which goes to 0 as $j \to \infty$. So, the idea is to define x_1 from x_0 by changing a_{j_1} for large enough j_1 to some large enough N_1, such that if $M_{n_1}^\phi$ is given an oracle for x_1, then it does not properly compute $\Phi(x_1)$, which in some sense "fools" the oracle TM $M_{n_1}^\phi$. To fool the machine $M_{n_2}^\phi$, we then change a digit $j_2 > j_1$ sufficiently far in the symbolic representation of x_1 to a large N_2 to get x_2, in such a way that neither $M_{n_2}^\phi$ nor any other M_k^ϕ for $k < n_2$ properly compute $\Phi(x_2)$. Continuing in this manner, we will arrive at a limiting number $x_\infty \in \Lambda$, with $\Phi(x_\infty) < \infty$ and such that none of the oracle TMs M_i^ϕ in our list properly compute x_∞.

As in the proof of Theorem 4.3, we will need to carefully control the value of $\Phi(x)$ from the symbolic expansion of x. For the proof of Theorem 5.2, we only need Lemma 4.1 from the previous section, which is proven in the Appendix under the weaker assumptions on Φ used in this section.

For the below proofs, we will say that $\Phi(x)$ is computable at x if there exists a TM M^ϕ such that if ϕ is an oracle for x, then on input n, M^ϕ outputs some y' for which $|\Phi(x) - y'| \le 2^{-n}$. This definition uses "$\le$" instead of the "$<$" which is used in the definition given in Section 2, but it is easy to see that the two definitions are equivalent.

Before starting the proof, we need the following elementary fact.

Lemma 5.3 *Write any number in Λ as $\omega = [a_1, a_2, \ldots]$. For any $\varepsilon > 0$, there is an $L > 0$ for which $n > L$ implies that for any sequence of natural numbers $(N_0, N_1, \ldots)$,*

$$|\omega - [a_1, a_2, \ldots, a_{n-1}, N_0, N_1, \ldots]| < \varepsilon.$$

Proof Let $M > 0$ be large enough that $\dfrac{s_1 - s_0}{\tau^M} < \varepsilon$ and let $L = \kappa \cdot M$, where κ, τ are from assumption (ii) on G. Noting that

$$\omega, [a_1, a_2, \ldots, a_{n-1}, N_0, N_1] \in (G_{a_1}^{-1} \circ G_{a_2}^{-1} \circ \cdots \circ G_{a_{n-1}}^{-1})((s_0, s_1))$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

and $|(G_{a_{i+\kappa-1}} \circ G_{a_{i+\kappa-2}} \circ \cdots \circ G_{a_i})'(\theta)| > \tau > 1 \implies |(G_{a_i}^{-1} \circ G_{a_{i+1}}^{-1} \circ \cdots \circ G_{a_{i+\kappa-1}}^{-1})'(\theta)| < \frac{1}{\tau} < 1$, as well as $|(G_{a_i}^1)'(\theta)| < 1$ from (ii), we have

$$\begin{aligned}
&\operatorname{length}((G_{a_1}^{-1} \circ \cdots \circ G_{a_{n-1}}^{-1})((s_0, s_1))) \\
&\quad \le \left(\prod_{i=0}^{M-1} \sup_{\theta \in \Lambda} |(G_{a_{i\kappa+1}}^{-1} \circ \cdots \circ G_{a_{i\kappa+\kappa}}^{-1})'(\theta)| \right) \cdot \left(\prod_{j=L+1}^{n-1} \sup_{\theta \in \Lambda} |(G_{a_j}^{-1})'(\theta)| \right) \cdot (s_1 - s_0) \\
&\quad < \frac{s_1 - s_0}{\tau^M} < \varepsilon.
\end{aligned}$$

■

In particular, we have the following.

Corollary 5.4 *For $\omega = [a_1, a_2, \ldots]$ as above, for any $\varepsilon > 0$, there is an $L > 0$ for which $n > L$ implies that $\forall N \in \mathbb{N}$,*

$$|\omega - [a_1, a_2, \ldots, a_{n-1}, N, a_{n+1}, \ldots]| < \varepsilon.$$

Before proceeding to the proof of the main result, we first define some notation. For any $x_i = [a_1^i, a_2^i, \ldots]$, let $\eta_k^i = [a_k^i, a_{k+1}^i, \ldots]$, noting that

$$\Phi(x_i) = \sum_{n=1}^{\infty} s(i) \cdot \left(\eta_0^i \eta_1^i \ldots \eta_{n-1}^i\right)^{\nu} \cdot u(\eta_n^i),$$

where $\nu, \rho > 0$ and $s(i) \in \{-1, 1\}$ with $s(i) = 1$ infinitely often. Let

$$f(i,k) = \sum_{n=1}^{k} s(i) \cdot \left(\eta_0^i \eta_1^i \ldots \eta_{n-1}^i\right)^{\nu} \cdot u(\eta_n^i),$$

noting that $\lim_{k\to\infty} f(i,k) = \Phi(x_i)$.

5.3 Proof of Theorem 5.2

We will first show inductively that there exist:

- nested initial segments $I_1 \subseteq I_2 \subseteq \ldots$, where each I_i has length p_i;
- for each $i = 1, 2, \ldots$, positive integers N^i and m_i;
- positive integers $k_1 < k_2 < \cdots$ and $l_1 < l_2 < \cdots$ and $\hat{a}_1 < \hat{a}_2 < \cdots$ and $n(0) < n(1) < n(2) < \cdots$, positive real numbers $\hat{\varepsilon}_1 > \hat{\varepsilon}_2 > \cdots$, and oracles $\phi_0, \phi_1, \ldots$;

such that if we let $x_i = [I_i, 1, \ldots, 1, N^i, 1, \ldots]$ for $i \in \mathbb{Z}^+$, where N^i is in the $(m_i + p_i)$-th position, then we have the following:

(1) ϕ_i is an oracle for x_i such that $|\phi_i(n) - x_i| < 2^{-(n+1)}$ for all n.

(2) ϕ_i agrees with the oracle ϕ_{j-1} on inputs $1, 2, \ldots, k_j$ for $j = 1, 2, \ldots, i$.

(3) Running $M_{l_i}^{\phi_i}(1), M_{l_i}^{\phi_i}(2), \ldots, M_{l_i}^{\phi_i}(\hat{a}_i)$ queries ϕ_i with parameters not exceeding k_i.

(4) Running $M_{l_i}^{\phi_i}(\hat{a}_i)$ yields a number A_{l_i} for which

$$A_{l_i} + 2^{-\hat{a}_i} \le \Phi(x_{i-1}) + 2^{-\hat{a}_i+1} < \Phi(x_i) < \Phi(x_{i-1}) + 2 \cdot 2^{-\hat{a}_i+1}.$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

(5) Running $M_{l_j}^{\phi_i}(\hat{a}_j)$ for $j = 1, 2, \ldots, i$ yields a number B_{l_j} for which $B_{l_j} + 2^{-\hat{a}_j} < \Phi(x_i)$.

(6) The TMs $M_k^{\phi_i}$ for $k \in \{1, \ldots, l_i\}$ all do not properly compute $\Phi(x_i)$; in particular, they all compute $\Phi(x_i)$ with an error of at least $\hat{\varepsilon}_i$.

(7) For $k \geq n(i)$, we have $|f(i,k) - \Phi(x_i)| < 2^{-i}$.

(8) For $k = 1, 2, \ldots, n(i-1)$, we have $|f(i,k) - f(i-1,k)| < 2^{-i}$.

5.3.1 Base case

There are countably many oracle TMs M^{ϕ}, where ϕ represents an oracle for x, so we can order them as $M_1^{\phi}, M_2^{\phi}, \ldots$. Let $x_0 = [1,1,1,\ldots]$. Given an oracle ϕ_0 for x_0 such that $|\phi_0(n') - x_0| < 2^{-(n'+1)}$ for all n', let $M_{l_1}^{\phi_0}$ be the first TM to compute $\Phi(x_0)$ (if no such TM exists, we are done). Since this is the first such TM, all of $M_1^{\phi_0}, M_2^{\phi_0}, \ldots, M_{l_1-1}^{\phi_0}$ do not properly compute $\Phi(x_0)$, so there are integers $a_1, a_2, \ldots, a_{l_1-1}$ and small positive real numbers $\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_{l_1-1}$ for which $M_k^{\phi_0}(a_k)$ outputs some number A_k with $|A_k - \Phi(x_0)| > 2^{-a_k} + \varepsilon_k$. Set $\hat{\varepsilon}_1 = \min(\varepsilon_1, \ldots, \varepsilon_{l_1-1})$; choose $\hat{n}_1$ large enough so that $2^{-\hat{n}_1+2} < \hat{\varepsilon}_1/2$, and set $\hat{a}_1 = \max(a_1, \ldots, a_{l_1-1}, \hat{n}_1)$.

Run $M_{l_1}^{\phi_0}(\hat{a}_1)$ with the oracle ϕ_0. This TM outputs a number A_{l_1} for which $|A_{l_1} - \Phi(x_0)| \leq 2^{-\hat{a}_1}$. Since the computation is performed in finite time, there is a $k_1 > 0$ such that ϕ_0 is only queried with parameters not exceeding k_1. We assume k_1 is large enough that the computations $M_{l_1}^{\phi_0}(k)$ query ϕ_0 for parameters not exceeding k_1 for $k = 1, 2, \ldots, \hat{a}_1 - 1$. Setting $n(0) = 1$, we additionally make k_1 large enough such that for any x_1 with $|x_1 - x_0| < 2^{-(k_1+1)}$, we have

$$|f(1,1) - f(0,1)| = \left|u(\eta_1^1) - u(\eta_1^0)\right| = |u(x_1) - u(x_0)| < 2^{-1}$$

by continuity of $u(\cdot)$. Hence, (8) is satisfied.

Now, for any x_1 such that $|x_1 - x_0| < 2^{-(k_1+1)}$, ϕ_0 is a valid oracle for x_1 up to parameter value k_1. In particular, we can create an oracle ψ for x_1 which agrees with ϕ_0 on $1, 2, \ldots, k_1$. Then, the execution of $M_{l_1}^{\phi_0}(\hat{a}_1)$ will be identical to that of $M_{l_1}^{\psi}(\hat{a}_1)$, so it will output the same value A_{l_1} which is a $2^{-\hat{a}_1}$-approximation for $\Phi(x_0)$.

Applying Corollary 5.4 with $\varepsilon = 2^{-(k_1+1)}$, we get $L_1 > 0$ such that

$$\forall m_1 > L_1, \forall N_1 \in \mathbb{N}, |\beta^{N_1} - x_0| < 2^{-(k_1+1)},$$

where β^{N_1} has all ones except an N_1 at the (m_1+1)-th position. Applying Lemma 4.1 with $I_1 = [1]$ and $\varepsilon = 2^{-\hat{a}_1+1}$, and making sure the integer $m = m_1$ we get from this lemma satisfies $m_1 > L_1$, we get some $\beta^{N_1} = \beta_1^{N_1}$ for which

$$\exists N_1 \in \mathbb{N} \text{ such that } |\beta_1^{N_1} - x_0| < 2^{-(k_1+1)} \quad [\text{since } m_1 > L_1]$$

yet

$$\Phi(x_0) + 2^{-\hat{a}_1+1} < \Phi(\beta_1^{N_1}) < \Phi(x_0) + 2 \cdot 2^{-\hat{a}_1+1},$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

where $\beta_1^{N_1}$ has all ones except an N_1 at the (m_1+1)-th position. Let $x_1 = \beta_1^{N_1}$, and let $\phi_1 = \psi$ be the oracle for x_1 which agrees with ϕ_0 on $1, 2, \ldots, k_1$ and which additionally satisfies $|\phi_1(n) - x_1| < 2^{-(n+1)}$ for all n. This oracle ϕ_1 satisfies (1) and (2). As previously stated, the execution of $M_{l_1}^{\phi_0}(\hat{a}_1)$ is identical to that of $M_{l_1}^{\phi_1}(\hat{a}_1)$, so the output will be the same number A_{l_1} such that $|A_{l_1} - \Phi(x_0)| \leq 2^{-\hat{a}_1}$. By construction, $M_{l_1}^{\phi_1}(1), M_{l_1}^{\phi_1}(2), \ldots, M_{l_1}^{\phi_1}(\hat{a}_1)$ only query ϕ_1 with parameters not exceeding k_1, satisfying (3). But then, by the above work,

$$A_{l_1} + 2^{-\hat{a}_1} \leq \Phi(x_0) + 2^{-\hat{a}_1} + 2^{-\hat{a}_1} = \Phi(x_0) + 2^{-\hat{a}_1+1} < \Phi(x_1),$$

satisfying (4) and also (5) by taking $B_{l_1} = A_{l_1}$. Thus, $M_{l_1}^{\phi_1}(\hat{a}_1)$, where ϕ_1 is an oracle for x_1, does not approximate $\Phi(x_1)$ with precision $2^{-\hat{a}_1}$, so this TM does not properly compute this number. Additionally, note that $M_k^{\phi_1}$ also does not compute $\Phi(x_1)$ for $k = 1, 2, \ldots, l_1 - 1$. Running $M_k^{\phi_1}(a_k)$ is identical to running $M_k^{\phi_0}(a_k)$ by our choice of k_1 in the construction of ϕ_1, so $M_k^{\phi_1}(a_k)$ outputs a number B_k for which $|B_k - \Phi(x_0)| > 2^{-a_k} + \varepsilon_k$. Since

$$|\Phi(x_1) - \Phi(x_0)| < 2^{-\hat{a}_1+2} < \frac{\hat{\varepsilon}_1}{2} \leq \frac{\varepsilon_k}{2},$$

we have

$$\begin{aligned} |B_k - \Phi(x_1)| &\geq |B_k - \Phi(x_0)| - |\Phi(x_0) - \Phi(x_1)| > (2^{-a_k} + \varepsilon_k) - \frac{\varepsilon_k}{2} \\ &= 2^{-a_k} + \frac{\varepsilon_k}{2} > 2^{-a_k}, \end{aligned}$$

satisfying (6). To show (7), note that $\lim_{k\to\infty} f(1,k) = \Phi(x_1)$ is finite, so there is some large enough $n(1)$ for which $k \geq n(1)$ implies $|f(1,k) - \Phi(x_1)| < 2^{-1}$ as required.

5.3.2 Induction step

Now, inductively, suppose there exist the following:

- nested initial segments $I_1 \subseteq \cdots \subseteq I_{i-1}$, where each I_j has length p_j;
- for each $j = 1, 2, \ldots, i-1$, positive integers N^j and m_j;
- positive integers $k_1 < k_2 < \cdots < k_{i-1}$ and $l_1 < l_2 < \cdots < l_{i-1}$ and $\hat{a}_1 < \hat{a}_2 < \cdots < \hat{a}_{i-1}$ and $n(0) < n(1) < \cdots < n(i-1)$, positive real numbers $\hat{\varepsilon}_1 > \hat{\varepsilon}_2 > \cdots > \hat{\varepsilon}_{i-1}$, and oracles $\phi_0, \phi_1, \ldots, \phi_{i-1}$;

such that if we let $x_j = [I_j, 1, \ldots, 1, N^j, 1, \ldots]$ for $j = 1, 2, \ldots, i-1$, where N^j is in the $(m_j + p_j)$-th position, then we have the following:

(1) ϕ_j is an oracle for x_j such that $|\phi_j(n) - x_j| < 2^{-(n+1)}$ for all n.

(2) ϕ_{i-1} agrees with the oracle ϕ_{j-1} on inputs $1, 2, \ldots, k_j$.

(3) Running $M_{l_j}^{\phi_j}(1), M_{l_j}^{\phi_j}(2), \ldots, M_{l_j}^{\phi_j}(\hat{a}_j)$ queries ϕ_j with parameters not exceeding k_j.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

(4) Running $M_{l_j}^{\phi_j}(\hat{a}_j)$ yields a number A_{l_j} for which

$$A_{l_j} + 2^{-\hat{a}_j} \le \Phi(x_{j-1}) + 2^{-\hat{a}_j+1} < \Phi(x_j) < \Phi(x_{j-1}) + 2 \cdot 2^{-\hat{a}_j+1}.$$

(5) Running $M_{l_j}^{\phi_{i-1}}(\hat{a}_j)$ yields a number B_{l_j} for which $B_{l_j} + 2^{-\hat{a}_j} < \Phi(x_{i-1})$.

(6) The TMs $M_k^{\phi_{i-1}}$ for $k \in \{1, \dots, l_{i-1}\}$ all do not properly compute $\Phi(x_{i-1})$; in particular, they all compute $\Phi(x_{i-1})$ with an error of at least $\hat{\varepsilon}_{i-1}$.

(7) For $k \ge n(i-1)$, we have $|f(i-1,k) - \Phi(x_{i-1})| < 2^{-(i-1)}$.

(8) For $k = 1, 2, \dots, n(i-2)$, we have $|f(i-1,k) - f(i-2,k)| < 2^{-(i-1)}$.

Let $M_{l_i}^{\phi_{i-1}}$ be the first TM, with some $l_i > l_{i-1}$ and with ϕ_{i-1} being the oracle for x_{i-1} from the induction hypothesis, which computes $\Phi(x_{i-1})$. We want to find an initial segment $I_i \supseteq I_{i-1}$ of some length p_i, positive integers $N^i, m_i, k_i > k_{i-1}, \hat{a}_i > \hat{a}_{i-1}, n(i) > n(i-1)$, and an oracle ϕ_i for x_i such that (1)–(8) are satisfied for i instead of $i-1$. It would follow from (5) and (6) that none of $M_1^{\phi_i}, M_2^{\phi_i}, \dots, M_{l_i}^{\phi_i}$ properly compute $\Phi(x_i)$.

Since $M_{l_i}^{\phi_{i-1}}$ is the first such TM with $l_i > l_{i-1}$, combined with (6) from the induction hypothesis, we get that all of $M_1^{\phi_{i-1}}, M_2^{\phi_{i-1}}, \dots, M_{l_i-1}^{\phi_{i-1}}$ do not properly compute $\Phi(x_{i-1})$. So, there are integers $a_1, a_2, \dots, a_{l_i-1}$ and small positive real numbers $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{l_i-1} < \hat{\varepsilon}_{i-1}$ for which $M_k^{\phi_{i-1}}(a_k)$ outputs some number A_k with $|A_k - \Phi(x_{i-1})| > 2^{-a_k} + \varepsilon_k$. Set $\hat{\varepsilon}_i = \min(\varepsilon_1, \dots, \varepsilon_{l_i-1})$, choose $\hat{n}_i$ large enough so that $2^{-\hat{n}_i+2} < \hat{\varepsilon}_i/2^i$, and set $\hat{a}_i = \max(a_1, \dots, a_{l_i-1}, \hat{a}_{i-1}, \hat{n}_i)$.

When run, the TM $M_{l_i}^{\phi_{i-1}}(\hat{a}_i)$ outputs a number A_{l_i} for which $|A_{l_i} - \Phi(x_{i-1})| \le 2^{-\hat{a}_i}$. The computations $M_{l_j}^{\phi_j}(1), M_{l_j}^{\phi_j}(2), \dots, M_{l_j}^{\phi_j}(\hat{a}_i)$ are performed in finite time, so there is a $k_i > 0$ such that ϕ_{i-1} is only queried with parameters not exceeding k_i for all of these computations; we can make k_i arbitrarily large, so assume $k_i > k_{i-1}$. We additionally make k_i large enough such that for any x_i with $|x_i - x_{i-1}| < 2^{-(k_i+1)}$ and any $k = 1, 2, \dots, n(i-1)$, we have

$$|f(i,k) - f(i-1,k)|$$
$$= \left| \sum_{n=1}^{k} s(i) \cdot \left(\eta_0^i \eta_1^i \dots \eta_{n-1}^i\right)^\nu \cdot u(\eta_n^i) - \sum_{n=1}^{k} s(i) \cdot \left(\eta_0^{i-1} \eta_1^{i-1} \dots \eta_{n-1}^{i-1}\right)^\nu \cdot u(\eta_n^{i-1}) \right| < 2^{-i}.$$

We can do this because the sums are finite and have continuous dependence on x_i, since u and G are $\mathcal{C}^1$ on the set S (from the definition of G). Hence, (8) is satisfied.

Now, for any x_i such that $|x_i - x_{i-1}| < 2^{-(k_i+1)}$, ϕ_{i-1} is a valid oracle for x_i up to parameter value k_i. In particular, we can create an oracle ψ for x_i which agrees with ϕ_{i-1} on $1, 2, \dots, k_i$ and so that $|\psi(n) - x_i| < 2^{-(n+1)}$ for all n. Then, the execution of $M_{l_i}^{\phi_{i-1}}(\hat{a}_i)$ will be identical to that of $M_{l_i}^{\psi}(\hat{a}_i)$, so it will output the same number A_{l_i} which is a $2^{-\hat{a}_i}$-approximation for $\Phi(x_{i-1})$.

Applying Corollary 5.4 with $\varepsilon = 2^{-(k_i+1)}$, we get $L_i > 0$ such that

$$\forall m_i > L_i, \forall N_i \in \mathbb{N}, |\beta^{N_i} - x_{i-1}| < 2^{-(k_i+1)},$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

where $\beta^{N_i} = [I_{i-1}, 1, \ldots, 1, N_{i-1}, 1, \ldots]$ agrees with x_{i-1} at all entries except having an N_{i-1} instead of a 1 at the $(m_{i-1} + p_{i-1})$-th position. Applying Lemma 4.1 with $I_i = [I_{i-1}, 1, \ldots, 1, N_{i-1}]$ (where N_{i-1} is at the $(m_{i-1} + p_{i-1})$-th position) and $\varepsilon = 2^{-\hat{a}_i+1}$, and making sure the integer $m = m_i$ we get from this lemma satisfies $m_i > L_i$, we get some $\beta^{N_i} = \beta_i^{N_i}$ for which

$$\exists N_i \in \mathbb{N} \text{ such that } |\beta_i^{N_i} - x_{i-1}| < 2^{-(k_i+1)} \quad [\text{since } m_i > L_i]$$

yet

$$\Phi(x_{i-1}) + 2^{-\hat{a}_i+1} < \Phi(\beta_i^{N_i}) < \Phi(x_{i-1}) + 2 \cdot 2^{-\hat{a}_i+1}.$$

Let $x_i = \beta_i^{N_i}$, and let $\phi_i = \psi$ be the oracle for x_i which agrees with ϕ_{i-1} on $1, 2, \ldots, k_i$. This ϕ_i satisfies (1) and (2). As previously stated, the execution of $M_{l_i}^{\phi_{i-1}}(\hat{a}_i)$ is identical to that of $M_{l_i}^{\phi_i}(\hat{a}_i)$, so the output will be the same number A_{l_i} such that $|A_{l_i} - \Phi(x_{i-1})| \le 2^{-\hat{a}_i}$. By construction, $M_{l_i}^{\phi_i}(1), M_{l_i}^{\phi_i}(2), \ldots, M_{l_i}^{\phi_i}(\hat{a}_i)$ only query ϕ_i with parameters not exceeding k_i, satisfying (3). But then, by the above work, we have

$$A_{l_i} + 2^{-\hat{a}_i} \le \Phi(x_{i-1}) + 2^{-\hat{a}_i} + 2^{-\hat{a}_i} = \Phi(x_{i-1}) + 2^{\hat{a}_i+1} < \Phi(x_i) < \Phi(x_{i-1}) + 2 \cdot 2^{-\hat{a}_i+1},$$

satisfying (4). Thus, $M_{l_i}^{\phi_i}(\hat{a}_i)$, where ϕ_i is an oracle for x_i, does not compute $\Phi(x_i)$ with precision $2^{-\hat{a}_i}$, so it does not properly compute this number. But now since ϕ_i agrees with ϕ_{i-1} on $1, 2, \ldots, k_i$, by (2) of the induction hypothesis, we have that it agrees with ϕ_{j-1} on $1, 2, \ldots, k_j$ for $j = 1, 2, \ldots, i$. By (3) of the induction hypothesis, running $M_{l_j}^{\phi_j}(\hat{a}_j)$ queries ϕ_j with parameters not exceeding $k_j \le k_i$, hence, the execution of $M_{l_j}^{\phi_j}(\hat{a}_j)$ is identical to that of $M_{l_j}^{\phi_i}(\hat{a}_j)$ for all j. Therefore, by (4), running $M_{l_j}^{\phi_i}(\hat{a}_j)$ gives a number $B_{l_j} = A_{l_j}$ such that

$$B_{l_j} + 2^{-\hat{a}_j} \le \Phi(x_{j-1}) + 2^{\hat{a}_j+1} < \Phi(x_j) < \Phi(x_{j+1}) < \cdots < \Phi(x_i).$$

Hence, (5) is satisfied.

Now, note that $M_k^{\phi_i}$ also does not compute $\Phi(x_i)$ properly for $k = 1, 2, \ldots, l_i - 1$. Running $M_k^{\phi_i}(a_k)$ is identical to running $M_k^{\phi_{i-1}}(a_k)$ by our choice of k_i in the construction of ϕ_i, so $M_k^{\phi_i}(a_k)$ outputs a number B_k for which $|B_k - \Phi(x_{i-1})| > 2^{-a_k} + \varepsilon_k$. Since

$$|\Phi(x_i) - \Phi(x_{i-1})| < 2^{-\hat{a}_i+2} < \frac{\hat{\varepsilon}_i}{2^i} \le \frac{\varepsilon_k}{2},$$

we have

$$\begin{aligned} |A_k - \Phi(x_i)| &\ge |A_k - \Phi(x_{i-1})| - |\Phi(x_{i-1}) - \Phi(x_i)| > (2^{-a_k} + \varepsilon_k) - \frac{\varepsilon_k}{2} \\ &= 2^{-a_k} + \frac{\varepsilon_k}{2} > 2^{-a_k}, \end{aligned}$$

satisfying (6). Finally, it remains to show (7) for i. Note that $\lim_{k\to\infty} f(i,k) = \Phi(x_i)$ is finite, so there is some $n(i) > n(i-1)$ for which $k \ge n(i)$ implies $|f(i,k) - \Phi(x_i)| < 2^{-i}$ as required. This completes the induction.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

5.3.3 Finalizing the argument

Let $[a_1, a_2, \dots] = x_\infty = \lim_{i\to\infty} x_i$. We claim that $\Phi(x_\infty) < \infty$ and $\Phi(x_\infty)$ is not computable by any TM.

We first show that $\lim_{n\to\infty} \Phi(x_i) < \Phi(x_0) + 4 < \infty$. From (4), we have for all $i \in \mathbb{N}$ that

$$\Phi(x_i) < \Phi(x_{i-1}) + 2 \cdot 2^{-i+1},$$

hence,

$$\Phi(x_i) < \Phi(x_0) + \sum_{j=1}^{i} 2^2 \cdot 2^{-j} \implies \sup_{i\in\mathbb{N}} \Phi(x_i) \le \Phi(x_0) + 4.$$

We now show that $\Phi(x_\infty)$ is finite and equals $\lim_{i\to\infty} \Phi(x_i)$. For each i, the sequence $(f(j, n(i)))_{j=1}^{\infty}$ is Cauchy by (8), hence, it converges to some $f_\infty(n(i))$. It is clear that

$$f_\infty(n(i)) = \sum_{n=1}^{n(i)} s(i) \cdot (\eta_0^\infty \eta_1^\infty \dots \eta_{n-1}^\infty)^\nu \cdot u(\eta_n^\infty)$$

by definition of x_∞ and by the way, each successive x_j was chosen. Since each $f(j, n(i)) \le \Phi(x_j)$, taking limits on both sides gives $f_\infty(n(i)) \le \Phi(x_0) + 4$. Thus, the sequence $(f_\infty(n(i)))_{i=1}^{\infty}$ is bounded from above, so the limit superior is finite. But now

$$\limsup_{i\to\infty} f_\infty(n(i)) = \limsup_{i\to\infty} \sum_{n=1}^{n(i)} s(i) \cdot (\eta_0^\infty \eta_1^\infty \dots \eta_{n-1}^\infty)^\nu \cdot u(\eta_n^\infty) \ge \Phi(x_\infty),$$

showing that $\Phi(x_\infty)$ is finite. Now, we claim that $\Phi(x_\infty) = \lim_{i\to\infty} \Phi(x_i)$. Let $\varepsilon > 0$ be given.

- Choose i_1 large enough so that
$$i > i_1 \implies |f_\infty(n(i)) - \Phi(x_\infty)| < \varepsilon/3.$$
- Choose $i_2 > i_1$ large enough so that $2^{-i_2} < \varepsilon/3$. By (7), we have
$$i > i_2 \implies |f(i, n(i)) - \Phi(x_i)| < \varepsilon/3.$$
- Set $i_3 = i_2 + 1$, so that $2^{-i_3+1} < \varepsilon/3$. Then, for $i > i_3$, repeated application of (8) yields
$$|f(i, n(i)) - f_\infty(n(i))| \le \sum_{j=i}^{\infty} |f(j+1, n(i)) - f(j, n(i))| < \sum_{j=i}^{\infty} 2^{-j} = 2^{-i+1} < \varepsilon/3.$$

Taking $i > \max\{i_1, i_2, i_3\}$, we finally get

$$\begin{aligned} &|\Phi(x_\infty) - \Phi(x_i)| \\ &\quad \le |\Phi(x_\infty) - f_\infty(k)| + |f_\infty(n(i)) - f(i, n(i))| + |f(i, n(i)) - \Phi(x_i)| < \varepsilon. \end{aligned}$$

It remains to show that $\Phi(x_\infty)$ is not computable by any of the TMs M_i^ϕ, where ϕ is an oracle for x_∞. Let ϕ_∞ be the oracle for x_∞ which agrees with ϕ_i on inputs $1, 2, \dots, k_{i+1}$. This is a valid construction of an oracle by (2) and since $\lim_{i\to\infty} k_{i+1} = \infty$. To show that

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

ϕ_∞ is indeed an oracle for x_∞, let $n \in \mathbb{Z}^+$ and set i such that $n+1 < k_i$. Since $n+1 < k_i$ implies that $n+j+1 < k_{i+j}$,

$$|\phi_\infty(n) - x_\infty| \le |\phi_\infty(n) - x_i| + |x_i - x_\infty| \le |\phi_i(n) - x_i| + \sum_{j=i}^{\infty} |x_{j+1} - x_j|$$
$$< 2^{-(n+1)} + \sum_{j=0}^{\infty} 2^{-(k_{i+j}+1)} \le 2^{-(n+1)} + \sum_{j=0}^{\infty} 2^{-(n+j+2)} = 2^{-n},$$

so ϕ_∞ is indeed an oracle for x_∞. Now, for a contradiction, suppose some TM $M_j^{\phi_\infty}$ computes $\Phi(x_\infty)$. We have two cases:

(1) $j = l_i$ for some i. Then, $M_{l_i}^{\phi_i}(\hat{a}_i)$ outputs B_i for which $B_i + 2^{-\hat{a}_i} < \Phi(x_i)$ by (5). This same number B_i is output when running $M_{l_i}^{\phi_\infty}(\hat{a}_i)$. But then

$$B_i + 2^{-\hat{a}_i} < \Phi(x_i) < \Phi(x_{i+1}) < \cdots < \Phi(x_\infty),$$

so we cannot have $|B_i - \Phi(x_\infty)| \le 2^{-\hat{a}_i}$, contradicting the assumption that $M_{l_i}^{\phi_\infty}$ computes $\Phi(x_\infty)$.

(2) $j \ne l_i$ for all i. Choose the smallest $i > 2$ for which $j < l_i$. Then, $M_j^{\phi_{i-1}}(a_j)$ outputs A_j for which $|A_j - \Phi(x_{i-1})| > 2^{-a_j} + \varepsilon_j$, where a_j and ε_j are from the i-th step of the induction. This same number A_j is output when running $M_j^{\phi_\infty}(a_j)$. By assumption, $|A_j - \Phi(x_\infty)| < 2^{-a_j}$, hence $|\Phi(x_{i-1}) - \Phi(x_\infty)| > \varepsilon_j$. But

$$|\Phi(x_k) - \Phi(x_{k+1})| < 2^{-\hat{a}_k+2} \le 2^{-\hat{n}_k+2} < \frac{\varepsilon_j}{2^k}$$

for all $k \ge i-1$, thus

$$|\Phi(x_{i-1}) - \Phi(x_\infty)| \le \sum_{k=i-1}^{\infty} |\Phi(x_k) - \Phi(x_{k+1})| < \sum_{k=i-1}^{\infty} \frac{\varepsilon_j}{2^k} \le \varepsilon_j,$$

a contradiction.

We have shown that none of the TMs $M_j^{\phi_\infty}$ compute $\Phi(x_\infty)$, where ϕ_∞ is an oracle for x_∞. This completes the proof of Theorem 5.2.

A Proofs of the main technical statements

A.1 Proof of Theorem 3.1

The most involved part of the proof will be showing that conditions (i)–(vii) on G are satisfied by the α-continued fraction expansion maps A_α for $\alpha \in [1/2, 1]$.

Lemma A.1 *Conditions (i)–(vii) are satisfied by both the Gauss map and by A_α for $\alpha \in [1/2, 1)$.*

Proof Properties (i), (iii), and (v)–(vii) are easy enough that we group them together and prove them at once, and then (ii) and (iv) are verified separately.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

(i), (iii), and (v)–(vii). For the Gauss map G, $G(J_i) = G((\frac{1}{i+1}, \frac{1}{i})) = (0,1) = (s_0, s_1)$. It is clear that G is decreasing on $J_1 = (1/2, 1)$. Since $J_i = (\ell_i, r_i) = (\frac{1}{i+1}, \frac{1}{i})$, we have

$$\frac{r_i - \ell_i}{\ell_i^2} < 2 =: D \text{ and } g(i) = \frac{r_i}{\ell_{i+1}} \xrightarrow{i\to\infty} 0.$$

Finally, since G is a composition of computable functions, it is computable.

Now, let $\alpha \in [1/2, 1)$, and consider the map A_α corresponding to α-continued fraction expansions. Set $n_1 := \lceil 1/(1-\alpha) \rceil$; this is the smallest integer for which $1/n_1 \le 1 - \alpha$ and $A_\alpha(1/n_1) = 0$. We restrict the domain and codomain of $A_\alpha(x)$ to the maximal invariant set $\Lambda \subseteq (s_0, s_1) := (0, 1/n_1) \subseteq (0, 1-\alpha)$, and henceforth consider $A_\alpha : \Lambda \to \Lambda$. It is clear that Λ is a countable disjoint union of connected intervals, which we label $J_1 = (\ell_1, r_1), J_2 = (\ell_2, r_2), \ldots$ with $r_1 > \ell_1 \ge r_2 > \ell_2 \ge \cdots$. From maximality of the invariant set and the fact that $|A'_\alpha(x)| > 1$ on Λ, (i) holds. For (vi), computability of A_α again follows since it is a composition of computable functions. We will make use of the following facts about A_α and the intervals J_j:

- A_α is decreasing on J_j for odd j and increasing on J_j for even j. In particular, since $A_\alpha(r_1) = A_\alpha(1/n_1) = 0$, $|A'_\alpha(x)| > 1$, and $A_\alpha(x) > 0$, A_α is decreasing on J_1 and so (iii) is satisfied.
- For $\alpha = 1/2$, it can be readily computed by solving the equations $A_\alpha(x) = 0$ and $A_\alpha(x) = 1$ that $n_1 = 2$ and $J_j = (\ell_j, r_j) = (\frac{2}{j+4}, \frac{2}{j+3})$.
- For $\alpha \in (1/2, 1)$, we have

$$\begin{aligned}
\ell_j &> \frac{2}{2(n_1 - 2 + k) + 3} \\
&= \frac{2}{j + 2n_1}, r_j = \frac{1}{n_1 - 1 + k} = \frac{2}{j + 2n_1 - 1} \text{ for } j = 2k-1 \text{ odd, and} \\
\ell_j &= \frac{1}{n_1 + k} = \frac{2}{2n_1 + j}, r_j < \frac{2}{2(n_1 - 2 + k) + 3} = \frac{2}{j + 2n_1 - 1} \text{ for } j = 2k \text{ even.}
\end{aligned}$$

It is easy to verify (v) and (vii) by showing that $\frac{r_i - \ell_i}{\ell_i^2}$ is bounded from above independently of i and that $g(i) \xrightarrow{i\to\infty} 0$.

(ii). For G, we note that $G'(x) = 1/x^2$, $G''(x) = 2/x^3$, where it is defined. Since G is decreasing on all J_j, we have $\tau_{i,1} = |G'(r_i)| = i^2$. Taking any $\sigma > 2$, $\tau_{i,1}^{-1} < \ell_i \sigma$ as needed. Now, set $\kappa = 2$. Then, where it is defined, it is easy to check that $(G^2)'' > 0$ and thus $\tau_{i,2} = |(G^2)'(\ell_i)| = (i+1)^2$. Hence, for any $1 < \tau < 2$, we have $\tau_{i,2}^{-1} < \ell_i \cdot \tau^{-1}$ as needed.

Now, again consider A_α, for $\alpha \in [1/2, 1)$. As for the Gauss map, $A'_\alpha(x) = 1/x^2$, $A''_\alpha(x) = 2/x^3 > 0$, where it is defined. Set $\kappa = 1$. Then, $\tau_{i,\kappa} = \tau_{i,1} = 1/r_i^2$.

- Let $\sigma > 1$ be large enough that for all i, $\sigma > \frac{2(2n_1 + i)}{(i + 2n_1 - 1)^2}$. It is easily verified that $\tau_{i,1}^{-1} < \ell_i \sigma$.
- Now, note that

$$\tau_{i,1}^{-1} = r_i^2 < \ell_i \tau^{-1} \iff \frac{2(2n_1 + i)}{(i + 2n_1 - 1)^2} < \tau^{-1}.$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Since the left-hand side tends to zero as $n_1 \to \infty$ for all $i \geq 1$, restricting Λ to a small enough invariant set under A_α, that is, making $r_1 = \frac{1}{n_1}$ small enough, gives existence of some $\tau > 1$ for which the above holds.

(iv). For the Gauss map, recalling that $\varphi = \frac{\sqrt{5}-1}{2} = [1, 1, \ldots]$, we have

$$\delta_G(N) = \frac{1}{(N+\varphi)(N+1+\varphi)} \text{ and so } \frac{r_{N+1}}{\ell_{N+1}} \cdot \frac{r_n - \ell_{N+1}}{\delta_G(N)} < D$$

for some constant D. We first show that (iv) holds for $A_{1/2}$, then for A_α, $\alpha \in (1/2, 1)$. We denote the symbolic expansion of a point in Λ generated by $A_{1/2}$ as $[a_1, a_2, a_3, \ldots]_{1/2}$, and that of a point in the invariant set generated by G as $[a_1, a_2, a_3, \ldots]_1$. Since the interval J_1 corresponding to $A_{1/2}$ is contained in the interval J_2 corresponding to G, we have $\psi := [1, 1, 1, \ldots]_{1/2} = [2, 2, 2, \ldots]_1 = \sqrt{2} - 1$. For $N = 2k - 1$ odd, the interval J_N corresponding to $A_{1/2}$ is contained in the interval $J_{(N+3)/2}$ corresponding to G, so

$$[N, 1, 1, \ldots]_{1/2} = [k+1, 2, 2, \ldots]_1 = [(N+3)/2, 2, 2, \ldots]_1 = \frac{1}{\frac{N+3}{2} + \psi} = \frac{2}{N+3+2\psi}.$$

Now, consider $N = 2k$ even. Observe that on J_N, $A_{1/2}(x) = 1 - G(x)$. We have

$$[N, 1, 1, \ldots]_{1/2} = (A_{1/2}|_{J_N})^{-1}([1, 1, \ldots]_{1/2}) = G^{-1}_{N/2+1}(1 - [2, 2, \ldots]_1) = \frac{2}{N+4-2\psi}.$$

Thus,

$$\begin{aligned} \delta_{A_{1/2}}(N) &= [N, 1, 1, \ldots]_{1/2} - [N+1, 1, 1, \ldots]_{1/2} \\ &= \frac{4 - 8\psi}{(N+3+2\psi)(N+5-2\psi)} \text{ if } N \text{ is odd, and} \\ \delta_{A_{1/2}}(N) &= \frac{8\psi}{(N+4-2\psi)(N+4+2\psi)} \text{ if } N \text{ is even.} \end{aligned}$$

Noting that $r_N - \ell_{N+1} = \frac{4}{(N+3)(N+5)}$ and $\frac{r_{N+1}}{\ell_{N+1}} = \frac{N+5}{N+4}$, it is clear that some constant D upper-bounds $\frac{r_{N+1}}{\ell_{N+1}} \cdot \frac{r_N - \ell_{N+1}}{\delta_{A_{1/2}}(N)}$. Finally, for $\alpha \in (1/2, 1)$, we have $A_{1/2} = A_\alpha$ on the invariant set Λ corresponding to A_α. Thus, the asymptotic behavior of ℓ_N, r_N, and $\delta_{A_\alpha}(N)$ is identical, so (iv) holds in this case as well. ∎

We can now prove Theorem 3.1 without too much difficulty.

Proof of Theorem 3.1. By Lemma A.1 and since the function u in the definition of $\mathcal{B}_{\alpha,u,v}$ satisfies $\lim_{x \to 0^+} u(x) = \infty$ by assumption, conditions (i)–(vii) on G and (i)–(v) on u hold for $\mathcal{B}_{\alpha,u,v}$. To prove this theorem, it remains to show that for $G = A_\alpha$, $\alpha \in [1/2, 1]$, conditions (i)–(v) hold for both $u(x) = \log^n(1/x)$, $n \in \mathbb{Z}^+$ and $u(x) = x^{-1}$. It is clear that both these functions are computable on $\mathbb{R}^+$ and tend to infinity near zero. If $\alpha \in [1/2, 1)$, then $s_1 < 1$, so (ii) in this case is satisfied. Thus, we only need to show (ii) in the case that G is the ordinary Gauss map, and (iii).

For $u(x) = x^{-1}$, (iii) is obvious. If $u(x) = \log^n(1/x)$, then $|u'(x)| = \frac{n}{x} \cdot \log^{n-1}(1/x)$. Since there is some $C_n > 0$ for which $\log^{n-1}(y) < C_n y$ for all large enough y, we have $\log^{n-1}(1/x) < \frac{C_n}{x}$ for all small enough x. Thus, $|u'(x)| < \frac{nC_n}{x^2}$ for all small enough x and

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

since $|u'(x)|$ is bounded for x which is bounded away from zero, there is some $C > 0$ for which $|u'(x)| < \dfrac{C}{(x - s_0)^2}$ for all $x \in (s_0, s_1)$ as needed.

Now, suppose G is the Gauss map. Then, $u \circ G_1^{-1} \circ G_N^{-1}(z)$ is decreasing with respect to N since $u(\cdot)$ is decreasing, so

$$\frac{u \circ G_1^{-1} \circ G_N^{-1}(z)}{u \circ G_1^{-1} \circ G_N^{-1}(w)} \geq \frac{u \circ G_1^{-1} \circ G_N^{-1}(1)}{u \circ G_1^{-1} \circ G_N^{-1}(0)} =: v(N).$$

A tedious computation shows that $v'(N) > 0$ and $v(N) > 0$ for all N and both $u(x) = \log^n(1/x)$ and $u(x) = x^{-1}$, therefore (ii) holds.

A.2 Proofs of the main lemmas

Here, we will prove the three main lemmas used in the above work. Lemma 4.1 will be proven under the weaker assumptions on G and u along with the $s(i)$ term in Section 5, and Lemmas 4.2 and 4.3 will be proven under the full assumptions in Section 3. We recall these lemmas below.

Lemma 4.1 *For any initial segment $I = [a_1, a_2, \ldots, a_n]$, write $\omega = [a_1, a_2, \ldots, a_n, 1, 1, 1, \ldots]$. Then, for any $\varepsilon > 0$, there is an $m > 0$ and an integer N such that if we write $\beta = [a_1, a_2, \ldots, a_n, 1, 1, \ldots, 1, N, 1, 1, \ldots]$, where the N is located in the $(n + m)$-th position, then*

$$\Phi(\omega) + \varepsilon < \Phi(\beta) < \Phi(\omega) + 2\varepsilon.$$

Lemma 4.2 *Write $\omega = [a_1, a_2, \ldots, a_n, 1, 1, 1, \ldots]$. Then, for any $\varepsilon > 0$, there is an $m_0 > 0$, which can be computed from $(a_1, a_2, \ldots, a_n)$ and ε, such that for any $m \geq m_0$ and for any tail $I = [a_{n+m}, a_{n+m+1}, \ldots]$,*

$$\Phi(\beta^I) > \Phi(\omega) - \varepsilon,$$

where

$$\beta^I = [a_1, a_2, \ldots, a_n, 1, 1, \ldots, 1, a_{n+m}, a_{n+m+1}, \ldots].$$

Lemma 4.3 *Let $\omega = [a_1, a_2, \ldots]$ be such that $\Phi(\omega) < \infty$. Write $\omega_k = [a_1, a_2, \ldots, a_k, 1, 1, \ldots]$. Then, for every $\varepsilon > 0$, there is an m such that, for all $k \geq m$,*

$$\Phi(\omega_k) < \Phi(\omega) + \varepsilon.$$

Write

$$\beta^N = [a_1, a_2, \ldots, a_n, 1, 1, \ldots, 1, N, 1, 1, \ldots],$$

where N is in the $(m + n)$-th position. The following preliminary lemmas are required for all of the main proofs below.

Lemma A.2 *For any $s_0 < a < s_1$, the function u is bounded on the interval $[a, s_1)$.*

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Proof By assumption on u, there is some $C > 0$ for which $-u'(x) < \frac{C}{(x-s_0)^2}$ for all $x \in [a, s_1)$. Integrating both sides gives $u(x) < \frac{\tilde{C}}{x-s_0} \leq \frac{\tilde{C}}{a-s_0}$ for another constant $\tilde{C} > 0$ and all $x \in [a, s_1)$. ∎

Lemma A.3 *Let β^N, β^1 be as before, let β^I, ω be as in Lemma 4.2, and let γ_1, γ_2 be two numbers whose symbolic representations coincide in the first $n+m-1$ terms $[a_1, a_2, \ldots, a_{n+m-1}]$ (in particular, we could have $\gamma_1 = \beta^N$ and $\gamma_2 = \beta^1$). Then, for g and m_g as in (vii), for κ as in (ii), and for any N, the following holds:*

(1) *For $i \leq n+m$,*

$$\left|\log \frac{\eta_i(\beta^N)}{\eta_i(\beta^{N+1})}\right| < \frac{g(N)}{\ell_1} \cdot \tau^{1+\frac{i-(n+m)}{\kappa}}.$$

(2) *For $i < n+m$,*

$$\left|\log \frac{\eta_i(\gamma_1)}{\eta_i(\gamma_2)}\right| < m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.$$

(3) *For all large enough m, there is some positive function f with $\lim_{m\to\infty} f(m) = 0$ such that for $i < n+m-1$,*

$$\left|\log \frac{u(\eta_i(\beta^I))}{u(\eta_i(\omega))}\right| < f(m).$$

In particular, we could take $\beta^I = \beta^1$ and $\omega = \beta^N$. Clearly, $0 < \sup_{m\in\mathbb{N}} f(m) =: M < \infty$.

Proof of (1) We will first show that for all i, we have

$$(*_1) \qquad |\eta_i(\beta^N) - \eta_i(\beta^{N+1})| < g(N) \cdot \tau^{1+\frac{i-(n+m)}{\kappa}}.$$

If $i = n+m$, since $[N, 1, 1, \ldots] \in J_N$, $[N+1, 1, 1, \ldots] \in J_{N+1}$, for all $N \in \mathbb{N}$, we have

$$1 < \frac{\eta_{n+m}(\beta^N)}{\eta_{n+m}(\beta^{N+1})} = \frac{[N, 1, 1, \ldots]}{[N+1, 1, 1, \ldots]} < \frac{r_N}{\ell_{N+1}} < 1 + g(N)$$

and so

$$|\eta_{n+m}(\beta^N) - \eta_{n+m}(\beta^{N+1})| < g(N) < g(N) \cdot \tau^{1+\frac{i-(n+m)}{\kappa}}.$$

Now, let $1 \leq j < \kappa$. We wish to show $*_1$ for $i = n+m-j$. Note that since $|G_1'(x)| > 1$ for all $x \in S$ by (ii) and G_1 is $\mathcal{C}^1$, the Inverse Function Theorem gives $|(G_1^{-1})'(x)| < 1$ for all $x \in J_1$. Repeatedly applying the Mean Value Theorem gives

$$\begin{aligned} |\eta_{n+m-j}(\beta^N) - \eta_{n+m-j}(\beta^{N+1})| &< |\eta_{n+m}(\beta^N) - \eta_{n+m}(\beta^{N+1})| \\ &< g(N) < g(N) \cdot \tau^{1-j/\kappa}. \end{aligned}$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

For $i = n + m - \kappa$, since $|(G_1^\kappa)'| > \tau$ by assumption (ii) on G, we similarly use Inverse Function Theorem and Mean Value Theorem (applied to the function $G_1^{-\kappa}$ this time) to get

$$\begin{aligned}|\eta_{n+m-\kappa}(\beta^N) - \eta_{n+m-\kappa}(\beta^{N+1})| &< \tau^{-1} \cdot |\eta_{n+m}(\beta^N) - \eta_{n+m}(\beta^{N+1})| \\ &< g(N) \cdot \tau^{1+\frac{i-(n+m)}{\kappa}}.\end{aligned}$$

To show $*_1$ for all other $i > n + m - j$ with $j > \kappa$, we apply the inequality

$$|\eta_{n+m-j}(\beta^N) - \eta_{n+m-j}(\beta^{N+1})| < \tau^{-1} \cdot |\eta_{n+m-j+\kappa}(\beta^N) - \eta_{n+m-j+\kappa}(\beta^{N+1})|$$

repeatedly until one of the cases $i \geq n + m - \kappa$ above is reached.

We now proceed to prove (1). If $i = n + m$, then as before for all $N \in \mathbb{Z}^+$, we have

$$1 < \frac{\eta_i(\beta^N)}{\eta_i(\beta^{N+1})} < 1 + g(N)$$

and so

$$\left|\log \frac{\eta_i(\beta^N)}{\eta_i(\beta^{N+1})}\right| < \log(1 + g(N)) < \log(e^{g(N)}) = g(N) < \frac{g(N)}{\ell_1} \cdot \tau^{1+\frac{i-(n+m)}{\kappa}},$$

where we used the inequality $1 + \xi < e^\xi$ for any $\xi > 0$.

Let $1 \leq i < n + m$, and assume $\eta_i(\beta^N) > \eta_i(\beta^{N+1})$; the complementary case is almost identical. We have $\eta_i(\beta^N), \eta_i(\beta^{N+1}) > \ell_1$ since they are in J_1 and so from $*_1$,

$$\begin{aligned}&\left|\frac{\eta_i(\beta^N) - \eta_i(\beta^{N+1})}{\eta_i(\beta^{N+1})}\right| \\ &< \frac{g(N)}{\ell_1} \cdot \tau^{1+\frac{i-(n+m)}{\kappa}} \text{ and so } 1 < \frac{\eta_i(\beta^N)}{\eta_i(\beta^{N+1})} < 1 + \frac{g(N)}{\ell_1} \cdot \tau^{1+\frac{i-(n+m)}{\kappa}}\end{aligned}$$

As above, this gives the desired inequality. ∎

Proof of (2) Note that $\eta_i(\gamma_1)$ and $\eta_i(\gamma_2)$ agree on the first digit of the symbolic expansion for each $1 \leq i < n + m$; we write this digit as N_i.

We will first show that for all i, we have

$$(*_2) \qquad \frac{\eta_i(\gamma_1)}{\eta_i(\gamma_2)} < 1 + m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.$$

If $\eta_i(\gamma_1) < \eta_i(\gamma_2)$, this is obviously true as the second term on the right-hand side is greater than zero, but to unify the argument, we will not split into cases.

For $i = n + m - 1$, using assumption (vii) on G, we compute

$$\frac{\eta_{n+m-1}(\gamma_1)}{\eta_{n+m-1}(\gamma_2)} < \frac{r_{N_{n+m-1}}}{\ell_{N_{n+m-1}}} < 1 + g(N_{n+m-1}) \leq 1 + m_g < 1 + m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.$$

For future use, note that the above inequalities yield

$$\eta_{n+m-1}(\gamma_1) - \eta_{n+m-1}(\gamma_2) \leq \eta_{n+m-1}(\gamma_2) \cdot m_g < m_g.$$

Now, let $1 \leq j < \kappa$. We want to show $*_2$ for $i = n + m - 1 - j$. Similar to the proof of (1), we note that $|(G^{-1}_{N_{n+m-1-j}})'(x)| < \tau^{-1}_{N_{n+m-1-j},1}$ for all $x \in J_{N_{n+m-1-j}}$ and (ii) gives

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

$|(G_{N_t}^{-1})'(x)| < 1$ for all $x \in J_{N_t}$, t arbitrary, so repeatedly applying the Mean Value Theorem,

$$\begin{aligned}
&\eta_{n+m-1-j}(\gamma_1) - \eta_{n+m-1-j}(\gamma_2) \\
&< |G_{N_{n+m-1-j}}^{-1} \circ \cdots \circ G_{N_{n+m-2}}^{-1}(\eta_{n+m-1}(\gamma_1)) - G_{N_{n+m-1-j}}^{-1} \circ \cdots \circ G_{N_{n+m-2}}^{-1}(\eta_{n+m-1}(\gamma_2))| \\
&< \tau_{N_{n+m-1-j},1}^{-1} \cdot |G_{N_{n+m-j}}^{-1} \circ \cdots \circ G_{N_{n+m-2}}^{-1}(\eta_{n+m-1}(\gamma_1)) \\
&\quad - G_{N_{n+m-j}}^{-1} \circ \cdots \circ G_{N_{n+m-2}}^{-1}(\eta_{n+m-1}(\gamma_2))| \\
&< \tau_{N_{n+m-1-j},1}^{-1} \cdot |\eta_{n+m-1}(\gamma_1) - \eta_{n+m-1}(\gamma_2)| < \tau_{N_{n+m-1-j},1}^{-1} \cdot m_g \\
&< \ell_{N_{n+m-1-j}} \cdot \sigma \cdot m_g < \eta_{n+m-1-j}(\gamma_2) \cdot \sigma \cdot m_g \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.
\end{aligned}$$

Rearranging gives $*_2$. To show $*_2$ holds for $i - l \cdot \kappa$ with $n + m - 1 - \kappa < i \le n + m - 1$ and $l \in \mathbb{Z}^+$, repeatedly apply the inequality

$$\begin{aligned}
&\eta_{i-j\cdot\kappa}(\gamma_1) - \eta_{i-l\cdot\kappa}(\gamma_2) \\
&< |G_{N_{i-l\cdot\kappa}}^{-1} \circ \cdots \circ G_{N_{i-l\cdot\kappa+\kappa-1}}^{-1}(\eta_{i-(l-1)\cdot\kappa}(\gamma_1)) - G_{N_{i-l\cdot\kappa}}^{-1} \circ \cdots \\
&\quad \circ G_{N_{i-l\cdot\kappa+\kappa-1}}^{-1}(\eta_{i-(l-1)\cdot\kappa}(\gamma_2))| \\
&= \left|\frac{1}{(G^\kappa)'(\xi)}\right| \cdot |\eta_{i-(l-1)\cdot\kappa}(\gamma_1) - \eta_{i-(l-1)\cdot\kappa}(\gamma_2)| \\
&< \tau^{-1} \cdot |\eta_{i-(l-1)\cdot\kappa}(\gamma_1) - \eta_{i-(l-1)\cdot\kappa}(\gamma_2)|
\end{aligned}$$

(where $\xi \in J_{N_{i-j\cdot\kappa}}$ is from the Mean Value Theorem and we made use of (ii)) until one of the base cases above is reached.

We now suppose without loss of generality that $\eta_i(\gamma_1) \ge \eta_i(\gamma_2)$. Using the inequality $1 + \xi < e^\xi$ for any $\xi > 0$ with $*_2$ gives

$$\begin{aligned}
0 &< \log \frac{\eta_i(\gamma_1)}{\eta_i(\gamma_2)} \\
&< \log\left(1 + m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}\right) < \log(e^{m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}}) = m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.
\end{aligned}$$

On the other hand, the inequalities $\frac{1}{1-\xi} > e^\xi$ and $1 > \frac{\eta_i(\gamma_2)}{\eta_i(\gamma_1)} > \frac{1}{1 + m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}}$ give

$$\begin{aligned}
0 > \log \frac{\eta_i(\gamma_2)}{\eta_i(\gamma_1)} &> \log\left(\frac{1}{1 + m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}}\right) \\
&> \log(e^{-m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}}) = -m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.
\end{aligned}$$

Combining these together yields

$$\left|\log \frac{\eta_i(\gamma_1)}{\eta_i(\gamma_2)}\right| < m_g \cdot \sigma \cdot \tau^{1+\frac{i+1-(n+m)}{\kappa}}.$$

■

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Proof of (3) Let $(\delta(m))_{m=1}^{\infty}$ be a sequence of positive numbers such that $|\eta_i(\beta^I) - \eta_i(\omega)| < \delta(m)$ for all $0 < i < n+m-1$ and $N \in \mathbb{N}$, and $\lim_{m\to\infty} \delta(m) = 0$. We claim that $\eta_i(\beta^I)$ is uniformly bounded away from 0 and 1 for all I and $0 < i < n+m-1$. Note that

$$\begin{aligned}
\eta_i(\beta^I) &= [1,1,\ldots] && \text{for } n < i < n+m-1,\\
\eta_i(\beta^I) &= [a_n,1,\ldots] && \text{for } i = n, \text{ and}\\
\eta_i(\beta^I) &= [a_i,a_{i+1},\ldots] && \text{for } 0 < i < n.
\end{aligned}$$

Call these first two digits p_i and q_i, noting that $p_i = q_i = p_{n+1} = q_{n+1}$ if $n < i < n+m-1$. Let

$$\begin{aligned}
\eta_{\min} &= \min_{0<i<n+m-1}[p_i,q_i,\ldots] = \min_{0<i\le n+1}[p_i,q_i,\ldots] = \min_{0<i\le n+1}(G_{p_i}^{-1}\circ G_{q_i}^{-1})((s_0,s_1)),\\
\eta_{\max} &= \max_{0<i<n+m-1}[p_i,q_i,\ldots] = \max_{0<i\le n+1}[p_i,q_i,\ldots] = \max_{0<i\le n+1}(G_{p_i}^{-1}\circ G_{q_i}^{-1})((s_0,s_1)).
\end{aligned}$$

It is clear that if $p_i \neq 1$ or $q_i \neq 1$, then $s_0 < \eta_{\min} < \eta_{\max} < s_1$. If $p_i = q_i = 1$, by assumption (iii) on G, we have

$$(G_{p_i}^{-1}\circ G_{q_i}^{-1})((s_0,s_1)) = G_{p_i}^{-1}(G_{q_i}^{-1}(s_1),s_1) = (G_{q_i}^{-1}(s_1),(G_{p_i}^{-1}\circ G_{q_i}^{-1})(s_1))$$

with $s_0 < G_{q_i}^{-1}(s_1) < (G_{p_i}^{-1}\circ G_{q_i}^{-1})(s_1) < s_1$. Hence, $s_0 < \eta_{\min} < \eta_{\max} < s_1$ in all cases. Thus,

$$\eta_i(\beta^I) \in [\eta_{\min},\eta_{\max}] \subseteq (s_0,s_1).$$

Clearly, $\eta_i(\omega) \in [\eta_{\min},\eta_{max}]$ as well. Now, letting

$$\tilde{u} = \inf_{x\in[\eta_{\min},\eta_{\max}]} u(x),$$

we have $\tilde{u} > 0$ since u is continuous and positive on the compact subset $[\eta_{\min},\eta_{\max}]$.

Since u is continuous, it is uniformly continuous on $[\eta_{\min},\eta_{\max}]$, hence admits a continuous and strictly increasing modulus of continuity $\sigma : [0,\infty) \to [0,\infty)$ such that $\lim_{t\to 0}\sigma(t) = \sigma(0) = 0$ and for all $x_1, x_2 \in [\eta_{\min},\eta_{\max}]$, $|u(x_1) - u(x_2)| \le \sigma(|x_1 - x_2|)$.

Now, fix m large enough that $1 - \dfrac{\sigma(\delta(m))}{\tilde{u}} > 0$, and let $0 < i < n+m-1$ and $N \in \mathbb{N}$. Then,

$$\begin{aligned}
&|u(\eta_i(\beta^I)) - u(\eta_i(\omega))| \le \sigma(|\eta_i(\beta^I) - \eta_i(\gamma_2)|) < \sigma(\delta(m))\\
&\quad\Longrightarrow \left|\frac{u(\eta_i(\beta^I))}{u(\eta_i(\omega))} - 1\right| < \frac{\sigma(\delta(m))}{u(\eta_i(\omega))} \le \frac{\sigma(\delta(m))}{\tilde{u}}\\
&\quad\Longrightarrow \left|\log\left(\frac{u(\eta_i(\beta^I))}{u(\eta_i(\omega))}\right)\right| < \max\left\{\left|\log\left(1 - \frac{\sigma(\delta(m))}{\tilde{u}}\right)\right|, \left|\log\left(1 + \frac{\sigma(\delta(m))}{\tilde{u}}\right)\right|\right\}\\
&\quad=: f(m),
\end{aligned}$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

with

$$\lim_{m\to\infty} f(m) = 0 \text{ since } \lim_{\delta(m)\to 0^+} \log\left(1 \pm \frac{\sigma(\delta(m))}{\tilde{u}}\right) = 0$$

and $f(m) > 0$ since $\dfrac{\sigma(\delta(m))}{\tilde{u}} > 0$ for all m. ∎

Lemma A.4 *There is some $\rho < 1$ such that for any $k > 1$ and $x \in \Lambda$, $\eta_{k-1}(x) \cdot \eta_k(x) < \rho$.*

Proof If $s_1 < 1$, we can take $\rho = s_1$, so suppose $s_1 = 1$. By the decreasing assumption (iii) on G_1, it is easily checked that there are some $a_*, b_* \in (s_0, s_1)$ for which $G(S \cap [a_*, 1)) \subseteq (s_0, b_*)$.

- If $G^{k-1}(x) < a_*$, then since $G^k(x) < 1$, we have $G^k(x) \cdot G^{k-1}(x) < a_* < 1$.
- If $a_* \le G^{k-1}(x) < 1$, then $G^k(x) = G(G^{k-1}(x)) < b_*$ and so $G^k(x) \cdot G^{k-1}(x) < b_* < 1$.

Taking $\rho = \max\{a_*, b_*\}$ completes the proof. ∎

A.3 Proof of Lemmas 4.2 and 4.3

Lemma A.5 *There is a constant D such that for any $\gamma_1, \gamma_2 \in \Lambda$ with $\gamma_1 = [a, \ldots]$, $\gamma_2 = [a, \ldots]$ for some $a \in \mathbb{Z}^+$, we have*

$$|u(\gamma_1) - u(\gamma_2)| < D.$$

Proof If $\gamma_1 = \gamma_2$, we are done; suppose $\gamma_1 < \gamma_2$. Define $x(a) := \gamma_1$ and $h(a) = \gamma_2 - \gamma_1 > 0$. By Taylor's theorem,

$$|u(\gamma_1) - u(\gamma_2)| = h(a) \cdot |u'(x(a)) + R_1(x(a))| \le \frac{h(a)}{(x(a) - s_0)^2} + h(a) \cdot |R_1(x(a))|,$$

where we used assumption (iii) on u and where R_1 is the first-order Taylor remainder. But now

$$\frac{h(a)}{(x(a) - s_0)^2} = \frac{\gamma_2 - \gamma_1}{(\gamma_1 - s_0)^2} < \frac{r_a - \ell_a}{(\ell_a - s_0)^2} < C_1$$

for some constant C_1 independent of a, where we used assumption (v) on G. Since

$$h(a) \cdot |R(x(a))| \to 0 \text{ as } a \to \infty,$$

$h(a) \cdot |R(x(a))| < C_2$ for some constant C_2. Taking $D = C_1 + C_2$ completes the proof. ∎

We are now ready to prove Lemmas 4.2 and 4.3.

Proof of Lemma 4.2 We will first show that such an m_0 exists, and then give an algorithm to compute it.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Note that the sum in the expression for $\Phi(\omega)$ converges, because its tail converges:

$$\sum_{i=n+1}^{\infty} (\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega))$$
$$\le u(\eta_{n+1}(\omega))\cdot(\eta_n(\omega))^{\nu}\cdot\sum_{i=n+1}^{\infty}(\eta_{n+1}(\omega))^{\nu}<\infty$$

since $(\eta_{n+1}(\omega))^{\nu} < s_1 \le 1$ for $\nu > 0$. Hence, there is an $m_1 > 1$ such that the tail of the sum

$$\sum_{i\ge n+m_1}(\eta_1(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega)) < \frac{\varepsilon}{2}.$$

If needed, additionally make m_1 large enough that Lemma A.3 (3) applies for all $m \ge m_1$. We will show how to choose $m_0 > m_1$ to satisfy the conclusion of the lemma.

By Lemma A.3 (2) and (3), for any β^I and any $i \le n + m_1$, we have

$$\left|\log\frac{\left(\eta_0(\beta^I)\dots\eta_{i-1}(\beta^I)\right)^{\nu}\cdot u(\eta_i(\beta^I))}{(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega))}\right|$$
$$=\left|\nu\cdot\log\frac{\eta_0(\beta^I)\dots\eta_{i-1}(\beta^I)}{\eta_0(\omega)\dots\eta_{i-1}(\omega)}+\log\frac{u(\eta_i(\beta^I))}{u(\eta_i(\omega))}\right|$$
$$<\nu\cdot\left(\sum_{j=1}^{i-1}m_g\cdot\sigma\cdot\tau^{1+\frac{j+1-(n+m)}{\kappa}}\right)+f(m)<\nu\sigma\cdot m_g\cdot\tau^{1+\frac{2-(n+m)}{\kappa}}\left(\frac{\tau^i-1}{\tau^{1/\kappa}-1}\right)+f(m)$$
$$<\nu\sigma\cdot m_g\cdot\tau^{1+\frac{2-(n+m)}{\kappa}}\left(\frac{\tau^{n+m_1}}{\tau^{1/\kappa}-1}\right)+f(m)\le\nu\sigma\cdot m_g\cdot\frac{\tau^{1+m_1+\frac{2-m}{\kappa}}}{\tau^{1/\kappa}-1}+f(m).$$

We can choose m_0 sufficiently large so that $\exp(\nu\sigma\cdot m_g\cdot\frac{\tau^{1+m_1+\frac{2-m}{\kappa}}}{\tau^{1/\kappa}-1}+f(m))>1-\frac{\varepsilon}{2\cdot\Phi(\omega)}$ for $m \ge m_0$ and also

$$\left(\eta_1(\beta^I)\dots\eta_{i-1}(\beta^I)\right)^{\nu}\cdot u(\eta_i(\beta^I))$$
$$>\left(1-\frac{\varepsilon}{2\cdot\Phi(\omega)}\right)\cdot(\eta_1(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega))$$

for $i \le n + m_1$. Now, for any β^I, we have

$$\Phi(\beta^I)\ge\sum_{i=1}^{n+m_1-1}\left(\eta_1(\beta^I)\dots\eta_{i-1}(\beta^I)\right)^{\nu}\cdot u(\eta_i(\beta^I))$$
$$>\sum_{i=1}^{n+m_1-1}\left(1-\frac{\varepsilon}{2\cdot\Phi(\omega)}\right)\cdot(\eta_1(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega))$$
$$>\left(1-\frac{\varepsilon}{2\cdot\Phi(\omega)}\right)\left(\Phi(\omega)-\frac{\varepsilon}{2}\right)>\Phi(\omega)-\varepsilon.$$

Since the symbolic representation of ω ends with all ones, for any specific $(a_1, a_2, \dots, a_n)$, we can compute $\Phi(\omega)$ by Proposition 4.4. This allows us to compute

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

m_1 by iteratively checking whether

$$\sum_{i\geq n+m_1} (\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega)) < \frac{\varepsilon}{2}$$

for each $m_1 > 1$. Then, for each candidate $m_0 > m_1$, since we can compute $\Phi(\omega)$ and $(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega))$ and left-compute $\left(\eta_0(\beta^I)\dots\eta_{i-1}(\beta^I)\right)^{\nu}\cdot u(\eta_i(\beta^I))$ for each individual i by assumption (v) on u and computability assumption (vi) on G, we can iteratively check whether the following two conditions hold:

(1) $$\exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau^{1+m_1+\frac{2-m}{\kappa}}}{\tau^{1/\kappa}-1}+f(m)\right) > 1-\frac{\varepsilon}{2\cdot\Phi(\omega)}.$$

(2) For every $i \leq n+m_1$,

$$\left(\eta_0(\beta^I)\dots\eta_{i-1}(\beta^I)\right)^{\nu}\cdot u(\eta_i(\beta^I)) > \left(1-\frac{\varepsilon}{2\cdot\Phi(\omega)}\right)\cdot(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega)).$$

We only need to be able to left-compute $\left(\eta_0(\beta^I)\dots\eta_{i-1}(\beta^I)\right)^{\nu}\cdot u(\eta_i(\beta^I))$ because of the strict inequality in condition (2). Since we have shown that there must exist $m_1 > m_0$ for which these hold, this algorithm terminates and we eventually find the m_1 we are looking for. ∎

Proof of Lemma 4.3 Let $i \leq k-1$ for some integer $k > 1$. We will first show that

$$\rho_1 < \frac{u(\eta_i(\omega_k))}{u(\eta_i(\omega))} < \rho_2$$

for some $0 < \rho_1 < \rho_2 < \infty$, independent of i and k. Since $i \leq k-1$, the symbolic representations of $\eta_i(\omega_k), \eta_i(\omega)$ coincide in the first two terms, call them a and b. So, we can write

$$x := \eta_i(\omega_k) = [a, b, T_1],$$
$$y := \eta_i(\omega) = [a, b, T_2],$$

where T_1, T_2 are the infinite tails of the symbolic expansions of $\eta_i(\omega_k), \eta_i(\omega)$ respectively. Since u is $\mathcal{C}^1$, we have

$$u(y) = u(x) + \int_x^y u'(t)\,dt$$

and so by assumption (iii) on u and (v) on G,

$$\left|\frac{u(y)}{u(x)}-1\right| = \frac{1}{u(x)}\left|\int_x^y u'(t)\,dt\right| \leq \frac{1}{u(x)}\left|\int_x^y \frac{C}{(t-s_0)^2}\,dt\right|$$
$$= \frac{C}{u(x)}\left|\frac{1}{x-s_0}-\frac{1}{y-s_0}\right|$$
$$< \frac{C}{u(x)}\left(\frac{1}{\ell_i-s_0}-\frac{1}{r_i-s_0}\right) < \frac{C}{u(x)}\cdot\frac{r_i-\ell_i}{(\ell_i-s_0)^2} < \frac{\tilde{C}}{u(x)}.$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

for some constant $\tilde{C} > 0$. By assumption (i) on u, there is some $l_1 \in (0,1)$ such that $\frac{1}{u(\xi)} < \frac{1}{2\tilde{C}}$ for $\xi \in (0, l_1)$. Thus, if a is large enough such that $x, y \in (0, l_1)$, say $a \geq N_1 \in \mathbb{Z}^+$, then

$$\left| \frac{u(x)}{u(y)} - 1 \right| < \frac{\tilde{C}}{2\tilde{C}} = \frac{1}{2}.$$

If $1 < a < N_1$, then $x, y \in [l_1', l_2']$ for some $0 < l_1' < l_2' < 1$, thus $u(x), u(y)$ are bounded away from 0 and ∞, so $\rho_1' < \frac{u(x)}{u(y)} < \rho_2'$ for some $0 < \rho_1' < \rho_2' < \infty$.

Now, suppose $a = 1$. If $s_1 < 1$, then $x, y \in J_1$ with J_1 bounded away from 0 and 1, thus u is bounded away from 0 and infinity on J_1. Now, suppose $s_1 = 1$. It follows immediately from assumption (ii) on u that for large enough b, say $b \geq N_2$, $\frac{u(x)}{u(y)}$ is bounded away from 0 and infinity. Therefore, $\rho_1'' < \frac{u(x)}{u(y)} < \rho_2''$ for some $0 < \rho_1'' < \rho_2'' < \infty$.

Putting all this together, letting $\rho_1 = \min\{\frac{1}{2}, \rho_1', \rho_1''\} > 0$ and $\rho_2 = \max\{\frac{1}{2}, \rho_2', \rho_2''\} < \infty$, we have for any ω_k and ω that $\rho_1 < \frac{u(\omega_k)}{u(\omega)} < \rho_2$ and so $\left|\log \frac{u(\omega_k)}{u(\omega)}\right| < D_1 < \infty$ for some constant $D_1 > 0$.

Returning to the proof of the lemma, since the sum in the expression for $\Phi(\omega)$ converges, we can split this sum as

$$\Phi(\omega) = \underbrace{\sum_{i=1}^{s} \left(\eta_0(\omega) \dots \eta_{i-1}(\omega)\right)^{\nu} \cdot u(\eta_i(\omega))}_{\text{“head”}} + \underbrace{\sum_{i=s+1}^{\infty} \left(\eta_0(\omega) \dots \eta_{i-1}(\omega)\right)^{\nu} \cdot u(\eta_i(\omega))}_{\text{“tail”}}$$

so that “tail” $< \dfrac{\varepsilon}{2\left(\exp\left(\nu\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1} + D_1\right)\right)}$. If needed, additionally make s large enough so that

$$\frac{\rho^{\nu s/2}}{1-\rho^{\nu/2}} < \frac{\varepsilon}{4\left(\exp\left(\nu\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1}\right) \cdot D_2 + u(\varphi)\right)},$$

where $\varphi = [1, 1, \dots]$, ρ is as in Lemma A.4, and D_2 is the constant from Lemma A.5.

Since $\sum_{i=1}^{s} \left(\eta_0(\cdot) \dots \eta_{i-1}(\cdot)\right)^{\nu} \cdot u(\eta_i(\cdot))$ is a finite sum of functions continuous on a neighborhood of ω, it is continuous on some interval containing ω. Noting that $\omega_k \to \omega$, there is some $m > s$ such that for any $k \geq m$,

$$\sum_{i=1}^{s} \left(\eta_0(\omega_k) \dots \eta_{i-1}(\omega_k)\right)^{\nu} \cdot u(\eta_i(\omega_k)) < \sum_{i=1}^{s} \left(\eta_0(\omega) \dots \eta_{i-1}(\omega)\right)^{\nu} \cdot u(\eta_i(\omega)) + \varepsilon/4.$$

We now want to bound the change from

$$\left(\eta_0(\omega_k) \dots \eta_{i-1}(\omega_k)\right)^{\nu} \cdot u(\eta_i(\omega_k)) \text{ to } \left(\eta_0(\omega) \dots \eta_{i-1}(\omega)\right)^{\nu} \cdot u(\eta_i(\omega))$$

for $i > s$. We consider these terms individually for $s < i \leq k-1$, $i = k$, and $i \geq k+1$.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

- For $s < i \le k-1$. Taking $n+m := k+1$ in Lemma A.3 (2), since ω_k and ω coincide in the first $k = n+m-1$ terms, we have

$$\begin{aligned}
&\left|\log \frac{(\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k))^{\nu}\cdot u(\eta_i(\omega_k))}{(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega))}\right| \\
&= \left|\nu\cdot\log\frac{\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k)}{\eta_0(\omega)\dots\eta_{i-1}(\omega)} + \log\frac{u(\eta_i(\omega_k))}{u(\eta_i(\omega))}\right| \\
&< \nu\cdot\left(\sum_{j=1}^{i-1} m_g\cdot\sigma\cdot\tau^{1+\frac{j-k}{\kappa}}\right) + D_1 < \nu\sigma\cdot m_g\cdot\frac{\tau^{1+\frac{i-k}{\kappa}}}{\tau^{1/\kappa}-1} + D_1 \le \nu\sigma\cdot m_g\cdot\frac{\tau^{1-\frac{1}{\kappa}}}{\tau^{1/\kappa}-1} + D_1.
\end{aligned}$$

  Hence, in this case, each term can increase by a factor of at most $\exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau^{1-\frac{1}{\kappa}}}{\tau^{1/\kappa}-1} + D_1\right)$.

- For $i = k$. We have

$$|u(\eta_i(\omega)) - u(\eta_i(\omega_k))| = |u([a_k,\dots]) - u([a_k,\dots])| < D_2,$$

  where we had previously gotten D_2 from Lemma A.5. Now, by Lemma A.3 (2),

$$\begin{aligned}
\left|\log\frac{(\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k))^{\nu}}{(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}}\right| &= \nu\cdot\left|\log\frac{\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k)}{\eta_0(\omega)\dots\eta_{i-1}(\omega)}\right| \\
&< \nu\cdot\left(\sum_{j=1}^{i-1} m_g\cdot\sigma\cdot\tau^{1+\frac{j-k}{\kappa}}\right) \\
&< \nu\sigma\cdot m_g\cdot\frac{\tau^{1+\frac{i-k}{\kappa}}}{\tau^{1/\kappa}-1} = \nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1},
\end{aligned}$$

  so $(\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k))^{\nu} < \exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1}\right)\cdot(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}$. Combining these estimates and using the value $\rho < 1$ from Lemma A.4, we get

$$\begin{aligned}
&(\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k))^{\nu}\cdot u(\eta_i(\omega_k)) \\
&\le \exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1}\right)\cdot(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot(u(\eta_i(\omega)) + D_2) \\
&< \exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1}\right)\cdot(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega)) \\
&\quad + \exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1}\right)\cdot D_2\cdot\rho^{\nu(i-1)/2}.
\end{aligned}$$

- For $i \ge k+1$. Noting that $u(\eta_i(\omega_k)) = u(\eta_i(\omega_k))$, by Lemma A.4, we have

$$(\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k))^{\nu}\cdot u(\eta_i(\omega_k)) < \rho^{\nu(i-1)/2}\cdot u(\varphi).$$

Therefore, for any $i > s$,

$$\begin{aligned}
&(\eta_0(\omega_k)\dots\eta_{i-1}(\omega_k))^{\nu}\cdot u(\eta_i(\omega_k)) \\
&< \exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1} + D_1\right)\cdot(\eta_0(\omega)\dots\eta_{i-1}(\omega))^{\nu}\cdot u(\eta_i(\omega)) \\
&\quad + \left(\exp\left(\nu\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1}\right)\cdot D_2 + u(\varphi)\right)\cdot\rho^{\nu(i-1)/2}.
\end{aligned}$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Finally, noting that

$$\sum_{i=s+1}^{\infty} \rho^{v(i-1)/2} = \frac{\rho^{vs/2}}{1-\rho^{v/2}},$$

for such a k, we have

$$\begin{aligned}
\Phi(\omega_k) &= \text{“head”}(\omega_k) + \text{“tail”}(\omega_k) \\
&< \left(\text{“head”}(\omega) + \frac{\varepsilon}{4}\right) + \left(\exp\left(v\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1} + D_1\right) \cdot \text{“tail”}(\omega)\right. \\
&\quad \left. + \left(\exp\left(v\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1}\right) \cdot D_2 + u(\varphi)\right) \cdot \frac{\rho^{vs/2}}{1-\rho^{v/2}}\right) \\
&< \text{“head”}(\omega) + \frac{\varepsilon}{4} + \exp\left(v\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1} + D_1\right) \cdot \frac{\varepsilon}{2\left(\exp\left(v\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1} + D_1\right)\right)} \\
&\quad + \left(\exp\left(v\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1}\right) \cdot D_2 + u(\varphi)\right) \cdot \frac{\varepsilon}{4\left(\exp\left(v\sigma \cdot m_g \cdot \frac{\tau}{\tau^{1/\kappa}-1}\right) \cdot D_2 + u(\varphi)\right)} \\
&= \text{“head”}(\omega) + \frac{\varepsilon}{4} + \frac{\varepsilon}{2} + \frac{\varepsilon}{4} < \Phi(\omega) + \varepsilon.
\end{aligned}$$

■

A.4 Proof of Lemma 4.1

Write

$$\Phi^-(\omega) = \Phi(\omega) - s(n+m) \cdot (\eta_1(\omega)\eta_2(\omega)\dots\eta_{n+m-1}(\omega))^v \cdot u(\eta_{n+m}(\omega)).$$

The value of the integer $m > 0$ is yet to be determined.

Lemma A.6 *For any ω of the form as in Lemma 4.1 and for any $\varepsilon > 0$, there is an $m_0 > 0$ such that, for any N and any $m \geq m_0$,*

$$|\Phi^-(\beta^N) - \Phi^-(\beta^1)| < \frac{\varepsilon}{4}.$$

Proof Noting that $\eta_{n+m-1}(\beta^N) \in (\ell_1, s_1)$ for all $N \in \mathbb{Z}^+$, by Lemma A.2, we have some $C > 0$ such that $0 \leq u(\eta_{n+m-1}(\beta^N)) < C$ for all N. Now, the sum in the expression for $\Phi(\beta^1)$ converges because its tail converges absolutely:

$$\sum_{i=n+1}^{\infty} \left|s(i) \cdot \left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^v \cdot u(\eta_i(\beta^1))\right| = u(\beta^1) \cdot \sum_{i=n+1}^{\infty} \left((\beta^1)^v\right)^i < \infty$$

since $(\beta^1)^v < s_1 \leq 1$ for $v > 0$. Hence, there is an $m_1 > 1$, make it large enough that Lemma A.3 (3) applies for all $m \geq m_1$, such that the tail of the sum satisfies

$$\sum_{i \geq n+m_1} s(i) \cdot \left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^v \cdot u(\eta_i(\beta^1)) < \frac{\varepsilon}{4(1+D)},$$

where $M > 0$ is as in Lemma A.3 (3), and D is defined as

$$D = \exp\left(v\left(\frac{m_g \sigma \tau^3}{\tau^{1/\kappa}-1} + \left|\log\left(\frac{C}{u(\ell_1)}\right)\right|\right) + M\right).$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

We will now show how to choose $m_0 > m_1$ to satisfy the conclusion of the lemma.

We bound the influence of the change from β^1 to β^N using Lemma A.3 (2) and (3). The influence on each of the "head elements" ($i < n + m_1$) is bounded by

$$\begin{aligned}
&\left|\log \frac{s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^{\nu}\cdot u(\eta_i(\beta^1))}{s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^{\nu}\cdot u(\eta_i(\beta^N))}\right| \\
&\quad= \left|\nu\cdot\log\frac{\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)}{\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)} + \log\frac{u(\eta_i(\beta^1))}{u(\eta_i(\beta^N))}\right| \\
&\quad< \nu\cdot\left(\sum_{j=1}^{i-1} m_g\cdot\sigma\cdot\tau^{1+\frac{j+1-(n+m)}{\kappa}}\right) + f(m) \le \nu\sigma\cdot m_g\cdot\frac{\tau^{m_1+\frac{2-m}{\kappa}}}{\tau^{1/\kappa}-1} + f(m).
\end{aligned}$$

By making m sufficiently large (i.e., by choosing a sufficiently large m_0), we can ensure that

$$\begin{aligned}
1-\frac{\varepsilon}{4(1+D)\cdot\Phi(\beta^1)} < \frac{s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^{\nu}\cdot u(\eta_i(\beta^N))}{s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^{\nu}\cdot u(\eta_i(\beta^1))} < 1 \\
+\frac{\varepsilon}{4(1+D)\cdot\Phi(\beta^1)}.
\end{aligned}$$

Hence,

$$\begin{aligned}
&\Big|s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^{\nu}\cdot u(\eta_i(\beta^N)) \\
&\qquad - s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^{\nu}\cdot u(\eta_i(\beta^1))\Big| \\
&\quad< \frac{\varepsilon}{4(1+D)\cdot\Phi(\beta^1)}\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^{\nu}\cdot u(\eta_i(\beta^1)).
\end{aligned}$$

Adding the inequalities for $i = 1, 2, \dots, n + m_1 - 1$, we obtain

$$\begin{aligned}
&\left|\sum_{i=1}^{n+m_1-1} s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^{\nu}\cdot u(\eta_i(\beta^N))\right. \\
&\qquad\left. - \sum_{i=1}^{n+m_1-1} s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^{\nu}\cdot u(\eta_i(\beta^1))\right| \\
&< \frac{\varepsilon}{4(1+D)\cdot\Phi(\beta^1)}\sum_{i=1}^{n+m_1-1}\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^{\nu}\cdot u(\eta_i(\beta^1)) = \frac{\varepsilon}{4(1+D)}.
\end{aligned}$$

Thus, the influence on the "head" of Φ^- is bounded by $\frac{\varepsilon}{4(1+D)}$. To bound the influence on the "tail," we consider three kinds of terms $s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^{\nu}\cdot u(\eta_i(\beta^N))$: those for which $n + m_1 \le i \le n + m - 2$, $i = m + n - 1$, and $i \ge m + n + 1$ (recall that the $i = n + m$ term is not in Φ^-).

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

- For $n+m_1 \le i \le n+m-2$. By Lemma A.3 (2) and (3), and by the work above,

$$\begin{aligned}&\left|\log \frac{s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^v\cdot u(\eta_i(\beta^1))}{s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^v\cdot u(\eta_i(\beta^N))}\right|\\ &< v\cdot\left(\sum_{j=1}^{i-1} m_g\cdot\sigma\cdot\tau^{1+\frac{j+1-(n+m)}{\kappa}}\right)+f(m)\\ &\le v\sigma\cdot m_g\cdot\tau^{1+\frac{2-(n+m)}{\kappa}}\left(\frac{\tau^i}{\tau^{1/\kappa}-1}\right)+f(m)\le v\sigma\cdot m_g\cdot\frac{\tau}{\tau^{1/\kappa}-1}+M.\end{aligned}$$

  Where $M>0$ is as in Lemma A.3 (3). Hence, in this case, each term can increase or decrease by a factor of at most $\exp\left(\frac{v\sigma m_g\tau}{\tau^{1/\kappa}-1}+M\right)$.
- For $i=n+m-1$. Recall that for all $N\in\mathbb{Z}^+$, $0\le u(\eta_{n+m-1}(\beta^N))<C$. Thus, we have

$$\begin{aligned}&\log\frac{s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^v\cdot u(\eta_i(\beta^N))}{s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^v\cdot u(\eta_i(\beta^1))}\le\log\frac{\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^v\cdot C}{\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^v\cdot u(\ell_1)}\\ &<v\cdot\left(\sum_{j=1}^{n+m-2} m_g\cdot\sigma\cdot\tau^{1+\frac{j+1-(n+m)}{\kappa}}+\log\left(\frac{C}{u(\ell_1)}\right)\right)<v\cdot\left(\frac{m_g\cdot\sigma\cdot\tau^2}{\tau^{1/\kappa}-1}+\log\left(\frac{C}{u(\ell_1)}\right)\right).\end{aligned}$$

  Hence, this term could increase or decrease by a factor of at most $\exp\left(v\cdot\left(\frac{m_g\sigma\tau^2}{\tau^{1/\kappa}-1}+\log\left(\frac{C}{u(\ell_1)}\right)\right)\right)$.
- For $i\ge n+m+1$. Note that the η_j for $j>n+m$ are not affected by the change, and the change decreases η_{n+m}, so that $\eta_{n+m}(\beta^N)\le\eta_{n+m}(\beta^1)$ and thus $\left(\eta_{n+m}(\beta^N)\right)^v\le\left(\eta_{n+m}(\beta^1)\right)^v$. Hence,

$$\begin{aligned}&\log\frac{s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^v u(\eta_i(\beta^N))}{s(i)\cdot\left(\eta_0(\beta^1)\dots\eta_{i-1}(\beta^1)\right)^v u(\eta_i(\beta^1))}=\log\frac{\left(\eta_0(\beta^N)\dots\eta_{n+m}(\beta^N)\right)^v}{\left(\eta_0(\beta^1)\dots\eta_{n+m}(\beta^1)\right)^v}\\ &\le\log\frac{\left(\eta_0(\beta^N)\dots\eta_{n+m-1}(\beta^N)\right)^v}{\left(\eta_0(\beta^1)\dots\eta_{n+m-1}(\beta^1)\right)^v}<v\cdot\sum_{j=1}^{n+m-1} m_g\cdot\sigma\cdot\tau^{1+\frac{j+1-(n+m)}{\kappa}}<\frac{v\cdot m_g\cdot\sigma\cdot\tau^3}{\tau^{1/\kappa}-1}.\end{aligned}$$

  So, in this case, each term could increase or decrease by a factor of at most $\exp\left(\frac{vm_g\sigma\tau^3}{\tau^{1/\kappa}-1}\right)$.

We see that after the change, each term of the tail could increase or decrease by a factor of

$$\exp\left(v\left(\frac{m_g\sigma\tau^3}{\tau^{1/\kappa}-1}+\left|\log\left(\frac{C}{u(\ell_1)}\right)\right|\right)+M\right)=:D$$

at most. So the value of the tail remains in the interval $\left[-\frac{D\cdot\varepsilon}{4(1+D)},\frac{D\cdot\varepsilon}{4(1+D)}\right]$, hence the change in the tail is bounded by $\frac{D\cdot\varepsilon}{4(1+D)}$.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Therefore, the total change in Φ^- is bounded by

$$\text{change in the "head"} \;+\; \text{change in the "tail"} < \frac{\varepsilon}{4(1+D)} + \frac{D\cdot\varepsilon}{4(1+D)} = \frac{\varepsilon}{4}.$$

■

Lemma A.7 *For any ε and for the same $m_0(\varepsilon)$ as in Lemma A.6, for any $m \geq m_0$ and N,*

$$|\Phi^-(\beta^N) - \Phi(\beta^{N+1})| < \frac{\varepsilon}{2}.$$

Proof We have

$$|\Phi^-(\beta^N) - \Phi(\beta^{N+1})| \leq |\Phi^-(\beta^N) - \Phi(\beta^1)| + |\Phi^-(\beta^{N+1}) - \Phi(\beta^1)| < \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \frac{\varepsilon}{2}.$$

■

We will now take a closer look at the term

$$\Phi^1(\omega) := s(n+m)\cdot(\eta_1(\omega)\dots\eta_{n+m-1}(\omega))^\nu \cdot u(\eta_{n+m}(\omega)) = \Phi(\omega) - \Phi^-(\omega).$$

Lemma A.8 *There exists some m, which can be made arbitrarily large, such that for any N,*

$$\Phi^1(\beta^{N+1}) - \Phi^1(\beta^N) < \frac{\varepsilon}{2}.$$

Proof We assume throughout that $s(n+m) = 1$. By assumption on $s(\cdot)$, there are infinitely many m such that this holds, so we can take m arbitrarily large within the course of the proof.

According to Lemma A.3 (1),

$$\left|\log \frac{s(n+m)\cdot\left(\eta_0(\beta^{N+1})\dots\eta_{n+m-1}(\beta^{N+1})\right)^\nu}{s(n+m)\cdot(\eta_0(\beta^N)\dots\eta_{n+m-1}(\beta^N))^\nu}\right| < \nu\cdot\sum_{i=1}^{n+m-1}\frac{g(N)}{\ell_1}\cdot\tau^{1+\frac{i-(n+m)}{\kappa}}$$
$$= \nu\cdot g(N)\cdot\tau^{1+\frac{1-(n+m)}{\kappa}}\cdot\frac{\tau^{n+m}-1}{\tau^{1/\kappa}-1} < \nu\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}.$$

Hence,

$$s(n+m)\cdot\left(\eta_0(\beta^{N+1}\dots\eta_{n+m-1}(\beta^{N+1}))\right)^\nu$$
$$< s(n+m)\cdot\left(\eta_0(\beta^N\dots\eta_{n+m-1}(\beta^N))\right)^\nu\cdot\exp\left(\nu\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}\right)$$

and

$$\Phi^1(\beta^{N+1}) < \Phi^1(\beta^N)\cdot\exp\left(\nu\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}\right)\cdot\frac{u(\eta_{n+m}(\beta^{N+1}))}{u(\eta_{n+m}(\beta^N))}.$$

We require an auxiliary inequality. It is straightforward to show that for any $y\in(0,1]$ and any $d > e^c/c$, we have $e^{cy} < 1 + dcy$. Taking $y = g(N)/(m_g\cdot\frac{\tau^2}{\tau^{1/\kappa}-1})$ and $c = \nu\cdot m_g\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}$, for any $d > e^{\nu\cdot m_g\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}}/(\nu\cdot m_g\cdot\frac{\tau^2}{\tau^{1/\kappa}-1})$, we have $e^{\nu\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}} < 1 +$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

$dv \cdot g(N) \cdot \frac{\tau^2}{\tau^{1/\kappa}-1}$. Hence, for any such d,

$$\Phi^1(\beta^{N+1}) - \Phi^1(\beta^N) < \Phi^1(\beta^N)\left(e^{v\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}} \cdot \frac{u(\eta_{n+m}(\beta^{N+1}))}{u(\eta_{n+m}(\beta^N))} - 1\right)$$
$$< \Phi^1(\beta^N)\left(\left(1 + dv \cdot g(N) \cdot \frac{\tau^2}{\tau^{1/\kappa}-1}\right) \cdot \frac{u(\eta_{n+m}(\beta^{N+1}))}{u(\eta_{n+m}(\beta^N))} - 1\right).$$

For $\rho \in (0,1)$ as in Lemma A.4, we have

$$\Phi^1(\beta^N) < \rho^{v\cdot(n+m-2)/2} \cdot u\left(\eta_{n+m}(\beta^N)\right).$$

Thus,

$$\Phi^1(\beta^{N+1}) - \Phi^1(\beta^N) < \rho^{v\cdot(n+m-2)/2} \cdot u\left(\eta_{n+m}(\beta^N)\right)$$
$$\cdot \left(\left(1 + dv \cdot g(N) \cdot \frac{\tau^2}{\tau^{1/\kappa}-1}\right) \cdot \frac{u(\eta_{n+m}(\beta^{N+1}))}{u(\eta_{n+m}(\beta^N))} - 1\right)$$
$$= \rho^{v\cdot(n+m-2)/2} \cdot \left(\left[u\left(\eta_{n+m}(\beta^{N+1})\right) - u\left(\eta_{n+m}(\beta^N)\right)\right]\right.$$
$$\left. + dv \cdot g(N) \cdot \frac{\tau^2}{\tau^{1/\kappa}-1} \cdot u\left(\eta_{n+m}(\beta^{N+1})\right)\right)$$

For the remainder of the proof, we will show that this product can be made less than $\frac{\varepsilon}{2}$ by choosing m large enough. Since the first term $\to 0$ as $m \to \infty$, it is enough to show that the product of the last two terms is bounded from above by a constant.

By assumption on u, we have $-u'(x) < \dfrac{C_1}{(x-s_0)^2}$ for some constant $C_1 > 0$. Integrating both sides gives $u(x) < \dfrac{C_2}{x-s_0}$ for another constant $C_2 > 0$. Letting $x(N) := \eta_{n+m}(\beta^{N+1})$ and $h(N) := \eta_{n+m}(\beta^N) - \eta_{n+m}(\beta^{N+1}) > 0$, we have

$$\frac{1}{h(N)} \cdot \left(\left[u\left(\eta_{n+m}(\beta^{N+1})\right) - u\left(\eta_{n+m}(\beta^N)\right)\right]\right.$$
$$\left. + dv \cdot g(N) \cdot \frac{\tau^2}{\tau^{1/\kappa}-1} \cdot u\left(\eta_{n+m}(\beta^{N+1})\right)\right)$$
$$= -\left[u'(x(N)) + R_1(x(N))\right] + \frac{x(N)\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}}{h(N)} \cdot \frac{dv \cdot u(x(N))}{x(N)},$$

where $R_1(\cdot)$ is the Taylor remainder term with $\lim_{N\to\infty} R_1(x(N)) = 0$.

We now want to find a lower bound on $h(N)$. Recalling assumption (iv) on G, we have

$$h(N) = [N,1,1,\ldots] - [N+1,1,1,\ldots] = G_N^{-1}(\varphi) - G_{N+1}^{-1}(\varphi) = \delta_G(N) > 0.$$

Taking $D > 0$ as in assumption (iv) and recalling the definition of g from (vii), we have

$$\frac{x(N)\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}}{h(N)} < \frac{\eta_{n+m}(\beta^{N+1})}{\delta_G(N)} \cdot \frac{r_N - \ell_{N+1}}{\ell_{N+1}} \cdot \frac{\tau^2}{\tau^{1/\kappa}-1} < D \cdot \frac{\tau^2}{\tau^{1/\kappa}-1}.$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

Since $\lim_{N\to\infty} R_1(x(N)) = 0$, we have $|R_1(x(N))| < C_3$ for some $C_3 > 0$. So,

$$\begin{aligned}&-\left[u'(x(N)) + R_1(x(N))\right] + \frac{x(N)\cdot g(N)\cdot \frac{\tau^2}{\tau^{1/\kappa}-1}}{h(N)}\cdot\frac{dv\cdot u(x(N))}{x(N)}\\ &< \frac{C_1}{(x(N))^2} + C_3 + \frac{D\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}\cdot dv}{x(N)}\cdot\frac{C_2}{(x(N))} < \frac{C_4}{(x(N))^2} < \frac{C}{h(N)}\end{aligned}$$

for some $C_4, C > 0$, where the last inequality holds true because

$$\begin{aligned}h(N) &< r_N - \ell_{N+1} < \ell_{N+1}\cdot g(N) < \eta_{n+m}(\beta^{N+1})\cdot g(N)\\ &= x(N)\cdot g(N) < (x(N))^2\cdot m_g.\end{aligned}$$

Therefore, we have shown that for all N,

$$\left[u\left(\eta_{n+m}(\beta^{N+1})\right) - u\left(\eta_{n+m}(\beta^N)\right)\right] + dv\cdot g(N)\cdot\frac{\tau^2}{\tau^{1/\kappa}-1}\cdot u\left(\eta_{n+m}(\beta^{N+1})\right) < C$$

for some constant $C > 0$, finishing the proof of the lemma. ∎

Lemmas A.7 and A.8 yield the following.

Lemma A.9 *There exists some m, which can be made arbitrarily large, such that for any N,*

$$\Phi(\beta^{N+1}) - \Phi(\beta^N) < \varepsilon.$$

Proof Using Lemmas A.7 and A.8, for sufficiently large m with $s(n+m) = 1$, we have

$$\Phi(\beta^{N+1}) - \Phi(\beta^N) \le \Phi^-(\beta^{N+1}) - \Phi^-(\beta^N) + \Phi^1(\beta^{N+1}) - \Phi^1(\beta^N) < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon.$$

∎

To complete the proof of Lemma 4.1, we will need the following statement.

Lemma A.10 *For any m satisfying $s(n+m) = 1$, we have*

$$\lim_{N\to\infty}\Phi(\beta^N) = \infty.$$

Proof We first prove that $\lim_{N\to\infty}\Phi^1(\beta^N) = \infty$. By Lemma A.3 (2),

$$\begin{aligned}\left|\log\frac{s(n+m)\cdot\left(\eta_0(\beta^N)\dots\eta_{n+m-1}(\beta^N)\right)^\nu}{s(n+m)\cdot\left(\eta_0(\beta^1)\dots\eta_{n+m-1}(\beta^1)\right)^\nu}\right| &< \nu\cdot\sum_{j=1}^{n+m-1} m_g\cdot\sigma\cdot\tau^{1+\frac{j+1-(n+m)}{\kappa}}\\ &< \frac{\nu\cdot m_g\cdot\sigma\cdot\tau^3}{\tau^{1/\kappa}-1}.\end{aligned}$$

Hence,

$$\begin{aligned}&s(n+m)\cdot\left(\eta_0(\beta^N)\dots\eta_{n+m-1}(\beta^N)\right)^\nu\\ &> \frac{1}{\exp\left(\frac{\nu m_g\sigma\tau^3}{\tau^{1/\kappa}-1}\right)}\cdot s(n+m)\cdot\left(\eta_0(\beta^1)\dots\eta_{n+m-1}(\beta^1)\right)^\nu\end{aligned}$$

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

and

$$\Phi^1(\beta^N) > \frac{1}{\exp\left(\frac{\nu m_g \sigma \tau^3}{\tau^{1/\kappa}-1}\right)} \cdot \frac{u(\eta_{n+m}(\beta^N))}{u(\eta_{n+m}(\beta^1))}\Phi^1(\beta^1).$$

Since $\lim_{x\to s_0} u(x) = \infty$, the latter expression tends to ∞ as $N \to \infty$. Now, write

$$\Phi(\beta^N) = \sum_{i=1}^{n+m-1} s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^\nu \cdot u(\eta_i(\beta^N)) + \Phi^1(\beta^N)$$
$$+ \sum_{i=n+m+1}^{\infty} s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^\nu \cdot u(\eta_i(\beta^N)).$$

For $i \in \{1, 2, \dots, n+m-1\}$ and any $N > 0$ we have $\eta_i(\beta^N) \in J_1$, thus $\eta_i(\beta^N) > \ell_1 > s_0$. So, using Lemma A.2, we can bound the first term above independently of N:

$$\left|\sum_{i=1}^{n+m-1} s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^\nu \cdot u(\eta_i(\beta^N))\right|$$
$$< \sum_{i=1}^{n+m-1}\left|\underbrace{(1\dots1)^\nu}_{i-1 \text{ times}}\cdot \sup_{x\in[\ell_1,1)} u(x)\right| < \infty.$$

The third term converges, since the sum converges absolutely:

$$\sum_{i=n+m+1}^{\infty}\left|s(i)\cdot\left(\eta_0(\beta^N)\dots\eta_{i-1}(\beta^N)\right)^\nu \cdot u(\eta_i(\beta^N))\right| = u(\beta^1)\cdot\sum_{i=n+1}^{\infty}\left((\beta^1)^\nu\right)^i < \infty$$

because $(\beta^1)^\nu < s_1 \le 1$ for $\nu > 0$. This bound on the third term does not depend on N. Thus, the first and third terms are bounded independently of N while the second term is $\Phi^1(\beta^N) \to \infty$ as $N \to \infty$, so $\Phi(\beta^N) \to \infty$ as needed. ∎

We are now ready to prove Lemma 4.1.

Proof of Lemma 4.1 Choose m as provided by Lemma A.8. Increase N by one at a time starting with $N = 1$. We know that $\Phi(\beta^1) = \Phi(\omega) < \Phi(\omega) + \varepsilon$ and, by Lemma A.10, there exists an M with $\Phi(\beta^M) > \Phi(\omega) + \varepsilon$. Let N be the smallest such M. Then, $\Phi(\beta^{N-1}) \le \Phi(\omega) + \varepsilon$, and by Lemma A.8,

$$\Phi(\beta^N) < \Phi(\beta^{N-1}) + \varepsilon \le \Phi(\omega) + 2\varepsilon.$$

Hence,

$$\Phi(\omega) + \varepsilon < \Phi(\beta^N) < \Phi(\omega) + 2\varepsilon.$$

Choosing $\beta = \beta^N$ completes the proof. ∎

Acknowledgements The authors would like to thank Stefano Marmi for sharing his insights into Brjuno-like functions.

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press

References

[BCM24] A. Bakhtawar, C. Carminati, and S. Marmi, *Global and local minima of α-Brjuno functions*. Monatshefte für Mathematik. **207**(2025), no. 2, 197–230.
[BM12] M. Balazard and B. Martin, *Comportement local moyen de la fonction de Brjuno*. Fundamenta Mathematicae. **218**(2012), 193–224.
[BY09] M. Braverman and M. Yampolsky, Computability of Julia sets, Algorithms and Computation in Mathematics, 23, Springer Berlin Heidelberg, 2009.
[Brj71] A. D. Brjuno, *Analytic form of differential equations, I, II*. Trudy Mosc. Mat. Ob. **25**(1971), 119–262.
[LMNN10] L. Luzzi, S. Marmi, H. Nakada, and R. Natsui, *Generalized Brjuno functions associated to α-continued fractions*. J. Approx. Theory **162**(2010), no. 1, 24–41.
[Mar22] S. Marmi, Small divisors, stability of orbits, and modular forms, University of Toronto, Toronto, ON, Colloquium talk, 2022.
[MMY97] S. Marmi, P. Moussa, and J.-C. Yoccoz, *The Brjuno functions and their regularity properties*. Commun. Math. Phys. **186**(1997), 265–293.
[MMY06] S. Marmi, P. Moussa, and J.-C. Yoccoz, *Some properties of real and complex Brjuno functions*. Front. Number Theory, Phys., Geom. **1**(2006), 603–625.
[Tur36] A. Turing, *On computable numbers, with an application to the Entscheidungsproblem*. Proc. London Math. Soc. **2**(1936), no. 42, 230–265.
[Wil33] J. R. Wilton, *An approximate functional equation with applications to a problem of Diophantine approximation*. J. Reine Angew. Math. **169**(1933), 219–237.
[Yoc96] J.-C. Yoccoz, *Théorem de Siegel nombres de Bruno et polynômes quadratiques*. Astérisque **231**(1996), 3–88.

University of Toronto, Canada
e-mail: yampol@math.toronto.edu ivan.shevchenko@mail.utoronto.ca

https://doi.org/10.4153/S0008414X2510179X Published online by Cambridge University Press