Ergod. Th. & Dynam. Sys. (2005), **25**, 227–256
DOI: 10.1017/S0143385704000173
© *2005 Cambridge University Press*
Printed in the United Kingdom

Dynamical properties of the Pascal adic transformation

XAVIER MÉLA† and KARL PETERSEN‡

† *IML, 163 avenue de Luminy, Case 907, 13288 Marseille Cedex 09, France*
(e-mail: mela@iml.univ-mrs.fr)
‡ *Department of Mathematics, CB 3250 Phillips Hall, University of North Carolina, Chapel Hill, NC 27599, USA*
(e-mail: petersen@math.unc.edu)

(*Received* 20 *May* 2003 *and accepted in revised form* 15 *October* 2003)

Abstract. We study the dynamics of a transformation that acts on infinite paths in the graph associated with Pascal's triangle. For each ergodic invariant measure the asymptotic law of the return time to cylinders is given by a step function. We construct a representation of the system by a subshift on a two-symbol alphabet and then prove that the complexity function of this subshift is asymptotic to a cubic, the frequencies of occurrence of blocks behave in a regular manner, and the subshift is topologically weak mixing.

1. *Introduction*

Adic transformations were introduced by A. Vershik as combinatorial models of the cutting and stacking constructions familiar in ergodic theory [**38, 39, 44**]. They move sequences in a transverse manner to the usual shift transformation, much as the horocycle flow is transverse to the geodesic flow (cf. [**17, 34**]). An adic transformation acts on the space of infinite paths on an infinite graded graph, or Bratteli diagram, and the dynamics of these transformations can provide information about the associated C^*-algebras or families of group representations (see, e.g., [**37, 41–43**]). Stationary adic transformations (in which after the first, or root, level all levels of the graph have the same number of vertices and the same pattern of connections to the next level) correspond to odometers and substitution subshifts [**7, 13, 16, 21, 35, 36, 44**]. Every minimal homeomorphism of the Cantor set is topologically conjugate to a particular type of adic transformation [**15**], and every ergodic measure-preserving transformation on a Lebesgue space is isomorphic to an adic transformation with a unique non-atomic invariant measure [**39, 40**]. The families of invariant sets for adic transformations correspond to tail fields in probability theory and ergodic theory, so ergodicity of invariant and quasi-invariant measures for adic systems is equivalent to 0,1 laws, which guarantee the triviality of these

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

tail fields—see [**27, 28, 32, 33**]. An especially regular and simple-looking non-stationary adic is the one based on the Pascal triangle regarded as a graded graph. Its σ-algebra of invariant sets corresponds to the exchangeable or symmetric σ-algebra in $\{0,1\}^{\mathbb{N}}$, the sets fixed by any permutation of finitely many coordinates (whereas the usual tail σ-algebra consists of the sets invariant under any change of finitely many coordinates).

In this paper we establish several dynamical properties of the Pascal adic transformation. It is known that the set of non-atomic ergodic invariant measures for this system is a one-parameter family corresponding to the Bernoulli measures on $\{0,1\}^{\mathbb{N}}$, as explained below. For each ergodic invariant measure we identify the asymptotic law of the return time to cylinder sets determined by finite initial path segments (Theorem 3.2). The original Pascal transformation is not defined everywhere, which means that we are dealing with a non-compact topological dynamical system. Attempts at compactification or at forming quotients lead to discontinuities. To overcome these difficulties, we use a countable family of substitutions to produce a subshift on an alphabet of two symbols, $\{a, b\}$, which represents the Pascal adic except for countably many points (Theorem 4.1). This subshift consists of all subwords of all 'basic words' formed by concatenating words, rather than adding integers, in Pascal's triangle. The basic word found at place k in row n has length equal to the binomial coefficient $C(n, k)$ found at the same place in the actual Pascal triangle, and the structure of the word conveys some information about the history of its formation and therefore also some extra information about the binomial coefficient. Not only does this subshift have zero entropy (it was known before that all the invariant measures for the Pascal adic transformation have entropy zero), but we can determine its complexity function: the number of n-blocks is asymptotic to $n^3/6$ (Theorem 4.6). While the subshift supports uncountably many ergodic invariant measures, it has a property that we call *directional unique ergodicity*: once a ray in Pascal's triangle beginning at the root is fixed, when we consider occurrences of a given block B only in basic blocks near that ray, the limiting frequency of occurrences exists and equals the measure of the cylinder set $[B]$ according to the ergodic invariant measure parameterized by the angle of the ray (Theorem 4.7). Finally, we use a characterization of weak mixing by Keynes and Robertson [**19**] and Weyl's theorem on uniform distribution to show that the subshift is topologically weakly mixing (Theorem 4.19).

2. *The Pascal adic transformation*

We define the Pascal adic transformation first in terms of its graph, then we give the cutting and stacking model to which it is isomorphic.

2.1. *The graph construction.*

The *Pascal graph* is an infinite planar graph divided into levels $n = 0, 1, \ldots$, with a root vertex at level 0 labeled $(0, 0)$, and $n + 1$ vertices at each level n labeled (n, k) for $k = 0, \ldots, n$. From each vertex (n, k) leave two edges; one goes to $(n+1, k+1)$ and is labeled by 1, and the other goes to $(n+1, k)$ and is labeled by 0—see Figure 1. The space X considered is the set of infinite paths going from the root down the graph, i.e. the set of all $(n, k_n)_{n\geq 1}$, where $0 \leq k_n \leq n$ and $k_{n+1} = k_n$ or $k_n + 1$. The labelling of the edges produces a natural one-to-one correspondence between X and

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 1. The Pascal graph. The number of finite paths from the root to a vertex (n, k) is given by the binomial coefficient $C(n, k) = n!/[k!(n-k)!]$.

the set $\{0, 1\}^{\mathbb{N}}$ of infinite sequences of 0's and 1's. The space X is compact for the product topology, and we denote by $\mathcal{B}$ the Borel σ-algebra. Let d be the usual metric on the space $\{0, 1\}^{\mathbb{N}}$ (letting $d(x, y) = (n+1)^{-1}$ whenever x and y disagree for the first time below the nth level). A cylinder set in X is a set of the type $\{x \in X : x_{i_1} = a_1, x_{i_2} = a_2, \ldots, x_{i_s} = a_s\}$, and the family of cylinder sets generates $\mathcal{B}$. For convenience we will often denote by $[a_1 a_2 \cdots a_s]$ the cylinder set $\{x \in X : x_1 = a_1, x_2 = a_2, \ldots, x_s = a_s\}$. We will refer to a point $x \in X$ by $(n, k_n(x))_{n \geq 1}$ or by $x_1 x_2 x_3 \cdots$, where $x_1, x_2, \ldots$ are the successive labels of the edges of x and $k_n(x) = \sum_{i=1}^{n} x_i$. We put a partial order on X, writing $x < y$ for $x, y \in X$, whenever x and y coincide below a certain level n and $x_n < y_n$. In other words, x and y are comparable with respect to this partial order if for some $n \in \mathbb{N}$ $n_j = y_j$ for all $j > n$ and there is a permutation $\pi \in \mathcal{S}_n$ such that $\pi(x_1, \ldots, x_n) = (y_1, \ldots, y_n)$. Let $X_{\min}$ and $X_{\max}$ be respectively the sets of minimal and maximal paths. We have

$$X_{\max} = \{x^i_{\max} = 0^i 1^\infty : i \geq 1\} \cup \{x^0_{\max} = 0^\infty, x^\infty_{\max} = 1^\infty\}$$
$$X_{\min} = \{x^i_{\min} = 1^i 0^\infty : i \geq 1\}.$$

Definition 2.1. The *Pascal adic transformation* is defined from $X \setminus X_{\max}$ onto $X \setminus X_{\min}$ by $Tx =$ smallest y greater than x. For every $x \in X \setminus X_{\max}$ there are positive integers n, m, and $x' \in \{0, 1\}^{\mathbb{N}}$ such that $x = 0^n 1^m 10 x'$. Hence an equivalent definition (illustrated in Figure 2) of T is

$$\boldsymbol{T(0^n 1^m 10 x') = 1^m 0^n 01 x'}.$$

Note that if $x = 1^k 0^{n-k} \cdots$, i.e. x coincides with the minimal path through the vertex (n, k), then $T^i x$ for $i = 1, \ldots, C(n, k) - 1$ goes through all $C(n, k)$ finite paths from $(0, 0)$ to (n, k).

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 2. T permutes finitely many coordinates and leaves the others unchanged; it 'carries' a path transversely to the shift transformation. Illustrated in the figure are $x = 0011\mathbf{10}0100\cdots$ and $Tx = 1100\mathbf{01}0100\cdots$.

There is a natural way to extend T bijectively on the whole space X by sending maximal paths to minimal ones:

$$\begin{aligned} Tx^{i}_{\max} &:= x^{i}_{\min} \quad \text{for all } i \geq 1, \\ Tx^{0}_{\max} &:= x^{0}_{\max}, \\ Tx^{\infty}_{\max} &:= x^{\infty}_{\max}. \end{aligned}$$

Unfortunately, this extension is not continuous at the points $x^{i}_{\max}$, $x^{\infty}_{\max}$.

A simple observation is that if a path x has a 'kink' at level n, i.e. if $x_{n+1}x_{n+2} = 10$—see Figure 3—then x comes back close to itself after $C(n, k_n(x))$ steps (see the following lemma).

LEMMA 2.2. (The 'kink' lemma) *If $x \in X$ is a path such that $x_{n+1}x_{n+2} = 10$, then $T^{C(n,k_n(x))}x$ and x coincide along the first n coordinates.*

Proof. Consider the following two paths (depicted in Figure 3) lying in the orbit of x:

$$\begin{aligned} x^{+} &= 0^{n-k_n(x)}1^{k_n(x)}10x_{n+3}x_{n+4}\cdots, \\ x^{-} &= 1^{k_n(x)}0^{n-k_n(x)}01x_{n+3}x_{n+4}\cdots. \end{aligned}$$

The adic takes the path x to x^{+} after l iterations, for a certain integer l. Then the adic applied once to x^{+} brings us to x^{-}. And, finally, it takes m iterations of the adic, for a certain integer m, to match up the first n coordinates of x. The total number of iterations we have made is simply the total number of finite paths from $(0, 0)$ to $(n, k_n(x))$, so that $l + 1 + m = C(n, k_n(x))$, establishing the lemma. □

Denote the orbit of a point $x \in X$ by $\mathcal{O}(x) = \{T^n x : n \in \mathbb{Z}\}$. (X, T) is not quite a minimal topological dynamical system (in the sense of a homeomorphism between

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 3. The 'kink' lemma.

compact spaces), but the kink lemma implies that if 10 appears infinitely many times in x, then x has a dense orbit. Assuming T has been extended to the whole space as above (sending maximal points to minimal ones), we have the following.

PROPOSITION 2.3. *T is a homeomorphism from $X_{\max}$ onto $X_{\min}$, and for every $x \in X$, exactly one of the following holds:*

(i) $\mathcal{O}(x) = \{x^0_{\max}\}$ *or* $\mathcal{O}(x) = \{x^\infty_{\max}\}$ *(x is a fixed point);*

(ii) *there exists $n \geq 1$ such that $x^n_{\max} \in \mathcal{O}(x)$ (the orbit of x is infinite but not dense);*

(iii) $\overline{\mathcal{O}(x)} = X$ *(x has a dense orbit).*

2.2. *Ergodic measures.* If $\tilde{p} = (p, 1-p)$ is the probability on $\{0, 1\}$ which gives mass p to 0 and mass $1-p$ to 1, then the product measure $\mu_p = \tilde{p}^{\otimes\mathbb{N}}$ on $\{0, 1\}^{\mathbb{N}}$ is called a *Bernoulli measure* and is often denoted by $\mathcal{B}(p, 1-p)$.

As noted in [**27, 28**] the result that the invariant ergodic Borel probability measures for the Pascal adic are the Bernoulli measures $\mathcal{B}(p, 1-p)$ is well known; it has been proved by using the ergodic theorem or the martingale convergence theorem. A more geometric and calculation-free approach developed in [**23**] permits extension of these results to a wider class of systems, the generalized Pascal adics. The statement that any T-invariant ergodic measure is a Bernoulli measure can be attributed to de Finetti in the context of exchangeable processes. The converse, stating that every Bernoulli measure is ergodic for the Pascal adic, follows from a result by Hajian *et al.* on a system isomorphic to the Pascal adic defined by interval splitting [**14**]. The connection with adics was made by Vershik [**39**].

In fact the Pascal adic is totally ergodic (every power T^n is ergodic). This is equivalent to saying that T does not have any eigenvalues (other than 1) which are roots of unity, which follows from the self-similar structure of Pascal's triangle modulo any prime (a consequence of a result of Lucas [**2, 22**]).

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 4. The binomial transformation, or cutting and stacking construction of the Pascal adic.

2.3. *The cutting and stacking equivalent.* Start by dividing the unit interval into two equal pieces. At each step, the stacks are divided into two equal halves, and the right half of each stack is placed on the bottom of the left half of the following stack—see Figure 4. If we repeat indefinitely, the resulting map T_{b} (which maps every open interval of each stack to the one above it) is defined everywhere except at the dyadic rationals (which correspond to the paths which are eventually diagonal in the graph construction). Denote by m the Lebesgue measure, and let $\mathcal{B}([0,1])$ be the σ-algebra of Borel sets in $[0,1]$. $([0,1], \mathcal{B}([0,1]), T_{\mathrm{b}}, m)$ is a measure-preserving system which we will refer to as the *binomial transformation*.

PROPOSITION 2.4. *The systems* $([0,1], \mathcal{B}([0,1]), T_b, m)$ *and* $(X, \mathcal{B}, T, \mu_{1/2})$ *are isomorphic.*

Proof. The isomorphism $\psi : [0,1] \setminus \{\text{dyadic rationals}\} \to X \setminus \{y : x \in X_{\min} \cup X_{\max}$ and $y \in \mathcal{O}(x)\}$ is defined by $\psi\big(\sum_{i=1}^{\infty} x_i 2^{-i}\big) = x_1 x_2 \cdots$. By induction one can check that at step n the bottom and top levels of the ith stack in the cutting and stacking are mapped respectively to the cylinders $[1^{n-i+1}0^{i-1}]$ and $[0^{i-1}1^{n-i+1}]$. Note also that two points with one exactly above the other have dyadic expansions which eventually coincide. If x is not a dyadic rational then for some level n it belongs to the right half of the top level in some stack, i.e. $\psi(x) = 0^{i-1}1^{n-i+1}0x_{n+1}x_{n+2}\cdots$ for some $i \in \{0, \ldots, n-1\}$. Therefore $\psi(T_{\mathrm{b}}(x)) = 1^{n-i}0^i 1 x_{n+1}x_{n+2}\cdots = T(\psi(x))$. Furthermore, ψ takes the Lebesgue measure m to the Bernoulli measure $\mu_{1/2}$, since the measures of the levels at step n are equal to 2^{-n}. □

Remark 1. The inverse of this isomorphism also carries any Bernoulli measure $\mathcal{B}(p, 1-p)$ on $\{0,1\}^{\mathbb{N}}$ to the Cantor measure m_p on $[0,1]$. Another point of view is to cut and stack with proportions p and $1-p$; then Lebesgue measure carries to the Bernoulli measure $\mathcal{B}(p, 1-p)$.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 5. The plot of the Pascal adic (or binomial transformation).

We used *Matlab* to produce the plot of the binomial transformation—see Figure 5. Note the symmetry with respect to the line $y = 1 - x$, which can be stated as

$$T_{\mathrm{b}}(1 - T_{\mathrm{b}}x) = 1 - x.$$

In the adic point of view this is equivalent to

$$TSTx = Sx,$$

where S is the transformation of $\{0, 1\}^{\mathbb{N}}$ which interchanges 0's and 1's. We state this observation as follows.

PROPOSITION 2.5. *The Pascal adic T is conjugate to its inverse via the map S which interchanges symbols: $TS = ST^{-1}$.*

Proof. Let $x \in X \setminus X_{\max}$. We can write x in the form $x = 0^i 1^j \mathbf{10} x'$, where $i, j \geq 0$ and x' is an infinite string of 0's and 1's. Then

$$\begin{aligned} TST(x) &= TS(1^j 0^i \mathbf{01} x') \\ &= T(0^j 1^i \mathbf{10} S(x')) \\ &= 1^i 0^j \mathbf{01} S(x') = S(x). \end{aligned}$$

□

3. *Limit laws for return times into cylinders*

Considerable attention has been devoted recently to determining the asymptotic laws of return times or hitting times to 'shrinking targets': see, e.g., [**1, 4, 5, 8, 20, 25, 29, 31**] and the references that they contain. We establish for the Pascal adic that the limit laws of return times into typical cylinders, when properly scaled, are piecewise constant.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 6. The first time an element of $C^n_{l,m}$ returns to U_n is when it enters the cylinder $[\omega_1 \cdots \omega_n 1^m 0^l 01]$.

For simplicity we assume that μ is the Bernoulli measure $\mathcal{B}(1/2, 1/2)$, but the steps below can be adapted to the general case. Fix a generic point $\omega \in X$. Denote by U_n the cylinder generated by the coordinates $\omega_1, \omega_2, \ldots, \omega_n$ (as above we use the notation $U_n = [\omega_1 \omega_2 \cdots \omega_n]$). Let τ_n be the first return time (or entrance time) to U_n, i.e.

$$\tau_n(x) = \inf\{k \geq 1 : T^k x \in U_n\}.$$

We are interested in the asymptotics of the return times τ_n, the question being what is the limit of the following function when the right scaling c_n is chosen:

$$G_n(t) = \frac{1}{\mu(U_n)} \mu\{x \in U_n : c_n \mu(U_n) \tau_{U_n}(x) > t\}.$$

Consider the cylinders $C^n_{l,m} = [\omega_1 \cdots \omega_n 0^l 1^m 10]$. Then $U_n = \bigcup_{l,m \geq 0} C^n_{l,m}$ (disjoint union up to a set of μ-measure zero). For $x \in C^n_{l,m}$ and $k_n = \sum_{j=1}^n \omega_j$—see Figure 6—the first return time to U_n is given by the following formula.

PROPOSITION 3.1. $\tau_n(x) = C(n + l, k_n) + C(n + m, k_n + m) - C(n, k_n)$.

Proof. If $x \in C^n_{l,m}$, it will return to U_n for the first time when it enters the cylinder $[\omega_1 \ldots \omega_n 1^m 0^l 01]$. After a certain number N_1 of iterations x gets mapped to $[0^{n-k_n} 1^{k_n} 0^l 1^m 10]$. It takes $C(n+l, k_n) - C(n, k_n)$ more iterations to bring x to the cylinder $[0^{n-k_n+l} 1^{k_n+m} 10]$. Applying the adic one more time takes us to $[1^{k_n+m} 0^{n-k_n+l} 01]$. Similarly, after $C(n + m, k_n + m) - C(n, k_n)$ more iterations we are in $[1^{k_n} 0^{n-k_n} 1^m 0^l 01]$ and after a certain number N_2 of iterations x finally re-enters U_n. To conclude, note that $N_1 + N_2 + 1 = C(n, k_n)$. □

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

Letting $t^n_{l,m} = (1/2^n)c_n\tau_n$ for $n \geq 1, l, m \geq 0$, we have

$$G_n(t) = 2^n \sum_{l,m\geq 0} \mu\{x \in C^n_{l,m} : t^n_{l,m} > t\} = 2^n \sum_{l,m\geq 0:t^n_{l,m}>t} \mu(C^n_{l,m})$$
$$= 2^n \sum_{l,m\geq 0:t^n_{l,m}>t} \frac{1}{2^{n+l+m+2}} = \frac{1}{4} \sum_{l,m\geq 0:t^n_{l,m}>t} \frac{1}{2^{l+m}}.$$

Using Stirling's formula, we know that $C(n,k_n)/2^n \approx 1/\sqrt{2\pi(n-k_n)}$, so that when c_n grows faster than $\sqrt{n}$, $\lim_{n\to\infty} t^n_{l,m} = 0$ and therefore $\lim_{n\to\infty} G_n(t) = \mathbb{1}_{(-\infty,0]}(t)$; and when c_n grows slower than $\sqrt{n}$, then $\lim_{n\to\infty} t^n_{l,m} = \infty$, which implies that $\lim_{n\to\infty} G_n(t) = \mathbb{1}_{(-\infty,\infty)}$. The interesting scaling is $c_n = \sqrt{n}$. Then, using again Stirling's formula and the fact that $k_n/n \to 1/2\ \mu$ almost everywhere, we get

$$t^n_{0,0} = \frac{1}{2^n}\sqrt{n}C(n,k_n) = \frac{1}{2^n}\sqrt{n}\frac{n!}{k_n!(n-k_n)!}$$
$$\approx \frac{1}{2^n}\sqrt{n}\frac{(n^n/e^n)\sqrt{2\pi n}}{\frac{k_n^{k_n}}{e^{k_n}}\sqrt{2\pi k_n}[(n-k_n)^{n-k_n}/e^{n-k_n}]\sqrt{2\pi(n-k_n)}}$$
$$\approx \frac{1}{2^n\sqrt{2\pi}}\sqrt{n}\frac{n^{n+1/2}}{k_n^{k_n+1/2}(n-k_n)^{n-k_n+1/2}}$$
$$\approx \frac{1}{2^n\sqrt{2\pi}}\sqrt{\frac{n}{n-k_n}}\left(\frac{n}{k_n}\right)^{k_n+1/2}\left(\frac{n}{n-k_n}\right)^{n-k_n}$$
$$\underset{n\to\infty}{\longrightarrow} \sqrt{\frac{2}{\pi}}.$$

Then

$$t^n_{l,m} = \frac{1}{2^n}\sqrt{n}[C(n+m,k_n+m) + C(n+l,k_n) - C(n,k_n)]$$
$$= t^n_{0,0}\left[\frac{(n+m)\cdots(n+1)}{(k_n+m)\cdots(k+1)} + \frac{(n+l)\cdots(n+1)}{(n+l-k_n)\cdots(n+1-k_n)} - 1\right]$$
$$\underset{n\to\infty}{\longrightarrow} \sqrt{\frac{2}{\pi}}(2^l + 2^m - 1).$$

We can now easily deduce that G_n converges to a step function whose heights are computed below. Let $t_{i,j} = \sqrt{2/\pi}(2^i + 2^j - 1)$ for $i \geq 0$ and $0 \leq j \leq i$.

Assume that $t_{i,j} \leq t < t_{i,j+1}$ and $j+1 \leq i$; then

$$\lim_{n\to\infty} G_n(t) = \frac{1}{4}\lim_{n\to\infty} \sum_{l,m\geq 0:t^n_{l,m}>t} \frac{1}{2^{l+m}}$$
$$= \frac{1}{4}\sum_{l,m\geq 0:t_{l,m}>t} \frac{1}{2^{l+m}}$$
$$= \frac{1}{4}\left(\sum_{l=m\geq 0:t_{l,m}>t} \frac{1}{2^{l+m}} + 2\sum_{l>m\geq 0:t_{l,m}>t} \frac{1}{2^{l+m}}\right)$$

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

$$= \frac{1}{4}\Big(\sum_{l=i}^{\infty} \frac{1}{2^{2l}} + 2 \sum_{m=j+1}^{i-1} \frac{1}{2^{i+m}} + 2 \sum_{l=i+1}^{\infty} \sum_{m=0}^{l-1} \frac{1}{2^{l+m}} \Big)$$
$$= 2^{-1-2i}(2^{i+1} + 2^{i-j} - 2).$$

Similarly, if $t_{i,i} \le t < t_{i+1,0}$ (the case $i = j$), then

$$\lim_{n\to\infty} G_n(t) = \frac{1}{4}\Big(\sum_{l=i+1}^{\infty} \frac{1}{2^{2l}} + 2 \sum_{l=i+1}^{\infty} \sum_{m=0}^{l-1} \frac{1}{2^{l+m}} \Big)$$
$$= 2^{-2(i+1)}(2^{i+2} - 1).$$

To summarize, we have the following.

THEOREM 3.2. *$G_n(t) = (1/\mu(U_n))\mu\{x \in U_n : \sqrt{n}\mu(U_n)\tau_n(x) > t\}$ converges pointwise as $n \to \infty$ to the piecewise constant function*

$$\begin{cases} 1 & \text{if } t < 0, \\ 2^{-1-2i}(2^{i+1} + 2^{i-j} - 2) & \text{if } t_{i,j} \le t < t_{i,j+1} \text{ and } i > j+1, \\ 2^{-2(i+1)}(2^{i+2} - 1) & \text{if } t_{i,i} \le t < t_{i+1,0}, \end{cases}$$

where $t_{i,j} = \sqrt{2/\pi}(2^i + 2^j - 1)$ for $i \ge 0$ and $0 \le j \le i$.

Remark 2. If $p \ne 1/2$ and μ is the Bernoulli measure $\mathcal{B}(p, 1-p)$, then $G_n(t)$ defined as above still converges to a piecewise constant function, but the formula for the function is rather messy.

4. *A countable-substitution subshift*

Stationary adics are topologically conjugate to substitution or odometer symbolic dynamical systems [**7, 13, 21**]. Here we use the idea described in [**7, 16**] to associate a substitution to an adic to show that the Pascal adic is isomorphic to a subshift whose language is determined by countably many substitutions. For basic definitions and general information about substitution systems see [**12, 30**]. We will use the alphabet $\{a, b\}$ for our subshift to decrease confusion about the various representations of the Pascal adic.

Consider the substitutions $\zeta_i : \{0, 1, \ldots, i\} \to \{a, b\}^* =$ all finite words on the alphabet $\{a, b\}$ defined by

$$\zeta_i(i) = b, \quad \zeta_i(0) = a \quad \text{for all } i \ge 1,$$
$$\zeta_i(j) = \zeta_{i-1}(j)\zeta_{i-1}(j-1) \quad \text{for all } i \ge 2 \text{ and } 0 < j < i.$$

Denote by $\mathcal{L}(\zeta_i)$ the language associated to ζ_i, by which we mean the subset of $\{a, b\}^*$ consisting of all subwords of all the $\zeta_i(j), i \ge 1, 0 \le j \le i$. Let Σ be the subshift consisting of all sequences in $\{a, b\}^{\mathbb{Z}}$ all of whose subwords belong to $\mathcal{L}(\zeta) = \bigcup_i \mathcal{L}(\zeta_i)$. Figure 7 shows how the 'basic' words in the language $\mathcal{L}(\zeta)$ can be obtained by successive concatenations. It is not very hard to see that these basic words give the codings of the cylinders $[1^k 0^{n-k}]$ in the Pascal graph under the action of the Pascal adic transformation according to the first edge (left $\sim$ label 1 $\sim b$, right $\sim$ label 0 $\sim a$)—or, in the cutting and stacking representation, the coding according to visits to $[0, 1/2) \sim b$ or $[1/2, 1] \sim a$

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

$$
\begin{array}{ccccccccccc}
 & & & & b & & a & & & & \\
 & & & & b & & ba & & a & & \\
 & & & & b & & b^2a & & ba^2 & & a \\
 & & & & b & & b^3a & & b^2aba^2 & & ba^3 & & a \\
 & & & & b & & b^4a & & b^3ab^2aba^2 & & b^2aba^2ba^3 & & ba^4 & & a \\
 & & b & & b^5a & & b^4ab^3ab^2aba^2 & & b^3ab^2aba^2b^2aba^2ba^3 & & b^2aba^2ba^3ba^4 & & ba^5 & & a \\
\cdots & & & & \cdots & & & & \cdots & & & & \cdots & & & & \cdots
\end{array}
$$

FIGURE 7. 'The Pascal triangle of words', showing the construction of the words by successive concatenations from level 1 to level 6. The word written at the 'vertex' (n, k) of this triangle represents the coding of the cylinder $[1^k 0^{n-k}]$.




FIGURE 8. Coding by a's and b's in the cutting and stacking.

(see Figure 8). (Just note that the cylinder $[1^k 0^{n-k}]$ corresponds to the bottom level of the kth stack in the cutting and stacking at stage n, thus its coding is the word seen going up the stack (Figure 8), which is the word $\zeta_n(k)$ shown at vertex (n, k) in Figure 7.)

THEOREM 4.1. *Let (X, T) denote the Pascal adic. There are a countable subset $X' \subset X$ and a one-to-one Borel measurable map $\phi : X \setminus X' \to \Sigma$ such that $\phi \circ T = \sigma \circ \phi$ on $X \setminus X'$. Consequently, for all $\alpha \in (0, 1)$, for each non-atomic ergodic measure μ_α on (X, T), if $\nu_\alpha = \mu_\alpha \circ \phi^{-1}$, then (X, T, μ_α) and $(\Sigma, \sigma, \nu_\alpha)$ are measurably isomorphic.*

Proof. Let $P = \{[0], [1]\}$, where we denote as previously $[0] = \{x \in X : x_0 = 0\}$ and $[1] = \{x \in X : x_0 = 1\}$. Let X' be the countable subset of X consisting of all paths which are eventually diagonal, i.e. $x \in X'$ if and only if there exists $n \in \mathbb{N}$ such that either $x_k = 0$ for all $k \geq n$, or $x_k = 1$ for all $k \geq n$. Let ϕ be the coding of the Pascal adic (X, T) by the partition P, where $[0]$ is coded with a and $[1]$ with b. More precisely, for every $x \in X \setminus X'$ we define

$$\phi(x) = (\ldots, \omega_{-2}, \omega_{-1}, \omega_0, \omega_1, \omega_2, \ldots),$$

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 9. After N steps x and y are mapped to the 'maximal' cylinder $[0^l 1^m]$. At step $N+1$, x is mapped to $1^{m+j-1}0^l$ whereas y is mapped to $1^m 0^l$. This shows that x will then stay in the cylinder [1] coded by b for $m+j-1$ steps until it is mapped to the cylinder [0] coded by a, whereas it will take only m steps for y to get mapped into [0].

where

$$\omega_i = \begin{cases} a & \text{if } (T^i x)_0 = 0, \\ b & \text{if } (T^i x)_0 = 1. \end{cases}$$

The map ϕ intertwines T and σ (this is clear) and establishes a measure-theoretic isomorphism between (X, T, μ_α) and $(\Sigma, \sigma, \nu_\alpha)$. To check this, it is enough to show that ϕ is one-to-one, or, equivalently, that P is a generating partition for T. Let $x, y \in X$ be two different paths which are not eventually diagonal. Since $x \neq y$ there is a smallest integer n such that $x_n \neq y_n$, and we can assume that $x_n = 1$ and $y_n = 0$. In addition, since $x \notin X_{\max}$, $x_{n+j} = 0$ for some smallest $j \geq 1$. Consequently, the P-names of x and y coincide until x and y get mapped after N iterations to the cylinder $[0^l 1^m]$ for some $N, l, m \geq 1$ (where $m = k_{n-1}(x) = k_{n-1}(y)$, $l = n - k_{n-1}(x)$, and $0 \leq N \leq C(n, k_{n-1}(x))$). Then, since $T^N x = 0^l 1^{m+j+1} 0 \cdots$ and $T^N y = 0^l 1^m 0 \cdots$, it follows that the P-name of $T^N x$ is $b^{m+j+1} a \cdots$ whereas the P-name of $T^N y$ is $b^m a \ldots$, showing that $\phi(x) \neq \phi(y)$—see Figure 9. □

4.1. *Complexity.* It is easy to see that the subshift (Σ, σ) has topological entropy 0. Finer measures of the size or richness of symbolic dynamical systems can be drawn from asymptotics of the complexity function, which for each n gives the number p_n of n-blocks found in sequences in the system. See [**3, 9, 11, 12**] for some examples of results on the complexity functions of various systems. Usually one finds upper or lower estimates on

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

the growth rate of a complexity function. Here we show that for the subshift (Σ, σ) that we have associated to the Pascal adic, p_n is asymptotic to $n^3/6$.

Let $B_{n,j} = \zeta_j(n)$ denote the block at the vertex (n, j) in Figure 7 (for example, $B_{3,2} = b^2a$). We continue to refer to the $B_{n,j}$'s as *basic blocks*. These blocks satisfy the recurrence formula

$$B_{n,j} = B_{n-1,j}B_{n-1,j-1}, \text{ for } 0 \le j \le n. \tag{4.1}$$

Every basic block at level n can be written as the product (we use interchangeably 'product' and 'concatenation') of two basic blocks from level $n-1$, and every basic block at level $n-1$ can be written as the product of two basic blocks from level $n-2$, and so forth. Thus we have a *hierarchical decomposition* of the basic blocks. We can view any block $B_{n,j}$ as the product of basic blocks from level l for any $l \le n$, depending on how far back in the hierarchy we want to look.

Definition 4.2. We say $B \in \mathcal{L}(\zeta)$ is a *new block at level* l if B appears as a subblock of one of the basic blocks $B_{l,k}$ at level l, but does not appear as a subblock at level $l-1$ (and *a fortiori* at any lower levels).

We would like to count the number of different new n-blocks at level l for $l = 1, 2, \ldots$. Although the recursive construction of the blocks is simple, it is not clear how to count precisely the different n-blocks since a given n-block can appear many times at the same level. The concatenation of two consecutive basic blocks at level l will result in the formation of new blocks at level $l+1$, but how can we tell whether a block which overlaps two such basic blocks did not appear higher in the Pascal triangle of words (see Figure 7)? After which level will we have seen all the different n-blocks? The following lemma provides an answer to these questions.

LEMMA 4.3. *Let B be an n-subblock at level $l+1$ of $B_{l+1,j}$, for $2 \le j \le l-1$. Assume that l, j are such that $|B_{l-1,j-1}| \ge \max\{n-(l-j), n-(j-1)\}$ (which guarantees that B does not overlap $B_{l-1,j}$ or $B_{l-1,j-2}$). Then B is a new n-block at level $l+1$ if and only if B contains $a^{l-j}b^{j-1}$.*

Example 4.4. Observe in Figure 7 that the only new 5-blocks which appear at level 6 as subblocks of $B_{6,4}$, $B_{6,3}$, $B_{6,2}$ are

$$b\boldsymbol{ab^3}, \boldsymbol{ab^3}a, b\boldsymbol{a^2b^2}, \boldsymbol{a^2b^2}a, b\boldsymbol{a^3b}, \boldsymbol{a^3b}a.$$

They all contain the block $\boldsymbol{a^{5-j}b^{j-1}}$ for some j.

Proof. Suppose that B is a new block at level $l+1$ not containing the subblock $a^{l-j}b^{j-1}$. For example, in the case $B_{l,j} = B_{6,4}$ and $B_{l,j-1} = B_{6,3}$, we have the following picture:

$$B_{l,j} = b^4a|\overbrace{b^3ab^2ab\boldsymbol{a^2}}^{B_{l-1,j-1}} \quad \overbrace{\boldsymbol{b^3}ab^2aba^2}^{B_{l-1,j-1}}|b^2aba^2ba^3 = B_{l,j-1},$$
$$B_{l,j} = b^4a|b^3\underbrace{ab^2ab\boldsymbol{aa} \quad \boldsymbol{bbb}\,\boldsymbol{a}}_{B}\underbrace{ab^2aba^2|b^2}_{B}aba^2ba^3 = B_{l,j-1}.$$

(Note: We use '|' to symbolize where the concatenation at the previous level took place.)

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

Observe that a^{l-j} is a right factor of $B_{l,j}$, and that b^{j-1} is a left factor of $B_{l,j-1}$. Consequently, there exists k such that either $B = \omega b^k$ and $1 \le k \le j-2$, or $B = a^k\omega$ and $1 \le k \le l-j$, where ω is, respectively, a right or left factor of $B_{l-1,j-1}$. The latter follows from the hypothesis that $|B_{l-1,j-1}| \ge \max\{n-(l-j), n-(j-1)\} \ge |B|-k$. Suppose that $B = \omega b^k$ (the case $B = a^k\omega$ is similar). Since $B_{l,j-1} = B_{l-1,j-1}B_{l-1,j-2} = B_{l-1,j-1}b^{j-2}\cdots$, it follows that B is a subblock of $B_{l,j-1}$. This contradicts the fact that B is a new block at level $l+1$.

Conversely, observe that for $1 \le j \le l-1$, $B_{l,j} = b^j a \ldots b a^{l-j}$. Thus, the first time the block $a^{l-j}b^{j-1}$ will be seen is at level $l+1$, as a subblock of $B_{l,j}B_{l,j-1}$. □

As a corollary we get the following.

LEMMA 4.5. *All n-blocks are seen at level $n+2$ as subblocks.*

Proof. Assume there exists a new n-block B at level $n+3$. First, note that B cannot be a subblock of any of the following 'edge' blocks:

$$B_{n+3,n+3} = b, \quad B_{n+3,n+2} = b^{n+2}a, \quad B_{n+3,1} = ba^{n+2}, \quad B_{n+3,0} = a.$$

Otherwise, $B = b^n, b^{n-1}a, a^n$ or ba^{n-1}, and those blocks are already seen at level $n+1$. Therefore, B is coming from the concatenation of $B_{n+2,j}$ and $B_{n+2,j-1}$ for some j with $2 \le j \le n+1$. By Lemma 4.3 (applied in the case $l = n+2$), B must contain the subblock $a^{n+2-j}b^{j-1}$, which is impossible. □

THEOREM 4.6. $\lim_{n\to\infty}(p_n/n^3) = 1/6$.

Proof. Fix n. Consider the Pascal triangle of words from level 1 to level $n+1$. We will count the number of n-blocks by estimating how many new n-blocks are created from one level to the next by concatenation of two adjacent basic blocks. We divide the triangle into two disjoint regions: let *region I* be the subset of the triangle formed by all blocks up to level $\sqrt{2n}$, as well as all blocks below the level $\sqrt{2n}$ located at the vertices (l,k), for $\sqrt{2n} \le l \le n+1$ and $k = 0, 1, l-1, l$. Let *region II* be the complement of region I in the triangle considered—see Figure 10. Let p_{I} be the sum for $l = 1$ to $n+1$ of the number of new n-blocks at level l appearing as subblocks of the product of two adjacent basic blocks, one of them at least belonging to region I. Similarly, let p_{II} denote the sum for $\sqrt{2n} \le l \le n+1$ of the number of new n-blocks at level l appearing as subblocks of the product of two consecutive basic blocks, both of them belonging to region II. By Lemma 4.5, since all n-blocks are seen at level $n+2$, we have the following inequality:

$$p_{\mathrm{II}} \le p_n \le p_{\mathrm{I}} + p_{\mathrm{II}}.$$

In region II, note that the hypothesis of Lemma 4.3 is satisfied since if $B_{l,j}$ and $B_{l,j-1}$ are two consecutive blocks in that region $|B_{l-1,j-1}| \ge (\sqrt{2n}+1)\sqrt{2n}/2$. By Lemma 4.3, a subblock B of $B_{l,j}B_{l,j-1}$ is a new n-block at level $l+1$ if and only if $a^{l-j}b^{j-1}$ is a subblock of B. Therefore, the number of new n-subblocks of $B_{l,j}B_{l,j-1}$ is equal to $n-l$.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 10. In region II, two consecutive basic blocks $B_{l,j}$ and $B_{l,j-1}$ at level l will generate $n-l$ new n-blocks at level $l+1$, according to Lemma 4.3.

Thus,

$$\begin{aligned} p_{\mathrm{II}} &= \sum_{\sqrt{2n}<l<n+1} (n-l)(l-3) \\ &= n\sum_{l=1}^{n+1} l - \sum_{l=1}^{n+1} l^2 + o(n^3) \\ &= n\frac{(n+1)(n+2)}{2} - \frac{(n+1)(n+2)(2n+3)}{6} + o(n^3) \\ &= \frac{n^3}{2} - \frac{2n^3}{6} + o(n^3) = \frac{n^3}{6} + o(n^3). \end{aligned}$$

For region I, a coarse approximation gives

$$p_{\mathrm{I}} \le \sum_{1\le l\le\sqrt{2n}} (n-1)(l-1) + 4 \sum_{\sqrt{2n}<l\le n+1} (l-1) = o(n^3).$$

It follows that

$$\frac{n^3}{6} + o(n^3) \le p_n \le \frac{n^3}{6} + o(n^3). \qquad \square$$

An immediate corollary is that the countable-substitution subshift (Σ, σ) has topological entropy

$$h_{\mathrm{top}}(\Sigma, \sigma) = \lim_{n\to\infty} \frac{\log(p_n)}{n} = 0.$$

The variational principle implies that the measure-theoretic entropy $h(\sigma)$ with respect to every measure ν_α is zero. Since entropy is an isomorphism invariant, this proves that the Pascal adic has zero entropy for every Bernoulli measure μ_α.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

4.2. *Directional unique ergodicity.* For any two blocks B and C in the language $\mathcal{L}(\zeta)$, let $\mathrm{freq}(B, C)$ denote the frequency of occurrences of B in C. In particular, if $\omega \in \Sigma$, and if ω_{-N}^{N} denotes the block $\omega_{-N}\omega_{-N+1}\cdots\omega_0\cdots\omega_{N-1}\omega_N$, then, provided that $2N+1$ is greater than $|B|$ (the length of B), we have

$$\mathrm{freq}(B, \omega_{-N}^{N}) = \frac{1}{2N+1}\sum_{i=-N}^{N-|B|} \mathbb{1}_{[B]} \circ \sigma^i(\omega).$$

The ergodicity of each $\nu_\alpha = \phi(\mu_\alpha)$ (carried by the isomorphism) implies therefore that for every block $B \in \mathcal{L}(\zeta)$ and ν_α-a.e. $\omega \in \Sigma$

$$\lim_{N\to\infty} \mathrm{freq}(B, \omega_{-N}^{N}) = \nu_\alpha(B). \tag{4.2}$$

Let x be a path in the Pascal graph going through the vertices $(n, k_n(x))$, and let $\omega = \phi(x)$. If x is not eventually diagonal, then there are sequences $i_n(x) \nearrow \infty$ and $j_n(x) \nearrow \infty$ such that $B_{n,k_n(x)} = \omega_{-i_n(x)}^{j_n(x)}$, where B_{n,k_n} is the basic block at vertex (n, k_n) in the Pascal triangle of words. Therefore, (4.2) implies that for μ_α almost every x

$$\lim_{n\to\infty} \mathrm{freq}(B, B_{n,k_n(x)}) = \nu_\alpha(B).$$

In other words, when (n, k_n) for $n = 1, 2, \ldots$ are the vertices of a generic path, which goes down the Pascal graph at an 'angle' α, we have $\mathrm{freq}(B, B_{n,k_n}) \to \nu_\alpha(B)$. We strengthen this statement as follows.

THEOREM 4.7. *For any block $B \in \mathcal{L}(\zeta)$, any $\alpha \in (0, 1)$, and* any *sequence $k_n \to \infty$ such that $k_n/n \to \alpha$ we have*

$$\lim_{n\to\infty} \mathrm{freq}(B, B_{n,k_n}) = \nu_\alpha(B).$$

To prove Theorem 4.7 we determine explicitly where a given block B is made and how many times it appears in each basic block. First, we introduce some notation and recall the key structure of the basic blocks. For a fixed block $B \in \mathcal{L}(\zeta)$, denote by $a(B, n, k)$ the number of occurrences of B in $B_{n,k}$. If no confusion is possible we will simply denote it by $a(n, k)$, so that

$$\mathrm{freq}(B, B_{n,k_n}) = \frac{a(n, k_n)}{|B_{n,k_n}|} = \frac{a(n, k_n)}{C(n, k_n)}. \tag{4.3}$$

By induction it is easy to show that

$$B_{n,k} = \underbrace{\boldsymbol{b^k}a\cdots b\boldsymbol{a^{n-k-1}}}_{B_{n-1,k}}\underbrace{\boldsymbol{b^{k-1}}a\cdots b\boldsymbol{a^{n-k}}}_{B_{n-1,k-1}}.$$

This decomposition characterizes the basic blocks $B_{n,k}$; if one sees the *telltale* block $b\boldsymbol{a^{n-k-1}b^{k-1}}a$, it is always within $B_{n,k}$, at the 'join', as seen above. The previous structure is easily seen by writing only the beginning and ending of the basic blocks in the Pascal triangle of words; for example, at level n we have:

$$\underbrace{\boldsymbol{b}}_{B_{n,n}} \quad \underbrace{\boldsymbol{b^{n-1}a}}_{B_{n,n-1}} \quad \cdots \quad \underbrace{\boldsymbol{b^k}a\cdots b\boldsymbol{a^{n-k}}}_{B_{n,k}} \quad \cdots \quad \underbrace{\boldsymbol{ba^{n-1}}}_{B_{n,1}} \quad \underbrace{\boldsymbol{a}}_{B_{n,0}}$$

Figure 12 shows how the structure is carried from one level to the next.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 11. For any block $B_{n,k}$ in the filled area the frequency of appearances of B in $B_{n,k}$ is ϵ-close to $\nu_\alpha(B)$ (for δ small and N large).

$$\overbrace{\boldsymbol{b^{k-1}}a\cdots b\boldsymbol{a^{n-k-1}}}^{B_{n-2,k-1}}$$

$$\underbrace{\boldsymbol{b^{k}}a\cdots b\boldsymbol{a^{n-k-1}}}_{B_{n-1,k}} \quad \underbrace{\boldsymbol{b^{k-1}}a\cdots b\boldsymbol{a^{n-k}}}_{B_{n-1,k-1}}$$

$$\boldsymbol{b^{k+1}}a\cdots b\boldsymbol{a^{n-k-1}} \quad \underbrace{\boldsymbol{b^{k}}a\cdots b\boldsymbol{a^{n-k}}}_{B_{n,k}} \quad \boldsymbol{b^{k-1}}a\cdots b\boldsymbol{a^{n-k+1}}$$

$$\boldsymbol{b^{k+2}}a\cdots b\boldsymbol{a^{n-k-1}} \quad \boldsymbol{b^{k+1}}a\cdots b\boldsymbol{a^{n-k}} \quad \boldsymbol{b^{k}}a\cdots b\boldsymbol{a^{n-k+1}} \quad \boldsymbol{b^{k-1}}a\cdots b\boldsymbol{a^{n-k+2}}$$

FIGURE 12.

LEMMA 4.8. *Let $B \in \mathcal{L}(\zeta)$. Then there is a unique vertex (n_0, k_0) such that B is a subblock of B_{n_0,k_0} and B does not appear in any other basic block in the 'rectangle' above (n_0, k_0) formed by all vertices (n, k) with $k < k_0$ and $n - k < n_0 - k_0$.*

Proof. Assume that B is a subblock of both B_{n_0,k_0} and $B_{n_0',k_0'}$, and that B does not appear in any other basic block in the 'rectangles' above (n_0, k_0) and (n_0', k_0') (in particular we can assume that $n_0' > n_0$ and $k_0' > k_0$)—see Figure 13. Recall that

$$B_{n_0,k_0} = b^{k_0}a\cdots ba^{n_0-k_0-1}|b^{k_0-1}\cdots ba^{n_0-k_0},$$
$$B_{n_0',k_0'} = b^{k_0'}a\cdots ba^{n_0'-k_0'-1}|b^{j_0'-1}\cdots ba^{n_0'-k_0'},$$

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 13.

so that B must contain a central subblock of both

$$b\,\overbrace{a\cdots a}^{n_0'-k_0'-1}\,|\,\overbrace{b\cdots\cdots\cdots b}^{k_0'-1}\,a$$
$$b\,\underbrace{a\cdots\cdots\cdots a}_{n_0-k_0-1}|\,\underbrace{b\ldots b}_{k_0-1}\,a.$$

(By 'central' we mean containing $a|b$, where '|' symbolizes the splitting into blocks of the above level.) The only way it can happen is if $B = a^i b^j$ with $i < n_0' - k_0' - 1$ and $j < k_0 - 1$. Therefore B had to appear at the vertex $(n_0' - k_0' + k_0, k_0)$ whose basic block is $\cdots b a^{n_0'-k_0'-1} b^{k_0-1} a \cdots$, contradicting our initial assumption. □

Remark 3. From the uniqueness in Lemma 4.8, it follows that if the block B appears for the first time at the vertex (n_0, k_0), then it will only appear as a subblock in the triangle below the vertex (n_0, k_0), as shown in Figure 14.

LEMMA 4.9. *If a block B appears in B_{n_0,k_0} for the first time, then it can appear at most twice (i.e. $a(n_0, k_0) = 1$ or 2).*

Proof. If B appears for the first time in B_{n_0,k_0}, then it is a central subblock of

$$b^{k_0}\cdots\cdots\underbrace{a^{n_0-k_0-1}|b^{k_0-1}\cdots\cdots}_{B}a^{n_0-k_0}.$$

There are two possibilities: either B contains the telltale block $ba^{n_0-k_0-1}|b^{k_0-1}a$, or it does not. In the first case, B cannot appear twice, since the telltale block appears only once in B_{n_0,k_0}. In the second case, the only way that B could appear twice is if it would start

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 14.

with $a^i b^{k_0-1}$ for some $i \leq n_0 - k_0 - 1$, and end with $a^{n_0-k_0-1}b^j$ for some $j \leq k_0 - 1$:

$$b^{k_0} \cdots\cdots \overbrace{a^{n_0-k_0-1}|b^{k_0-1}}^{B} \cdots\cdots a^{n_0-k_0},$$
$$b^{k_0} \cdots\cdots \underbrace{a^{n_0-k_0-1}|b^{k_0-1}}_{B} \cdots\cdots a^{n_0-k_0}.$$

Then clearly B cannot appear a third time. □

Given a block $B \in \mathcal{L}(\zeta)$, assume that B appears for the first time at vertex (n_0, k_0) in the Pascal triangle of words (this first appearance is unique by Lemma 4.8). Define the *triangle of appearances of* B to be

$$\begin{array}{ccccccccccc} & & & & & a(n_0, k_0) & & & & & \\ & & & & a(n_0+1, k_0+1) & & a(n_0+1, k_0) & & & & \\ & & & a(n_0+2, k_0+2) & & a(n_0+2, k_0+1) & & a(n_0+2, k_0) & & & \\ \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots & \cdots \end{array}$$

By Lemma 4.8, all other frequencies are zero, namely $a(n, k) = 0$ for all vertices (n, k) not equal to $(n_0 + i, k_0 + j)$ where $i \geq 0$ and $0 \leq j \leq i$—see Figure 14.

Example 4.10. The triangle of appearances of $B = a^2b^2$ is

$$\begin{array}{ccccc} & & 1 & & \\ & 2 & & 2 & \\ 3 & & 5 & & 2 \\ 4 & 9 & & 9 & 4 \\ \cdots & \cdots & & \cdots & \cdots \end{array}$$

where the first element in the triangle is $a(6, 3)$.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

It turns out that the triangle of appearances of a block B is either of the type of Example 4.10 or the Pascal triangle with certain 'initial diagonals' removed.

LEMMA 4.11. *Let $B \in \mathcal{L}(\zeta)$ and assume B appears for the first time in B_{n_0,k_0}. Let $i_0 = n_0 - k_0 - 1$ and $j = k_0 - 1$. Then there are exactly five possible cases for the triangle of appearances of B.*

Case 1: $B = \boldsymbol{a}^{i_0}\boldsymbol{b}^{j_0}$. The number of occurrences of B satisfies the recurrence relation

$$a(n,k) = a(n-1,k) + a(n-1,k-1) + 1,$$

and the triangle of occurrences of B is

$$\begin{array}{ccccc} & & 1 & & \\ & 2 & & 2 & \\ 3 & & 5 & & 2 \\ 4 & 9 & & 9 & 4 \\ \cdots & \cdots & & \cdots & \cdots \end{array}$$

It follows that $a(n,k) = C(n - n_0 + 2, k - k_0 + 1) - 1$.

Case 2: $B = \cdots b\boldsymbol{a}^{i_0}\boldsymbol{b}^{j_0}a \cdots$ (B contains the telltale block). The number of occurrences of B satisfies the recurrence relation

$$a(n,k) = a(n-1,k) + a(n-1,k-1),$$

and the triangle of occurrences of B is just the Pascal triangle

$$\begin{array}{ccccc} & & 1 & & \\ & 1 & & 1 & \\ 1 & & 2 & & 1 \\ 1 & 3 & & 3 & 1 \\ \cdots & \cdots & & \cdots & \cdots \end{array}$$

It follows that $a(n,k) = C(n - n_0, k - k_0)$.

Case 3: $B = \boldsymbol{a}^{i}\boldsymbol{B}_{\boldsymbol{n_0-2,k_0-1}}\boldsymbol{b}^{j}$ for some $i < i_0$ and $j < j_0$. The number of occurrences of B satisfies the recurrence relation

$$a(n,k) = a(n-1,k) + a(n-1,k-1),$$

and the triangle of occurrences of B is

$$\begin{array}{ccccc} & & 2 & & \\ & 3 & & 3 & \\ 4 & & 6 & & 4 \\ 5 & 10 & & 10 & 5 \\ \cdots & \cdots & & \cdots & \cdots \end{array}$$

It follows that $a(n,k) = C(n - n_0 + 2, k - k_0 + 1)$.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

Case 4: $B = \cdots b\boldsymbol{a}^{i_0}\boldsymbol{b}^{j}$ for some $j \leq j_0$, and B is not as in Case 3. The number of occurrences of B satisfies the recurrence relation

$$a(n,k) = a(n-1,k) + a(n-1,k-1),$$

and the triangle of occurrences of B is the following:

$$\begin{array}{ccccc} & & 1 & & \\ & 2 & & 1 & \\ 3 & & 3 & & 1 \\ 4 & 6 & & 4 & 1 \\ \cdots & & \cdots & & \cdots \end{array}$$

It follows that $a(n,k) = C(n - n_0 + 1, k - k_0)$.

Case 5: $B = \boldsymbol{a}^{i}\boldsymbol{b}^{j_0}a\cdots$ for some $i \leq i_0$, and B is not as in Case 3. The number of occurrences of B satisfies the recurrence relation

$$a(n,k) = a(n-1,k) + a(n-1,k-1),$$

and the triangle of occurrences of B is the following:

$$\begin{array}{ccccc} & & 1 & & \\ & 1 & & 2 & \\ 1 & & 3 & & 3 \\ 1 & 4 & & 6 & 4 \\ \cdots & & \cdots & & \cdots \end{array}$$

It follows that $a(n,k) = C(n - n_0 + 1, k - k_0 + 1)$.

Proof. If B appears for the first time in B_{n_0,k_0}, then clearly B is one of the following:

(i) $B = \boldsymbol{a}^{i_0}\boldsymbol{b}^{j_0}$;

(ii) $B = \cdots b\boldsymbol{a}^{i_0}\boldsymbol{b}^{j_0}a\cdots$;

(iii) $B = \cdots b\boldsymbol{a}^{i_0}\boldsymbol{b}^{j}$ for some $j \leq j_0$;

(iv) $B = \boldsymbol{a}^{i}\boldsymbol{b}^{j_0}a\cdots$ for some $i \leq i_0$.

In Case (i), which corresponds to Case 1, the block B appears only once in B_{n_0,k_0}, and is made infinitely many times further down. More precisely, every concatenation of two adjacent basic blocks below n_0 creates a unique block B. It follows that $a(n,k) = a(n-1,k) + a(n-1,k-1) + 1$, and the triangle of occurrences of B is

$$\begin{array}{ccccc} & & 1 & & \\ & 2 & & 2 & \\ 3 & & 5 & & 2 \\ 4 & 9 & & 9 & 4 \\ \cdots & & \cdots & & \cdots \end{array}$$

In the previous triangle, $a(n + n_0, k + k_0)$ represents the number of finite paths from the root $(0,0)$ to the vertex (n,k) in a modified Pascal graph with extra 'wormholes'—paths

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press




FIGURE 15.

connecting $(0, 0)$ directly to each (n, k)—see Figure 15. It is not hard to see from this graph that $a(n + n_0, k + k_0) + 1$ is equal to the sum of the binomial coefficients inside the rectangle determined by $(0, 0)$ and (n, k), i.e.

$$a(n + n_0, k + k_0) + 1 = \sum_{i=0}^{n-k} \sum_{j=i}^{k+i} C(j, j - i).$$

Using well-known properties of the binomial coefficients we get

$$\begin{aligned} a(n + n_0, k + k_0) + 1 &= \sum_{i=0}^{n-k} \sum_{j=i}^{k+i} C(j, j - i) \\ &= \sum_{i=0}^{n-k} C(k + i + 1, k + 1) \\ &= C(n + 2, k + 1), \end{aligned}$$

showing as announced that $a(n, k) = C(n - n_0 + 2, k - k_0 + 1) - 1$.

In Case (ii), a similar argument as in the proof of Lemma 4.9 shows that B can be made only once. Therefore $a(n, k) = a(n-1, k)+a(n-1, k-1)$, and the triangle of appearances of B is just the Pascal triangle. Since the first element in the triangle is $a(n_0, k_0)$, it follows that $a(n, k) = C(n - n_0, k - k_0)$, establishing Case 2.

A special case of (iii) and (iv) is Case 3, i.e when $B = \boldsymbol{a^i B_{n_0-2,k_0-1} b^j}$ for some $i < i_0$ and $j < j_0$. Here, B is made twice at (n_0, k_0), and infinitely many times along the edges of the triangle below (n_0, k_0). First, observe that

$$B_{n_0,k_0} = b^{k_0} \cdots a^{n_0-k_0-2} B_{n_0-2,k_0-1} | B_{n_0-2,k_0-1} b^{k_0-2} \cdots\cdots a^{n_0-k_0}.$$

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

Therefore B appears twice in B_{n_0,k_0}:

$$b^{k_0}\cdots\overbrace{a^{n_0-k_0-2}B_{n_0-2,k_0-1}|b^{k_0-1}}^{B}\cdots\cdots a^{n_0-k_0}$$
$$b^{k_0}\cdots\cdots\underbrace{a^{n_0-k_0-1}|B_{n_0-2,k_0-1}b^{k_0-2}}_{B}\cdots\cdots a^{n_0-k_0}.$$

Second, note that B can be made exactly once at each vertex along the edges of the triangle below (n_0, k_0):

$$B_{n_0+j,k_0}=\cdots\overbrace{a^{n_0-k_0+j}|B_{n_0-2,k_0-1}b^{k_0-2}}^{B}\cdots,$$

and

$$B_{n_0+j,k_0+j}=\cdots\underbrace{a^{n_0-k_0-2}B_{n_0-2,k_0-1}|b^{k+j}}_{B}\cdots,$$

for all $j\geq 1$. Thus $a(n,k)=a(n-1,k)+a(n-1,k-1)$, and the triangle of occurrences of B is

$$\begin{array}{ccccccc}
 & & & 2 & & & \\
 & & 3 & & 3 & & \\
 & 4 & & 6 & & 4 & \\
5 & & 10 & & 10 & & 5 \\
\cdots & & \cdots & & \cdots & & \cdots
\end{array}$$

which is a subtriangle of the Pascal triangle; hence the relation $a(n,k)=C(n-n_0+1,k-k_0)$ follows.

To conclude, assume that B is as in (iii) but not of the type $B=a^iB_{n_0-2,k_0-1}b^j$. Then $B=\cdots ba^{n_0-k_0-1}b^j$ appears only once in B_{n_0,k_0}, because in order to appear twice it would have to contain the entire block B_{n_0-2,k_0-1}. For the same reason it cannot be made again along the vertices (n_0+j,k_0) for $j\geq 1$. Any two adjacent basic blocks inside the triangle below (n_0,k_0) will not produce any more B's either, because they are all of the type $\cdots ba^i|b^ja\cdots$ where $i>i_0$. On the other hand, B is created once more at each vertex (n_0+j,k_0+j) for $j\geq 1$:

$$B_{n_0+j,k_0+j}=\cdots\cdots\underbrace{ba^{n_0-k_0-1}|b^{k_0+j}}_{B}\cdots.$$

Therefore the triangle of appearances is

$$\begin{array}{ccccccc}
 & & & 1 & & & \\
 & & 2 & & 1 & & \\
 & 3 & & 3 & & 1 & \\
4 & & 6 & & 4 & & 1 \\
\cdots & & \cdots & & \cdots & & \cdots
\end{array}$$

As previously this is a subtriangle of the Pascal triangle; thus $a(n,k)=C(n-n_0+1,k-k_0)$, and Case 4 is established. Case 5 is similar. □

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

LEMMA 4.12. *Let $r \geq 0$ and $0 \leq s \leq r$. If $k_n/n \to \alpha$ as $n \to \infty$, then*

$$\lim_{n\to\infty} \frac{C(n-r, k_n - s)}{C(n, k_n)} = \alpha^s (1-\alpha)^{r-s}.$$

Remark 4. This was the key calculation used by Hajian to prove the ergodicity of μ_α for the Pascal adic—see [**14**], and also [**28**, Theorem 2.7].

Proof of Theorem 4.7. By (4.3) we need to show that

$$\lim_{n\to\infty} \frac{a(n, k_n)}{C(n, k_n)} = \nu_\alpha(B), \tag{4.4}$$

when $k_n/n \to \alpha$. According to Lemma 4.11, $a(n, k_n)$ is equal to one of the following:

Case 1: $C(n - n_0 + 2, k - k_0 + 1) - 1$,
Case 2: $C(n - n_0, k - k_0)$,
Case 3: $C(n - n_0 + 2, k - k_0 + 1)$,
Case 4: $C(n - n_0 + 1, k - k_0)$,
Case 5: $C(n - n_0 + 1, k - k_0 + 1)$.

In each of these cases Lemma 4.12 shows that the limit in (4.4) equals respectively:

Case 1: $\alpha^{k_0-1}(1-\alpha)^{n_0-k_0-1}$,
Case 2: $\alpha^{k_0}(1-\alpha)^{n_0-k_0}$,
Case 3: $\alpha^{k_0-1}(1-\alpha)^{n_0-k_0-1}$,
Case 4: $\alpha^{k_0}(1-\alpha)^{n_0-k_0-1}$,
Case 5: $\alpha^{k_0-1}(1-\alpha)^{n_0-k_0}$.

On the other hand, $\nu_\alpha(B) = \sum \nu_\alpha(B_{n,k})$, where the sum is taken over all (n, k) such that B is created in the basic block $B_{n,k}$, and if B is created twice in $B_{n,k}$, then $\nu_\alpha(B_{n,k})$ appears twice in the sum. In particular, if $B = B_{n,k}$ we have $\nu_\alpha(B) = \alpha^k(1-\alpha)^{n-k}$. So, for example, if B is as in Case 3, we have

$$\begin{aligned}
\nu_\alpha(B) &= 2\nu_\alpha(B_{n_0,k_0}) + \sum_{i=1}^{\infty} \nu_\alpha(B_{n_0+i,k_0+i}) + \sum_{i=1}^{\infty} \nu_\alpha(B_{n_0+i,k_0}) \\
&= \sum_{i=0}^{\infty} \nu_\alpha(B_{n_0+i,k_0+i}) + \sum_{i=0}^{\infty} \nu_\alpha(B_{n_0+i,k_0}) \\
&= \sum_{i=0}^{\infty} \alpha^{k_0+i}(1-\alpha)^{n_0-k_0} + \sum_{i=0}^{\infty} \alpha^{k_0}(1-\alpha)^{n_0-k_0+i} \\
&= \alpha^{k_0}(1-\alpha)^{n_0-k_0}\left[\sum_{i=0}^{\infty} \alpha^i + \sum_{i=0}^{\infty} (1-\alpha)^i\right] \\
&= \alpha^{k_0}(1-\alpha)^{n_0-k_0}\left[\frac{1}{1-\alpha} + \frac{1}{\alpha}\right] \\
&= \alpha^{k_0}(1-\alpha)^{n_0-k_0}\frac{\alpha + 1 - \alpha}{\alpha(1-\alpha)} \\
&= \alpha^{k_0-1}(1-\alpha)^{n_0-k_0-1}.
\end{aligned}$$

Similar computations show that the identity (4.4) also holds in the other cases. □

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

Remark 5. We could have used the ergodic theorem to show that the limit in (4.4), which is known to exist because of Lemma 4.11, must be equal to $\nu_\alpha(B)$: as noted previously in §4.2, for μ_α almost every path x we have $k_n(x)/n \to \alpha$ and

$$\lim_{n\to\infty} \frac{a(n, k_n(x))}{C(n, k_n)} = \nu_\alpha(B).$$

Remark 6. We thank Anthony Quas for an insight which leads to a proof simpler than the preceding one. It is possible to show that after a while (that is, below a certain level) 'new' creations of a block B due to concatenations constitute a negligible fraction of the total number of appearances of B. More specifically, fix an allowed block $B \in \mathcal{L}(\Sigma)$ and a positive integer m large enough that B appears at level m of the Pascal triangle of words. Consider a level n with $n \geq m$. As mentioned above, any basic block $B_{n,k}$ at this level factors uniquely into basic blocks of level m. Let us denote as 'old' appearances of B in $B_{n,k}$ the ones that are contained entirely in one of these basic blocks of level m, and call the rest 'new' appearances. The new appearances were created by concatenations after level m and contain division points of the factorization of $B_{n,k}$ into basic blocks $B_{m,j}$.

Let $D(n, k)$ denote the length of $B_{n,k}$ when regarded as a word on the alphabet whose symbols are basic blocks of level m; thus $D(n, k)$ is one more than the number of division points in the factorization under discussion. We have

$$\begin{aligned} D(m, j) &= 1 \quad \text{for all } j = 0, 1, \ldots, m, \\ D(n, 0) = D(n, n) &= 1 \quad \text{for all } n \geq m, \\ D(n, k) = D(n-1, k-1) + D(n-1, k) &\quad \text{for all } n > m, k = 1, \ldots, n-1. \end{aligned}$$

Viewing the array whose (n, k) entry is $D(n, k)$ as the sum of $m + 1$ Pascal triangles originating at all the places in level m (since each of these places has an entry 1, which is the 'seed' for an entire Pascal triangle below it), we have

$$D(n, k) = \sum_{j=0}^{m} C(n-m, k-j).$$

By Lemma 4.12, for $n \gg m$ and $k/n \approx \alpha$,

$$\frac{D(n, k)}{C(n, k)} = \sum_{j=0}^{m} \frac{C(n-m, k-j)}{C(n, k)} \approx \sum_{j=0}^{m} \alpha^j (1-\alpha)^{m-j} \leq (\sup\{\alpha, 1-\alpha\})^{m+1},$$

so only a small fraction of the appearances of B in $B_{n,k}$ are 'new'.

The frequency of appearance of $B_{m,j}$ in $B_{n,k}$ is $C(n-m, k-j)/C(n, k)$, which has a limiting value as $n \to \infty, k/n \to \alpha$. Taking into account the number of times that our given block B appears in each basic word $B_{m,j}$ at level m, since we can ignore the 'new' appearances of B in $B_{n,k}$ we see that B also has a limiting frequency of appearance in $B_{n,k}$ as $n \to \infty, k/n \to \alpha$ (and we can compute it, getting the same answer as before).

4.3. *Topological weak mixing.* In this section we prove that the countable-substitution subshift version of the Pascal adic is topologically weakly mixing. Since the system is not minimal (for example, it contains the two fixed points of $\{a, b\}^{\mathbb{Z}}$), it is not enough

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

to prove that there are no non-constant continuous eigenfunctions (cf. [**26**]). Instead, we use a characterization of topological weak mixing provided by Keynes and Robertson [**19**] (see also [**18**]), along with Weyl's theorem on uniform distribution.

In all the following (X, T) denotes a topological dynamical system, i.e. X is a compact metric space and $T : X \to X$ is a homeomorphism. Recall that (X, T) is topologically ergodic (i.e. topologically transitive) if there is a point $x \in X$ with a dense orbit. Since the Pascal adic is ergodic for the Bernoulli measures μ_α, it follows that the (fully supported) image measure $\phi(\mu_\alpha)$ is ergodic for the substitution subshift, and therefore the system is topologically ergodic. Actually, if x is a path in the Pascal graph which is not eventually diagonal, then $\phi(x)$ has a dense orbit in Σ, so in fact all but countably many orbits in Σ are dense. We say that (X, T) is *topologically weakly mixing* if $(X \times X, T \times T)$ is topologically ergodic.

Definition 4.13. Let X be a topological space. Denote by $\mathcal{C}(X)$ the space of all continuous functions $f : X \to \mathbb{C}$, and by $\mathcal{B}(X)$ the set of bounded functions $f : X \to \mathbb{C}$ such that the set $\mathcal{C}(f)$ of points of continuity of f is residual.

Definition 4.14. Let $f, g \in \mathcal{B}(X)$. We say that f and g are *essentially equal*, and write $f \overset{\text{ess}}{=} g$, if $f = g$ on $\mathcal{C}(f) \cap \mathcal{C}(g)$.

Remark 7. If $f, g \in \mathcal{B}(X)$, then $f \overset{\text{ess}}{=} g$ if and only if $f = g$ on a dense set.

THEOREM 4.15. [**19**] *Let (X, T) be a topological dynamical system. The following are equivalent:*

(i) *(X, T) is topologically ergodic;*

(ii) *for every $f \in \mathcal{B}(X)$, if $f \circ T \overset{\text{ess}}{=} f$, then $f \overset{\text{ess}}{=}$ constant;*

(iii) *for every $f \in \mathcal{B}(X)$, if $f \circ T = f$ (everywhere), then $f \overset{\text{ess}}{=}$ constant.*

Remark 8. It is easy to show that if (X, T) is ergodic, and $\lambda \in \mathbb{C}$ is an eigenvalue for some $f \in \mathcal{B}(X)$, then $|\lambda| = 1$ and $|f| \overset{\text{ess}}{=}$ constant.

Definition 4.16. A Borel probability measure μ on X is called *closed ergodic* for T if every closed invariant subset of X has μ-measure 0 or 1.

Here is a criterion for weak mixing that we will use to show that the substitution dynamical system (Σ, σ) is topologically weakly mixing.

THEOREM 4.17. [**19**] *Let (X, T) be a topological dynamical system and suppose there exists a T-invariant Borel probability measure μ supported on all of X which is closed ergodic. The following are equivalent:*

(i) *(X, T) is topologically weakly mixing;*

(ii) *for every $f \in \mathcal{B}(X)$, if there is $\lambda \in \mathbb{C}$ such that $f \circ T = \lambda f$ (everywhere), then $f \overset{\text{ess}}{=}$ constant.*

Every measure ν_α is ergodic (so in particular closed ergodic) and has support equal to Σ, so Theorem 4.17 applies in our case. Given $f \in \mathcal{B}(\Sigma)$ and $\lambda \in \mathbb{C}$ such that $f \circ \sigma = \lambda f$ (everywhere), if we can show that $\lambda = 1$ then, combining Theorem 4.15 and Theorem 4.17, one would show that (Σ, σ) is topologically weakly mixing.

We need the following lemma.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

LEMMA 4.18. *Let (X, T) be a topological dynamical system (with underlying metric d). Let $f : X \to \mathbb{C}$ be a function with $\mathcal{C}(f) \neq \emptyset$ and such that $f \circ T = \lambda f$ for some $\lambda \in \mathbb{C}$. Then every point $x \in X$ with dense forward orbit and dense backward orbit is in $\mathcal{C}(f)$.*

Proof. Let $x \in X$ be a point such that both semiorbits $\{T^n x \mid n \geq 0\}$ and $\{T^n x \mid n \leq 0\}$ are dense. Let $z \in \mathcal{C}(f)$. Note that if $\lambda = 0$ then $f \equiv 0$, so we can assume that $|\lambda| > 0$. If $x \notin \mathcal{C}(f)$ then there exists $\alpha > 0$ such that for all $\delta > 0$ there is $y \in X$ such that

$$d(x, y) < \delta \quad \text{and} \quad |f(x) - f(y)| \geq \alpha. \tag{4.5}$$

Since $z \in \mathcal{C}(f)$, there exists $\eta > 0$ such that

$$\text{if } d(z, u) < \eta \quad \text{then } |f(z) - f(u)| < \alpha/4. \tag{4.6}$$

Since x has dense forward orbit we can find $n > 0$ such that $d(T^n x, z) < \eta/2$. By continuity of T^n there exists $\delta > 0$ such that

$$\text{if } d(x, y) < \delta \quad \text{then } d(T^n x, T^n y) < \eta/2.$$

For that previous δ, by (4.5), there exists y such that

$$d(x, y) < \delta \quad \text{and} \quad |f(x) - f(y)| \geq \alpha.$$

If $|\lambda| \geq 1$ then it follows that

$$|f \circ T^n x - f \circ T^n y| = |\lambda^n f(x) - \lambda^n f(y)| = |\lambda|^n |f(x) - f(y)| \geq \alpha.$$

If $|\lambda| < 1$ then the same inequality holds since x has dense backward orbit and we could have chosen $n < 0$. On the other hand, since $d(T^n x, z) < \eta/2$ and $d(T^n y, z) < \eta$, by (4.6) we get the following contradiction:

$$|f \circ T^n x - f \circ T^n y| < |f \circ T^n x - f(z)| + |f(z) - f \circ T^n y| < \alpha/2.$$

Thus, $x \in \mathcal{C}(f)$. □

THEOREM 4.19. *The countable substitution dynamical system (Σ, σ) is topologically weakly mixing.*

Proof. Let $f \in \mathcal{B}(\Sigma)$ be such that $f \circ \sigma = \lambda f$ for some $\lambda \in \mathbb{C}$. Assume that $\lambda \neq 1$. Since there are no rational eigenvalues (see [**27**]), $\lambda = e^{2\pi i \beta}$ for some irrational number β. Suppose that we can find $\omega \in \Sigma$ with the properties that:

(i) ω has dense orbit;

(ii) there exist $N_k \to \infty$ such that $\lambda^{N_k} \to -1$;

(iii) $\sigma^{N_k} \omega \to \omega$.

Then, by Lemma 4.18, since every point with dense orbit is a continuity point of f, the relation $f \circ \sigma^{N_k} \omega = \lambda^{N_k} f(\omega)$ would lead to a contradiction. To find such a point ω we use Weyl's theorem on uniform distribution. Since β is irrational, for every $k \geq 1$ the set $\{C(n, k)\beta : n \in \mathbb{N}\}$ is uniformly distributed modulo 1. Therefore there exist $n_1 < n_2 < \cdots < n_k < \cdots$ such that

$$|\lambda^{C(n_k, k)} + 1| < \frac{1}{k} \quad \text{for all } k. \tag{4.7}$$

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

Let x be the path in the Pascal graph defined by

$$x = 01^{n_1}01^{n_2-n_1-1}01^{n_3-n_2-1}0\cdots 1^{n_k-n_{k-1}-1}0\cdots .$$

Since x is not eventually diagonal, $\omega = \phi(x)$ has a dense orbit in Σ, and hence condition (i) is satisfied. Let $N_k = C(n_k, k)$. Condition (ii) follows from (4.7), and condition (iii) is guaranteed by the kink lemma (Lemma 2.2). □

5. *Questions*

Many properties of the Pascal adic and related systems remain to be determined. In particular, the question of weak mixing of the systems (X, T, μ_p) remains open. Since ergodicity of the Bernoulli measures μ_p under the Pascal adic map T implies the Hewitt-Savage 0,1 law, and weak mixing is stronger than ergodicity, results along these lines would have probabilistic implications.

Recently the loose Bernoulli property of each (X, T, μ_p) has been established by de la Rue and Janvresse [**6**]. We believe that these systems have infinite rank, and indeed that they do not have local rank 1 (see [**10**] for the definitions). Determination of the joinings, or even factors, of these systems would be of considerable interest, as would any more information about their spectra (simple? singular?).

Dynamical properties of a class of adic transformations generalizing the Pascal adic, which code adic transformations on certain shifts of finite type (see [**28**]), are studied in a forthcoming paper by the first-named author [**23**].

Acknowledgements. The first author was partially supported by the Millennium Nucleus in Information and Randomness, Programa Iniciativa Cientifica Milenio P01-005. We thank R. Burton, R. Kenyon, and A. Quas for helpful conversations. This paper is based on the Ph.D. dissertation of the first author [**24**], written under the direction of the second author.

References

[1] V. Afraimovich, J.-R. Chazottes and B. Saussol. Local dimensions for Poincaré recurrences. *Electron Res. Announc. Amer. Math. Soc.* **6** (2000), 64–74.

[2] R. Bollinger and C. Burchard. Lucas's theorem and some related results for extended Pascal triangles. *Amer. Math. Monthly* **97**(3) (1990), 198–204.

[3] J. Cassaigne, P. Hubert and S. Troubetzkoy. Complexity and growth for polygonal billiards. *Ann. Inst. Fourier (Grenoble)* **52** (2002), 835–847.

[4] Z. Coelho. Asymptotic laws for symbolic dynamical systems. *Topics in Symbolic Dynamics and Applications (Temuco 1997) (London Mathematical Society Lecture Note Series, 279)*. Cambridge University Press, Cambridge, 1997, pp. 123–165.

[5] P. Collet and A. Galves. Asymptotic distribution of entrance times for expanding maps of the interval. *Dynamical systems and applications—World Sci. Ser. Appl. Anal.* **4** (1995), 139–152.

[6] T. de la Rue and E. Janvresse. The Pascal adic transformation is loosely Bernoulli. *Annales de l'IHP, Probab. Stat.* **40** (2004), 133–139.

[7] B. Durand, F. Host and C. Skau. Substitutional dynamical systems, Bratteli diagrams and dimension groups. *Ergod. Th. & Dynam. Sys.* **19** (1999), 953–993.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

[8] F. Durand and A. Maass. Limit laws of entrance times for low-complexity Cantor minimal systems. *Nonlinearity* **14**(4) (2001), 683–700.
[9] S. Ferenczi. Rank and symbolic complexity. *Ergod. Th. & Dynam. Sys.* **16** (1996), 663–682.
[10] S. Ferenczi. Systems of finite rank. *Colloq. Math.* **73**(1) (1997), 36–65.
[11] S. Ferenczi. Complexity of sequences and dynamical systems. *Discrete Math.* **206** (1999), 145–154.
[12] N. P. Fogg. *Substitutions in Dynamics, Arithmetics, and Combinatorics (Lecture Notes in Mathematics, 1794)*. Springer, Berlin, 2002.
[13] A. Forrest. K-groups associated with substitution minimal systems. *Israel J. Math.* **98** (1997), 101–139.
[14] Y. Hajian, A. Ito and S. Kakutani. Invariant measures and orbits of dissipative transformations. *Adv. Math* **9** (1972), 52–65.
[15] I. Herman, R. Putnam and C. Skau. Ordered Bratteli diagrams, dimension groups and topological dynamics. *Internat. J. Math.* **3**(6) (1992), 827–864.
[16] B. Host. Substitution subshifts and Bratteli diagrams. *Topics in Symbolic Dynamics and Applications (London Mathematical Society Lecture Note Series)*. Eds. F. Blanchard, A. Maass and A. Nogueira. Cambridge University Press, Cambridge 1999.
[17] S. Ito. Construction of transversal flows for maximal Markov automorphisms. *Tokyo J. Math.* **1** (1978), 305–324.
[18] H. Keynes. *Lectures on Ergodic Theory*. University of Minnesota, 1971–1972.
[19] H. Keynes and J. Robertson. Eigenvalue theorems in topological transformation groups. *Trans. Amer. Math. Soc.* **139** (1969), 359–369.
[20] Y. Lacroix. Possible limit laws for entrance times of an ergodic aperiodic dynamical system. *Israel J. Math.* **132** (2002), 253–263.
[21] A. Livshitz. A sufficient condition for weak mixing of substitutions and stationary adic transformations. *Math. Notes* **44** (1988), 920–925.
[22] E. Lucas. Théorie des fonctions numériques simplement périodiques. *Amer. J. Math.* **1** (1878), 184–240.
[23] X. Méla. Dynamics of a class of nonstationary adic transformations, in preparation.
[24] X. Méla. Dynamical properties of the Pascal adic and related systems. *PhD Thesis*, University of North Carolina at Chapel Hill, 2002.
[25] F. Paccaut. Statistics of return times for weighted maps of the interval. *Ann. Inst. Henri Poincaré, Probabilités et Statistiques* **36**(3) (2000), 339–366.
[26] K. Petersen. Disjointness and weak mixing of minimal sets. *Proc. Amer. Math. Soc.* **24** (1970), 278–280.
[27] K. Petersen. Information compression and retention in dynamical processes. *Dynamics and Randomness (Proceedings of the December 2000 Workshop in Santiago, Chile)*. Eds. A. Maass, S. Martinez and J. San Martin. Kluwer Academic Publishers, Dordrecht, 2002, pp. 147–217.
[28] K. Petersen and K. Schmidt. Symmetric Gibbs measures. *Trans. Amer Math. Soc.* **349**(7) (1997), 2775–2811.
[29] B. Pitskel. Poisson limit law for Markov chains. *Ergod. Th. & Dynam. Sys.* **11**(3) (1991), 501–513.
[30] M. Queffelec. *Substitution Dynamical Systems—Spectral Analysis (Lecture Notes in Mathematics, 1294)*. Springer, Berlin, 1980.
[31] B. Saussol. On fluctuations and the exponential statistics of return times. *Nonlinearity* **14** (2001), 179–191.
[32] K. Schmidt. Invariant cocycles, random tilings, and the super-K and strong Markov properties. *Trans. Amer. Math. Soc.* **349** (1997), 2812–2825.
[33] K. Schmidt. Tail-fields of products of random variables and ergodic equivalence relations. *Ergod. Th. & Dynam. Sys.* **19** (1999), 1325–1341.
[34] Ja. G. Sinai. Probabilistic ideas in ergodic theory. *Amer. Math. Soc. Transl. Ser.* 2 **31** (1963), 62–84.
[35] B. Solomyak. On the spectral theory of adic transformations. *Adv. Soviet Math.* **9** (1992), 217–230.
[36] B. Solomyak. Substitutions, adic transformations, and beta-expansions. *Contemp. Math.* **135** (1992), 361–372.
[37] A. Vershik. Description of invariant measures for actions of some infinite groups. *Dokl. Akad. Nauk SSSR* **28** (1974), 749–752.
[38] A. Vershik. Uniform algebraic approximation of shift and multiplicative operators. *Dokl. Akad. Nauk SSSR* **218**(24) (1981), 526–529.
[39] A. Vershik. A theorem on the Markov periodic approximation in ergodic theory. *J. Sov. Math.* **28** (1985), 667–673.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press

[40] A. Vershik. The adic realizations of the ergodic actions with the homeomorphisms of the Markov compact and the ordered Bratteli diagrams (English translation). *J. Math. Sci.* (New York) **87**(6) (1997), 4054–4058.

[41] A. Vershik and S. Kerov. Asymptotics of the Plancherel measure of the symmetric group and the limiting form of Young tables. *Sov. Math. Dokl.* **18** (1977), 527–531.

[42] A. Vershik and S. Kerov. Asymptotic theory of characters of the symmetric group. *Func. Anal. Appl.* **15** (1981), 246–255.

[43] A. Vershik and S. Kerov. Asymptotics of the largest and typical dimensions of irreducible representations of a symmetric group. *Func. Anal. Appl.* **19** (1985), 21–31.

[44] A. Vershik and A. Livshitz. Adic models of ergodic transformations, spectral theory, substitutions, and related topics. *Adv. Soviet Math.* **9** (1992), 185–204.

https://doi.org/10.1017/S0143385704000173 Published online by Cambridge University Press