Math. Proc. Camb. Phil. Soc. (1987), **102**, 297
Printed in Great Britain

Relative semicharacteristic classes

BY JAMES F. DAVIS
Department of Mathematics, Indiana University, Bloomington, IN 47405, *U.S.A.*

(*Received* 18 *August* 1986; *revised* 21 *January* 1987)

§0. *Introduction*

In 1973 Ronnie Lee introduced the notion of semicharacteristic classes, which are invariants of the bordism group $\mathfrak{N}_*(B\pi)$ of closed manifolds equipped with a free action of a finite group π. In this paper we relativize his theory. Associated to a homomorphism $G \to \pi$ of finite groups, there is the relative bordism group $\mathfrak{N}_*(BG \to B\pi)$, which is the bordism group of compact manifolds M with a free π-action, so that the action on ∂M is induced from a free G-action, i.e. $\partial M = \pi \times_G N$ for some manifold N with a free G-action. The invariants defined here are invariants of this relative group.

Lee introduced his invariants to give homological restrictions on closed manifolds equipped with a free action of the dihedral group D_{2r}. We will apply our theory to study certain non-free actions. In §2 relative semicharacteristic classes are applied to prove:

THEOREM A. *Let D_{2r} ($r > 1$, r odd) act smoothly on a closed manifold M^{2m+1}, so that the generator g of $\mathbb{Z}_r \subset D_{2r}$ has no fixed points. Then if*

$$\sum_{i=0}^{m} \operatorname{trace}(g_*: H_i(M; \mathbb{F}_2) \to H_i(M; \mathbb{F}_2)) = 1 \in \mathbb{F}_2,$$

the singular set of the action has dimension $\geqslant m$. More precisely, if $\mathscr{S}$ denotes the singular set, then $H_m(\mathscr{S}; \mathbb{F}_2) \neq 0$.

COROLLARY B. *If D_{2r} ($r > 1$, r odd) acts smoothly on a closed manifold M^{2m+1} which has the* mod 2 *homology of a sphere S^{2m+1}, projective space $\mathbb{R}P^{4k+1}$, or a quaternionic space form $S^{8k+3}/Q(2^n)$ ($n \geqslant 3$), so that $\mathbb{Z}_r \subset D_{2r}$ acts freely, then the fixed set of the involution T ($\langle T \rangle = \mathbb{Z}_2 \subset D_{2r}$) has a component of dimension $\geqslant m$.*

Corollary B implies that there are no free dihedral actions on mod 2 homology spheres, a fact which was first proved by J. Milnor. R. Lee proved the general case of Theorem A for *free* dihedral actions. In the case where M is a mod 2 homology sphere, Smith theory implies that the fixed set M^T is a mod 2 homology sphere; since the singular set $\mathscr{S} = D_{2r} \times_{\mathbb{Z}_2} M^T$, all components of the singular set have the same dimension k. Thus the condition $H_m(\mathscr{S}; \mathbb{F}_2) \neq 0$ implies that $k = m$, and we are led to a new proof of a 'regularity' result of D. Montgomery and C. T. Yang:

COROLLARY C. *If D_{2r} (r odd, $r > 1$) acts smoothly on a* mod 2 *homology sphere M^{2m+1} such that $\mathbb{Z}_r \subset D_{2r}$ acts freely, then the singular set is m-dimensional.*

https://doi.org/10.1017/S0305004100067311 Published online by Cambridge University Press

§1. *Relative semicharacteristic classes*

Let π be a finite group. Fix a field K of characteristic 2. Let $K\pi$ denote the group ring.

Definition 1·1. A symmetric K-bilinear form $\phi\colon V \times V \to K$ on a finitely generated $K\pi$-module V is *Lee even* if

(i) $\phi(gx, gy) = \phi(x, y)$ for all $x, y \in V$, $g \in \pi$,

(ii) $\phi(tx, x) = 0$ for all $x \in V$ and for all elements t of π of order 2 (i.e. $t^2 = e$, $t \neq e$).

If ϕ is non-singular, we say the module V is *Lee even*.

Example (see [1]). If π acts freely on a compact manifold W^{2n}, then the intersection pairing $\phi\colon H_n(W; K) \times H_n(W; K) \to K$ is Lee even.

R. Lee defined the following Grothendieck group.

Definition 1·2. $\tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(\pi)$ is the abelian group with:

Generators: $[V]$, isomorphism classes of finitely generated $K\pi$-modules;

Relations: (*a*) if $0 \to V' \to V \to V'' \to 0$ is an exact sequence of finitely generated $K\pi$-modules, then $[V] - [V'] - [V''] = 0$,

(*b*) $[K\pi] = 0$,

(*c*) if V is Lee even, then $[V] = 0$.

Definition 1·3. If M^{2m+1} is a compact manifold with a π-action, the *Lee semicharacteristic* is

$$\chi_{\frac{1}{2}}(M; K) = \sum_{i=0}^{m} (-1)^i [H_i(M; K)] \in \tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(\pi).$$

Let $\mathfrak{N}_n(B\pi)$ denote the unoriented bordism group of pairs $(M^n, f\colon M^n \to B\pi)$ where M^n is a closed manifold. Let $\tilde{M}$ denote the induced π-cover of M. Equivalently $\mathfrak{N}_n(B\pi)$ is the unoriented bordism group of closed manifolds $\tilde{M}$ equipped with a free π-action.

THEOREM 1·4 (Lee). *The map* $(M, f) \mapsto \chi_{\frac{1}{2}}(\tilde{M}; K)$ *gives a well-defined homomorphism* $\chi_{\frac{1}{2}}\colon \mathfrak{N}_{2m+1}(B\pi) \to \tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(\pi)$.

Now for the relative version of the theory.

Definition 1·5. Let $\varphi\colon G \to \pi$ be a homomorphism of finite groups. Then

$$\tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(G \to \pi) = \tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(\pi) / \langle [K\pi \otimes_{KG} V]\colon V \text{ is a } f.g.\ KG\text{-module} \rangle.$$

If φ is an injection, then $\tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(G \to \pi) = \mathrm{coker}\,(\tilde{\varphi})$ where $\tilde{\varphi}\colon R_{\mathrm{GL},\,\mathrm{ev}}(G) \to R_{\mathrm{GL},\,\mathrm{ev}}(\pi)$ is defined by $\tilde{\varphi}[V] = [K\pi \otimes_{KG} V]$. In particular, if G is the trivial group then the relative group reduces to $R_{\mathrm{GL},\,\mathrm{ev}}(\pi)$.

Let $\mathfrak{N}_n(BG \to B\pi)$ denote the bordism group of triples

$$(M^n, f\colon M \to B\pi, f_\partial\colon \partial M \to BG)$$


where M is a compact n-dimensional manifold with boundary ∂M and

$$\begin{array}{ccc} \partial M & \hookrightarrow & M \\ \downarrow{\scriptstyle f_\partial} & & \downarrow{\scriptstyle f} \\ BG & \longrightarrow & B\pi \end{array}$$

commutes up to homotopy. Equivalently, we are given a compact manifold $\tilde{M}$ with a free π-action, so that the action on $\partial\tilde{M}$ is induced from a G-action, i.e. $\partial\tilde{M} = \pi \times_G N$

https://doi.org/10.1017/S0305004100067311 Published online by Cambridge University Press

for some manifold N with a free G-action. (Here $\pi \times_G N = \pi \times N/(\rho, gn) \sim (\rho\varphi(g), n)$.) A triple (M, f, f_∂) is zero in $\mathfrak{N}_n(BG \to B\pi)$ if there exists a quadruple $(W^{n+1}, \partial', g\colon W \to B\pi, g_{\partial'}\colon \partial' \to BG)$ where W and ∂' are compact manifolds, $\partial W = \partial' \cup M, \partial\partial' = \partial M = \partial' \cap M$, and the following diagram commutes up to homotopy:



There is the familiar long exact sequence

$$\ldots \to \mathfrak{N}_n(BG) \to \mathfrak{N}_n(B\pi) \to \mathfrak{N}_n(BG \to B\pi) \to \mathfrak{N}_{n-1}(BG) \to \ldots$$

Definition 1·6. A triple $(M^{2m+1}, f\colon M \to B\pi, f_\partial\colon \partial M \to BG)$ is *induced* if

$$\ker (i_*\colon H_m(\partial\tilde{M}; K) \to H_m(\tilde{M}; K)) = K\pi \otimes_{KG} V$$

for some KG-submodule V of $H_m(\overline{\partial M}; K)$. Here $\tilde{M}$ denotes the π-cover induced by f and $\overline{\partial M}$ denotes the G-cover induced by f_∂. The equality above is interpreted by identifying $H_m(\partial\tilde{M}; K)$ with the induced module $K\pi \otimes_{KG} H_m(\overline{\partial M}; K)$. Let $\mathfrak{N}I_{2m+1}(BG \to B\pi)$ denote the subgroup of $\mathfrak{N}_{2m+1}(BG \to B\pi)$ generated by induced triples.

The most common example of an induced triple would be when i_* is injective.

THEOREM 1·7. *There is a homomorphism*

$$\chi_{\frac{1}{2}}\colon \mathfrak{N}I_{2m+1}(BG \to B\pi) \to \tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(G \to \pi)$$

defined by sending any induced triple (M, f, f_∂) *representing a bordism class to* $\chi_{\frac{1}{2}}(\tilde{M}; K)$.

Proof. It suffices to show that $\chi_{\frac{1}{2}}(\tilde{M}; K) = 0 \in \tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(G \to \pi)$ for an induced triple (M, f, f_∂) which is the boundary of a quadruple $(W, \partial', g, g_{\partial'})$. By Lee's result $\chi_{\frac{1}{2}}(\partial\tilde{W}; K) = 0 \in \tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(\pi)$. Consider the Mayer–Vietoris exact sequence

$$0 \to \ker\alpha \cap \ker\beta \to H_m(\partial\tilde{M}; K) \xrightarrow{\alpha\oplus\beta} H_m(\tilde{M}; K) \oplus H_m(\tilde{\partial}'; K) \to H_m(\partial\tilde{W}; K) \to \ldots \to H_0(\partial\tilde{W}; K) \to 0.$$

(Here the tilde always represents the induced π-cover.) The Euler characteristic of an exact sequence is zero, hence $\chi_{\frac{1}{2}}(\tilde{M}; K)$ equals

$$-\chi_{\frac{1}{2}}(\tilde{\partial}'; K) + \sum (-1)^i [H_i(\partial\tilde{M}; K)] + (-1)^{m+1}[\ker\alpha \cap \ker\beta] + \chi_{\frac{1}{2}}(\partial\tilde{W}; K),$$

which is zero in $\tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(G \to \pi)$ since all the modules involved in the first three terms are induced.

Remark. An analogous theorem could be proved for oriented bordism groups with the target of the homomorphism being a relative version of Lee's orthogonal or symplectic Grothendieck groups.

§2. *Proof of Theorem A*

Let $D_{2r} = \langle g, T \mid g^r = T^2 = e, TgT = g^{-1}\rangle$ denote the dihedral group. It is a semi-direct product $D_{2r} = \mathbb{Z}_r \rtimes \mathbb{Z}_2 = \langle g\rangle \rtimes \langle T\rangle$.

https://doi.org/10.1017/S0305004100067311 Published online by Cambridge University Press

Given an action of π on X we define the singular set

$$\mathscr{S} = \mathscr{S}(X, \pi) = \{x \in X \mid \quad gx = x \quad \text{for some} \quad g \in \pi - \{e\}\}.$$

LEMMA 2·1. *If D_{2r} (r odd) acts on a space M so that $\mathbb{Z}_r$ acts freely, then $\mathscr{S}(M, D_{2r}) = D_{2r} \times_{\mathbb{Z}_2} \mathscr{S}(M, \mathbb{Z}_2)$. If in addition the action is smooth on a closed manifold M and $T(\mathscr{S})$ is a D_{2r}-invariant closed tubular neighbourhood of $\mathscr{S}(M, D_{2r})$, then $(M - \operatorname{int} T(\mathscr{S}))/D_{2r}$ represents an element of $\mathfrak{N}_n(B\mathbb{Z}_2 \to BD_{2r})$.*

Proof. A map $f\colon D_{2r} \times_{\mathbb{Z}_2} \mathscr{S}(M, \mathbb{Z}_2) \to \mathscr{S}(M, D_{2r})$ is given by $f(\rho, x) = \rho x$. We define f^{-1} as follows. For any $x \in \mathscr{S}(M, D_{2r})$, $g^i Tx = x$ for some $g^i \in \mathbb{Z}_r$. Furthermore, this element g^i is unique, since $g^i Tx = g^k Tx$ implies that $g^i = g^k$ since $\mathbb{Z}_r$ acts freely. Choosing j so that $2j \equiv i (\operatorname{mod} r)$ one easily checks that $Tg^{-j}x = g^{-j}x$, and we define $f^{-1}(x) = (g^j, g^{-j}x)$. One sees directly that f and f^{-1} are inverses. If the action is smooth on a manifold then

$$f\colon D_{2r} \times_{\mathbb{Z}_2} T(\mathscr{S}(M, \mathbb{Z}_2)) \to T(\mathscr{S}(M, D_{2r}))$$

is also a homeomorphism. The lemma follows.

LEMMA 2·2. *If $\pi = H \rtimes G$ with H of odd order then $\mathfrak{N}_*(BG \to B\pi) = 0$.*

Proof. If $\pi_1 \to \pi_2$ induces an isomorphism $H_*(\pi_1; \mathbb{Z}_{(2)}) \xrightarrow{\sim} H_*(\pi_2; \mathbb{Z}_{(2)})$ and $\mathscr{H}$ is a generalized homology theory, then $\mathscr{H}_*(B\pi_1) \otimes \mathbb{Z}_{(2)} \to \mathscr{H}_*(B\pi_2) \otimes \mathbb{Z}_{(2)}$ is an isomorphism. Indeed the map induces a $\mathbb{Z}_{(2)}$-isomorphism on the E_2-term of the Atiyah–Hirzebruch spectral sequence and hence on E_∞. Unoriented bordism $\mathfrak{N}_*$ is a generalized homology theory and $\mathfrak{N}_* = \mathfrak{N}_* \otimes \mathbb{Z}_{(2)}$ since it is 2-torsion.

Thus we must show $i_*\colon H_*(G; \mathbb{Z}_{(2)}) \to H_*(\pi; \mathbb{Z}_{(2)})$ is an isomorphism. Since, in the exact sequence

$$1 \to H \to \pi \xrightarrow{p} G \to 1,$$

p is split by $i\colon G \to \pi$, i_* is an injection. The order of H is odd, so a transfer argument shows $H_*(H; \mathbb{Z}_{(2)}) = 0$. The Lyndon–Hochschild–Serre spectral sequence then shows $p_*\colon H_*(\pi; \mathbb{Z}_{(2)}) \to H_*(G; \mathbb{Z}_{(2)})$ is an isomorphism, so i_* is surjective.

So the plan of attack on Theorem A is clear; one needs to compute $\tilde{R}_{\mathrm{Gl},\,\mathrm{ev}}(\mathbb{Z}_2 \to D_{2r})$ and $\chi_{\frac{1}{2}}(M - \operatorname{int} T(\mathscr{S}); K)$.

LEMMA 2·3. *Let V be a finitely generated KD_{2r}-module with r odd. Then $[V] = 0 \in \tilde{R}_{\mathrm{Gl},\,\mathrm{ev}}(\mathbb{Z}_2 \to D_{2r})$ if and only if $\operatorname{tr}(h\colon V \to V) = 0 \in K$ for all $h \in \mathbb{Z}_r - \{e\}$.*

Remark. Although this is true for any field K of characteristic 2, for simplicity we will only prove and use this in the case where K contains a primitive r-th root of unity ζ.

Proof. There is one 1-dimensional representation of D_{2r}, the trivial representation, denoted V_0. There are $(r-1)/2$ irreducible representations V_i $(1 \leqslant i \leqslant (r-1)/2)$ defined by

$$g \to \begin{bmatrix} \zeta^i & 0 \\ 0 & \zeta^{-i} \end{bmatrix}$$

$$T \to \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

The computations of Lee then show

$$\tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(D_{2r}) = (\mathbb{Z}_2)^{(r+1)/2}$$

with generators $[V_0], [V_1], \ldots, [V_{(r-1)/2}]$. A composition series for any $K\mathbb{Z}_2$-module has

https://doi.org/10.1017/S0305004100067311 Published online by Cambridge University Press

all factors isomorphic to K and we know $KD_{2r} \otimes_{K\mathbb{Z}_2} K \cong V_0 \oplus V_1 \oplus \ldots \oplus V_{(r-1)/2}$. It follows that

$$\tilde{R}_{\mathrm{GL},\,\mathrm{ev}}(\mathbb{Z}_2 \to D_{2r}) = (\mathbb{Z}_2)^{(r+1)/2}/\Delta(\mathbb{Z}_2),$$

where $\Delta\colon \mathbb{Z}_2 \to (\mathbb{Z}_2)^{(r+1)/2}$ is the diagonal map.

Let V be a finitely generated KD_{2r}-module, choose a composition series for V, and let a_i be the number of times V_i occurs as a factor module. Let

$$\bar{a} = (a_0, a_1, a_2, \ldots, a_{(r-1)/2}, a_{(r-1)/2}, \ldots, a_2, a_1).$$

Then $\mathrm{tr}\,(g^k\colon V \to V)$ is the kth component of $M\bar{a}$, where

$$M = \begin{bmatrix} 1 & \zeta & \ldots & \zeta^{r-1} \\ 1 & \zeta^2 & & \zeta^{2(r-1)} \\ \vdots & & & \\ 1 & \zeta^r & & \zeta^{r(r-1)} \end{bmatrix}.$$

M is a van der Monde determinant; in particular it is non-singular. The matrix obtained by deleting the last row thus has 1-dimensional kernel, generated by $(1, 1, \ldots, 1)$. So $\mathrm{tr}\,(g^k\colon V \to V) = 0 \in K$ for all $1 \leqslant k \leqslant r-1$ if and only if $\bar{a} \equiv (1, 1, \ldots, 1)$, which occurs if and only if $[V] = 0 \in \tilde{R}_{\mathrm{Gl},\,\mathrm{ev}}(\mathbb{Z}_2 \to D_{2r})$.

COROLLARY 2·4. *Let D_{2r} $(r > 1, r$ odd$)$ act freely and topologically on a compact topological manifold M^{2m+1}, so that the action on ∂M is induced from a $\mathbb{Z}_2$-action and so that $i_*\colon H_m(\partial\tilde{M}; \mathbb{F}_2) \to H_m(\tilde{M}; \mathbb{F}_2)$ is injective. Then*

$$\sum_{i=0}^{m} \mathrm{trace}\,(g_*\colon H_i(M; \mathbb{F}_2) \to H_i(M; \mathbb{F}_2)) = 0 \in \mathbb{F}_2.$$

Proof. By 1·7, 2·2, and 2·3 the above sum must be zero with $\mathbb{F}_2$ replaced by $\mathbb{F}_2[\zeta]$. But since trace is invariant under extension of fields it must also be zero with $\mathbb{F}_2$-coefficients.

Proof of Theorem A. Throughout the proof homology with K-coefficients is understood. Suppose D_{2r} acts smoothly on M^{2m+1} so that $\mathbb{Z}_r$ acts freely. We will assume that $H_m(\mathscr{S}) = 0$ and will show

$$\chi_{\frac{1}{2}}(M) = 0 \in \tilde{R}_{\mathrm{Gl},\,\mathrm{ev}}(\mathbb{Z}_2 \to D_{2r}), \qquad (*)$$

which implies by 2·3 that

$$\sum_{i=0}^{m} \mathrm{trace}\,(g_*\colon H_i(M) \to H_i(M)) = 0 \in K$$

and Theorem A follows.

Let $\mathscr{S}^k$ denote the k-dimensional component of the singular set. The boundary of $T(\mathscr{S}^k)$ is an S^{2m-k}-bundle over $\mathscr{S}^k$; from the Gysin sequence it follows that $H_m(\partial T(\mathscr{S}^k)) = 0$ since $H_m(\mathscr{S}^k) \cong H_{k-m}(\mathscr{S}^k) = 0$. By this and 2·1 $M - \mathrm{int}\,T(\mathscr{S})$ is induced, so by 1·7 and 2·2

$$\chi_{\frac{1}{2}}(M - \mathscr{S}) = 0 \in \tilde{R}_{\mathrm{Gl},\,\mathrm{ev}}(\mathbb{Z}_2 \to D_{2r}).$$

By the assumption on $\mathscr{S}$, $H^m(\mathscr{S}) = 0$, so there is an exact sequence

$$0 \to H^{m+1}(M, \mathscr{S}) \to H^{m+1}(M) \to H^{m+1}(\mathscr{S}) \to \ldots \to H^{2m+1}(M) \to 0.$$

Thus

$$\sum_{i=m+1}^{2m+1} (-1)^i [H^i(M, \mathscr{S})] - \sum_{i=m+1}^{2m+1} (-1)^i [H^i(M)] + \sum_{i=m+1}^{2m+1} (-1)^i [H^i(\mathscr{S})] = 0.$$

https://doi.org/10.1017/S0305004100067311 Published online by Cambridge University Press

Since the action on $\mathscr{S}$ is induced from a $\mathbb{Z}_2$-action, the last term vanishes. By Alexander–Poincaré duality the first two terms are equal to $\chi_{\frac{1}{2}}(M-\mathscr{S})$ and $\chi_{\frac{1}{2}}(M)$ respectively. Thus (*) holds and Theorem A follows.

Example. Let D_{2r} act on S^1 as symmetries of the regular n-gon, and act diagonally on the join

$$S^{2m+1} = S^1 * S^1 * \ldots * S^1.$$

The fixed set of an involution T is

$$S^m = S^0 * S^0 * \ldots * S^0.$$

It is easily verified that $M = S^{2m+1} - \operatorname{int} T(\mathscr{S})$ is not induced and $\chi_{\frac{1}{2}}(M; K) \neq 0 \in \tilde{R}_{\mathrm{Gl,\,ev}}(\mathbb{Z}_2 \to D_{2r})$ despite the fact that $\mathfrak{N}_*(\mathbb{Z}_2 \to D_{2r}) = 0$.

Remark. With considerably more difficulty, the hypothesis of a smooth action in Theorem A could be replaced by the hypothesis of a locally smooth action on a topological manifold. One needs a substitute for the $\mathbb{Z}_2$-invariant tubular neighbourhood of $\mathscr{S}(M, \mathbb{Z}_2)$ in Lemma 2·1. This is given by constructing a mapping cylinder neighbourhood of $\mathscr{S}(M, \mathbb{Z}_2)$ in $M/\mathbb{Z}_2$ by using Quinn's end theorem [3] (the point here is that $\tilde{K}_i(\mathbb{Z}[\mathbb{Z}_2]) = 0$ for $i \leqslant 0$, so the obstructions vanish), and then lifting to M to get an equivariant mapping cylinder neighbourhood of $\mathscr{S}(M, \mathbb{Z}_2)$ in M. The map from the boundary of the equivariant mapping cylinder neighbourhood to $\mathscr{S}(M, \mathbb{Z}_2)$ is not necessarily a sphere bundle, but can be shown to have the homotopy type of a spherical fibration. This allows one to use the Gysin sequence as in the proof of Theorem A.

The author would like to thank the Sonderforschungsbereich at Göttingen for their hospitality while this paper was being written and Sylvain Cappell for asking a question which started me thinking in this direction. This research has been partially supported by an NSF postdoctoral fellowship.

REFERENCES

[1] R. LEE. Semicharacteristic classes. *Topology* **12** (1973), 183–199.
[2] D. MONTGOMERY and C. T. YANG. Dihedral group actions I. *General Topology and Modern Analysis* (Academic Press, 1981), 295–304.
[3] F. QUINN. Ends of maps, I. *Ann. of Math.* **110** (1979), 275–331.

https://doi.org/10.1017/S0305004100067311 Published online by Cambridge University Press