Ergod. Th. & Dynam. Sys. (2002), **22**, 187–201
DOI: 10.1017/S0143385702000081
© *2002 Cambridge University Press*
Printed in the United Kingdom

Weak mixing for reparameterized linear flows on the torus

BASSAM R. FAYAD

Université Paris 13, Département de Mathématiques, 99 avenue J. B. Clément, 93430 Villetaneuse, France
(e-mail: fayadb@math.univ-paris13.fr)

(*Received* 3 *March* 2000 *and accepted in revised form* 15 *October* 2000)

Abstract. In this paper, we study the display of weak mixing by reparameterized linear flows on the torus $\mathbb{T}^d$, $d \geq 2$. We show that if the vector of the translation flow is Liouvillian (i.e. well approximated by rationals), then for a residual set of time change functions in the C^∞ topology, the reparameterized flow is weak mixing. If this is not the case, i.e. if the vector of the linear flow is Diophantine, it follows from a result of Kolmogorov on the two torus, and its generalization to any dimension by Herman, that any C^∞ reparameterization of the flow is C^∞ conjugate to a linear flow. More generally, in any given class of differentiability C^r for the time change function ϕ, we give the optimal arithmetical condition on the vector of the translation flow that guarantees the existence of a residual set in the C^r topology of weak mixing reparameterizations. In the real analytic case, the optimal arithmetical condition for the generic display of weak mixing under time change is also given.

As a consequence of our results on reparameterizations of Liouvillian linear flows, we obtain that an aperiodic smooth flow on the two-dimensional torus is in general weak mixing. We also deduce the existence on the torus of analytic diffeomorphisms that are rank one and weak mixing.

1. *Introduction*

1.1. We will denote by $\mathbb{T}^d$ the torus $\mathbb{R}^d/\mathbb{Z}^d$. Assume R_α is a minimal translation on $\mathbb{T}^d$ and consider on $\mathbb{T}^{d+1}$ the *irrational* translation flow:

$$\frac{dx}{dt} = (1, \alpha).$$

This flow is minimal and uniquely ergodic for the Haar measure on $\mathbb{T}^{d+1}$. Given $\phi \in C^r(\mathbb{T}^{d+1}, \mathbb{R}_+^*)$, $r \geq 1$, we define the *reparameterization*, or smooth time change, of this translation flow, with speed $1/\phi$, to be the flow given by

$$\frac{d\theta}{dt} = \frac{\alpha}{\phi(\theta, s)}, \quad \frac{ds}{dt} = \frac{1}{\phi(\theta, s)}.$$

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

The reparameterized flow is strictly ergodic (minimal and uniquely ergodic) and the invariant measure is $\phi(x)\,dx$, where dx denotes the Haar measure on $\mathbb{T}^{d+1}$ (see [**17**]). Considering the section $s = 0$, the reparameterized flow can be viewed as a special flow constructed over the translation R_α and under a suspension function with the same regularity as ϕ. To obtain results on time change for flows it is often more convenient to work with special flows (i.e. suspension flows) that are easier to handle, and then transfer the properties to reparameterizations. The exact definition of special flows will be given in §2 and the natural correspondence with reparameterization will be explained in §3.

Using reparameterizations of irrational flows on the two torus, Shklover, in 1967 [**18**], proved the existence of analytic flows that are weak mixing but not mixing. He actually shows the following result:

> For any real-analytic function on the circle $\mathbb{T}^1 = \mathbb{R}/\mathbb{Z}$, $f > 0$, which is not a polynomial, there exists $\alpha \in \mathbb{R} - \mathbb{Q}$ such that the special flow constructed over the rotation R_α of the circle with a suspension function equal to f is weak mixing.

By the natural correspondence with reparameterizations this yields the existence of analytic reparameterizations of irrational flows on $\mathbb{T}^2$ that are weak mixing. In the work of Shklover the suspension function is explicit, it could be any analytic function that is not a polynomial. Nevertheless, the rotation number α is obtained by going to the limit over rational numbers and the result is still a result of existence. In other words, by his method one cannot obtain the explicit condition on α that implies the existence of an analytic (respectively C^∞ or C^r, $r \in \mathbb{R}_+$) reparameterization of $R_{t(1,\alpha)}$ that is weak mixing. One of our aims here is to give these arithmetic conditions. In fact, given α with some arithmetical condition (Diophantine, Liouville, ...), we will find the optimal class of differentiability, corresponding to this condition, in which there exists a weak mixing reparameterization of $R_{t(1,\alpha)}$. We will actually prove the existence, in this class of differentiability, of a G_δ dense subset of reparameterizations of $R_{t(\alpha,1)}$ that are weak mixing. To obtain such generic dynamics we will use Baire category techniques inspired by some works of Herman and Fathi [**5, 10**]. In their turn, Herman and Fathi [**5**] were inspired by constructions via fast approximations developed in [**1**].

Explicit examples of weak mixing analytic special flows over irrational rotations of the circle were constructed by Katok [**14**]. Starting from these constructions of Katok, Iwanik obtained similar results to our Theorems 3 and 4 ([**12**] and [**13**]) in the case $d = 1$, in his study of smooth cocycles over irrational rotations of the circle.

In fact, our proofs show how to construct explicit examples of weak mixing reparameterizations (cf. Remark 1); and from the nature of weak mixing (cf. Proposition 2) and the fact that adding a trigonometrical polynomial to the time change function does not alter weak mixing, the existence of one weak mixing time change is enough to prove the existence of a residual set of them. Still, we preferred to present a categorical argument that fits perfectly with the nature of weak mixing.

For explicit examples and a direct and geometrical approach to prove mixing features of reparameterized flows, one can also refer to [**6**] where examples of mixing analytic reparameterizations of linear flows are produced on the torus $\mathbb{T}^3$.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

Finally, in our proof, we will insist on having the results in any dimension. Indeed, in a paper on invariant tori by symplectic diffeomorphisms [**9**], Herman proves the remarkable fact that, for a generic C^∞ exact symplectic perturbation of a completely integrable symplectic diffeomorphism, the dynamics on the generic invariant tori† of the perturbation is very different from what one usually expects, meaning different from the C^∞ diffeomorphisms C^∞conjugate to translations of $\mathbb{T}^n$ that we have on the tori given by KAM (Kolmogorov–Arnold–Moser) (also see [**20**] for a summary of this article). An example of such generic behavior that he finds is weak mixing, and he uses in his proof the result of Shklover for invariant tori of dimension $n = 2$, and the examples of Anosov–Katok [**1**] for $n \geq 3$. As remarked by Herman, rather than to use Anosov–Katok constructions it is easier to generalize Shklover's result to any dimension and obtain by the same token analytic examples, which is not yet possible by the methods in [**1**].

1.2. We recall first the definition of weak mixing and we introduce some notation.

A measure preserving flow $\{T^t\}$ on (L, μ) is said to be weak mixing if for all f, $g \in L^2(L, \mu)$ we have

$$\lim_{t\to\infty} \frac{1}{t} \int_0^t \left| \int_L f(T^u x) g(d)\, d\mu - \int_L f\, d\mu \int_L g\, d\mu \right| du = 0. \tag{1}$$

An equivalent definition is that for all measurable sets A and B

$$\mu\Big(T^{-t} A \bigcap B\Big) \to \mu(A)\mu(B), \tag{2}$$

when $|t|$ goes to infinity on a set of density one over $\mathbb{R}$.

One can also prove that a flow $\{T^t\}$ is weak mixing if and only if it does not have an eigenfunction, i.e. a measurable function h, not constant, such that $h(T^t x) = e^{i\lambda t} h(x)$, for some eigenvalue $\lambda \in \mathbb{R}$.

For the equivalence between the definitions, we refer to the book of Parry [**17**].

For $r \in \mathbb{R}^+ \bigcup \{+\infty\}$, we denote by $C^r(\mathbb{T}^d, \mathbb{R})$ the set of real functions on $\mathbb{R}^d$ of class C^r and $\mathbb{Z}^d$-periodic. The set $C^r(\mathbb{T}^d, \mathbb{R})$ is hence a Baire space for the C^r topology. If there is no ambiguity we might just denote these spaces by C^r and we will denote their norms by $\|\cdot\|_r$. Furthermore, if we denote by $C^r(\mathbb{T}^d, \mathbb{R}^*_+)$ the set of strictly positive functions in $C^r(\mathbb{T}^d, \mathbb{R})$, it is an open set for the C^0 norm, and in particular for the C^r norm, therefore it is a Baire space.

For $k = (k_1, \ldots, k_d) \in \mathbb{Z}^d$ and $\alpha = (\alpha_1, \ldots, \alpha_d) \in \mathbb{R}^d$ we will use the following notation:

$$\langle k, \alpha \rangle = \sum_{i=1}^{d} k_i \alpha_i, \qquad \|k\| = \sup_i |k_i|,$$
$$\|\!|\langle k, \alpha \rangle|\!\| = \inf_{p\in\mathbb{Z}} |\langle k, \alpha \rangle + p|.$$

† The generic invariant torus of the perturbation is not a KAM torus but is a C^0 limit of a KAM torus. The restriction on the dynamics of such a tori is that they should be in the closure of the C^∞ diffeomorphisms C^∞ conjugate to translations. That is the reason why Herman investigates, in [**9**], generic properties in the closure of diffeomorphisms C^∞ conjugate to translations. Reparameterized irrational flows are an excellent tool to this end.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

Given a vector $\alpha \in \mathbb{R}^d$, we say it is *β-Diophantine* if there exists a constant $C > 0$ such that for every $k \in \mathbb{Z}^d - \{0\}$ we have

$$|||\langle k, \alpha\rangle||| \geq \frac{C}{\|k\|^{d+\beta}}.$$

In this case we say that α is Diophantine or satisfies a Diophantine condition. If there is a constant $C > 0$ such that α verifies the above condition with $\beta = 0$ we say that α is of constant type. For commodity in stating the results we will allow the notation '0 Diophantine' for such vectors.

We recall the arithmetic decomposition of $\mathbb{R}^d = CD \sqcup L \sqcup \mathbb{Q}$, where $\mathbb{Q}$ designates the vectors with rationally dependent coordinates, CD the Diophantine vectors, and L the Liouville vectors (vectors that do not satisfy any Diophantine condition and are not in $\mathbb{Q}$). Again for commodity in stating the result we will use the notation α 'is not $+\infty$ Diophantine' for Liouville vectors.

Let $d \in \mathbb{N}$, $d \geq 1$. With the above notation, we show the following.

THEOREM 1. *Let $\beta \in \mathbb{R}^+ \bigcup \{+\infty\}$. Assume the irrational vector α in $\mathbb{R}^d$ is not β-Diophantine. Then, for a dense G_δ of ϕ in $C^{\beta+d}(\mathbb{T}^{d+1}, \mathbb{R}_+^*)$, the reparameterization of the translation flow $R_{t(1,\alpha)}$, with speed $1/\phi$ is weak mixing (for the unique invariant measure).*

The result is the best one can have as shown by the following proposition due to Kolmogorov (in dimension 2, i.e. $d = 1$, [**16**]) and generalized to any dimension by Herman in [**11**].

PROPOSITION 1. *If $\alpha \in \mathbb{T}^d$ is β-Diophantine and the function ϕ is C^r for some $r > \beta + d$, the reparameterization of the flow $R_{t(1,\alpha)}$, with speed $1/\phi$ is C^0 conjugate to a translation flow on $\mathbb{T}^{d+1}$.*

For a sketch of the proof to this proposition, cf. the remark on optimality in the next section devoted to special flows.

In the case of C^∞ reparameterizations ($\beta = +\infty$) the dichotomy is therefore complete between weak mixing and conjugation to translation flows and we have the following.

COROLLARY 1. *Let R_α be a minimal translation on $\mathbb{T}^d$. Then either one of two possibilities hold:*

(i) *the vector α is Diophantine and for any $\phi \in C^\infty(\mathbb{T}^{d+1}, \mathbb{R}_+^*)$, the reparameterization of the flow $R_{t(1,\alpha)}$, with speed $1/\phi$ is C^∞ conjugate to a translation flow on $\mathbb{T}^{d+1}$;*

(ii) *the vector α is Liouville and for a dense G_δ of $\phi \in C^\infty(\mathbb{T}^{d+1}, \mathbb{R}_+^*)$, the reparameterization of the flow $R_{t(1,\alpha)}$, with speed $1/\phi$ is weak mixing (for the unique invariant measure).*

For analytic reparameterizations, we have analogous results. First define for $h > 0$ the set $C_h^w(\mathbb{T}^d, \mathbb{R}_+^*)$ of strictly positive real functions on $\mathbb{R}^d$, $\mathbb{Z}^d$-periodic and that can be extended to holomorphic functions on the open band $B_h = \{z \in \mathbb{C}^d : \sup_i |\operatorname{Im} z_i| < h\}$. With the topology of the uniform convergence on compact sets, $C_h^w(\mathbb{T}^d, \mathbb{R}_+^*)$ is a Baire space and we have the following.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

THEOREM 2. *Given an irrational vector $\alpha \in \mathbb{R}^d$, if there exist $\delta > 0$ and a sequence of $k_n \in \mathbb{Z}^d$ such that*

$$|||\langle k_n, \alpha\rangle||| \leq e^{-\delta\|k_n\|},$$

then, for a dense G_δ subset of functions ϕ in the set of $C^w_{\delta/2\pi}$ strictly positive functions, the reparameterization of the translation flow $R_{t(1,\alpha)}$, with speed $1/\phi$ is weak mixing (for the unique invariant measure).

Here, the result is again optimal and we have a perfect dichotomy, because if α satisfies

$$\lim_{\|k\|\to\infty} \frac{-\log |||\langle k, \alpha\rangle|||}{\|k\|} = 0 \tag{3}$$

then, for any real analytic function $\phi > 0$, the reparameterization of $R_{t\alpha}$ by ϕ is analytically conjugate to a translation flow failing therefore to be weak mixing (the proof is based on the linearized equation and is analogous to the proof of Proposition 1, cf. §2).

Remark. The condition that implies conjugacy to a translation flow is weaker than the Brjuno condition.

This paper contains four sections. §§2 and 3 are dedicated to the proof to the Theorems 1 and 2. The essential results are obtained in §2 where we prove analogous theorems for special flows.

As we stated before, it is possible to obtain weak mixing as defined in (2), with a direct and a geometrical method, but it appears easier to prove the equivalent spectral characterization of weak mixing, i.e. the non-existence of eigenfunctions. In Lemma 2, we state a central criterion on the ceiling-function Birkhoff-sums that guarantees the non-existence of eigenfunctions for the special flow. Then we will use Baire category arguments and the stationary phase method to study when this criterion is fulfilled in relation with the arithmetics of α and the regularity of the ceiling function φ. In §3 we derive the results for reparameterizations.

In the last section we show that the general aperiodic flow of class C^∞ on the two-torus is weak mixing.

Finally, we have also added an appendix due to Michael Herman on the possibility of using time t_0 maps of the flows considered above to obtain analytic diffeomorphisms on the torus, combining the weak mixing property with the rank one property.

2. *Weak mixing for special flows*

We recall the definition of a special flow. Given a Lebesgue space L, a measure preserving transformation T on L and an integrable strictly positive real function defined on L we define the special flow over T and under the *ceiling function* φ by inducing on $L \times \mathbb{R}/\sim$, where $\sim$ is the identification $(x, s + \varphi(x)) \sim (T(x), s)$, the action of

$$L \times \mathbb{R} \to L \times \mathbb{R}$$
$$(x, s) \to (x, s + t).$$

If T preserves a unique probability measure μ then the special flow will preserve a unique probability measure that is the normalized product measure of μ on the base and the Lebesgue measure on the fibres.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

In this section α is an irrational vector in $\mathbb{R}^d$, $d \geq 1$, and we consider special flows constructed over the translation R_α of the torus $\mathbb{T}^d$. We will prove the following.

THEOREM 3. *If the vector $\alpha \in \mathbb{R}^d$ is not β-Diophantine then there exists a dense G_δ for the $C^{\beta+d}$ topology, of functions $\varphi \in C^{\beta+d}(\mathbb{T}^d, \mathbb{R}^*_+)$, such that the special flow constructed over R_α with the ceiling function φ is weak mixing.*

In the analytic case we have the following.

THEOREM 4. *If the vector $\alpha \in \mathbb{R}^d$ is such that there exist $\delta > 0$ and a sequence of $k_n \in \mathbb{Z}^d$ satisfying*

$$|||\langle k_n, \alpha\rangle||| \leq e^{-\delta\|k_n\|},$$

then, for a dense G_δ of functions φ in $C^w_{\delta/2\pi}(\mathbb{T}^d, \mathbb{R}^*_+)$, the special flow constructed over R_α with the ceiling function φ is weak mixing.

Proof of optimality. Like the theorems on reparameterization, Theorems 3 and 4 are optimal, and this is easy to check. Indeed, if the cohomological equation

$$\varphi(\theta) - \int_{\mathbb{T}^d} \varphi = \chi(\theta) - \chi(\theta + \alpha), \tag{4}$$

has a continuous solution χ, the special flow $\{R_\alpha, \varphi\}$ is C^0 conjugate to a special flow over R_α with a constant ceiling function, i.e. to a translation flow on $\mathbb{T}^{d+1}$ (cf. [**15**] or [**16**] for a simple proof). But the study of the linearized equation (4) using Fourier expansions (see [**10**, Annexe §8, p. 229]) proves the existence of a continuous solution χ when $\alpha \in \mathbb{T}^d$ is β-Diophantine† and the function φ is C^r with $r > \beta + d$. In the case of ceiling functions of class C^∞, and Diophantine vectors α, (4) always has a C^∞ solution, and the conjugacy of $\{R_\alpha, \varphi\}$ to a standard special flow with a constant ceiling function is of class C^∞. Finally, the condition on the vector α that implies *a priori* the existence of solutions in the real analytic case is the one stated in (3). □

We start the proof of Theorem 3 with a very general fact about weak mixing.

PROPOSITION 2. *Given $\alpha \in \mathbb{R}^d$, in every space $C^r(\mathbb{T}^d, \mathbb{R}^*_+)$, $C^\infty(\mathbb{T}^d, \mathbb{R}^*_+)$ or $C^w_h(\mathbb{T}^d, \mathbb{R}^*_+)$, the set of strictly positive functions φ for which the special flow $\{R_\alpha, \varphi\}$ is weak mixing is a G_δ set.*

Proof. We can find a proof of this result in the introductory book to ergodic theory of Halmos [**7**], in his proof of the second category theorem. □

In light of Proposition 2, to prove Theorem 3, it is enough to show that the set of strictly positive functions φ for which the flow is weak mixing is dense in the set $C^{\beta+d}(\mathbb{T}^d, \mathbb{R}^*_+)$. Still, in our proof of density that is based on a Baire category argument and a sufficient criterion implying weak mixing, we will obtain that the above set contains a G_δ dense set. We then use Proposition 2 to conclude that it is a G_δ dense set.

† The exact result as stated in [**10**] is the following. Assume α is β-Diophantine and let $\theta = d + \beta$ (when $\beta = 0$ take $\theta = d + \epsilon$). Then if $r > \theta$, and $\varphi \in C^r(\mathbb{T}^d)$, there exists $\psi \in C^{r-\theta}(\mathbb{T}^d)$ satisfying the equation (4) if $r - \theta$ is not integer; if $r - \theta$ is integer then the solution ψ is $C^{r-\theta-1}$ and 'smooth in the sense of Zygmund'. In both cases it is continuous.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

First we state a classical general lemma on weak mixing for ergodic special flows, the proof of which can be found in [**2**] for example. In this lemma $\{T^t\}$ will be the special flow constructed from an ergodic automorphism T of a Lebesgue space L and from a function $f > 0$.

LEMMA 1. *The flow $\{T^t\}$ is weak mixing if and only if, for any λ in $\mathbb{R}^*$ the equation*

$$h(T(x)) = e^{i\lambda f(x)}h(x) \tag{5}$$

does not have a non-zero measurable solution h.

Now we return to the special flow constructed over the rotation automorphism R_α of the torus $\mathbb{T}^d$ with the ceiling function φ and from the lemma we just stated we obtain a criterion, on the Birkhoff sums of φ, which guarantees weak mixing for the special flow. We will always denote the Birkhoff sums by

$$\varphi_m(\theta) = \sum_{k=0}^{m-1} \varphi(\theta + k\alpha).$$

LEMMA 2. (Criterion for weak mixing) *If for every g in $L^2(\mathbb{T}^d, \mathbb{C})$, and for every λ in $\mathbb{R}^*$, we have*

$$\inf_{m\in\mathbb{N}} \left| \int_{\mathbb{T}^d} e^{i\lambda\varphi_m(\theta)} g(\theta)\, d\theta \right| = 0,$$

then the flow over R_α with the ceiling function φ is weak mixing.

Proof of Lemma 2. Since R_α is ergodic any measurable solution of (5) has constant modulus and is therefore in L^2. Suppose h is a solution of (5), we have

$$h(\theta + m\alpha) = e^{i\lambda\varphi_m(\theta)}h(\theta)$$

and for any $k \in \mathbb{Z}^d$

$$\begin{aligned} e^{-i2\pi m\langle k,\alpha\rangle} \int_{\mathbb{T}^d} h(\theta)e^{i2\pi\langle k,\theta\rangle}\, d\theta &= \int_{\mathbb{T}^d} h(\theta + m\alpha)e^{i2\pi\langle k,\theta\rangle}\, d\theta, \\ &= \int_{\mathbb{T}^d} e^{i\lambda\varphi_m(\theta)}h(\theta)e^{i2\pi\langle k,\theta\rangle}\, d\theta. \end{aligned}$$

Should the condition of the lemma be satisfied we would have

$$\inf_{m\in\mathbb{N}} \left| \int_{\mathbb{T}^d} e^{i\lambda\varphi_m(\theta)}h(\theta)e^{i2\pi\langle k,\theta\rangle}\, d\theta \right| = 0,$$

which implies that $\int_{\mathbb{T}^d} h(\theta)e^{i2\pi\langle k,\theta\rangle}\, d\theta = 0$ for any $k \in \mathbb{Z}^d$, hence h is zero. □

Remark. It is enough to check the condition of the lemma for a countable set of C^∞ functions χ_j, dense in L^2.

Given the irrational vector $\alpha \in \mathbb{T}^d$ and a positive number, β, that can be zero or infinite define

$$\mathcal{A}(\alpha) = \left\{ \varphi \in C^{\beta+d}(\mathbb{T}^d, \mathbb{R}) : \forall|\lambda| \in \mathbb{R}^*, \forall j \in \mathbb{N}, \inf_{m\in\mathbb{N}} \left| \int_{\mathbb{T}^d} e^{i\lambda\varphi_m(\theta)}\chi_j(\theta)\, d\theta \right| = 0 \right\}.$$

Theorem 3 will then follow from the following.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

PROPOSITION 3. *If α is not β-Diophantine, then $\mathcal{A}(\alpha)$ is a G_δ dense subset of $C^{\beta+d}(\mathbb{T}^d, \mathbb{R})$.*

Indeed, in light of Lemma 2, Proposition 3 implies that the set of strictly positive functions φ for which the flow is weak mixing contains a G_δ dense subset. Together with Proposition 2, this proves Theorem 3.

Proof of Proposition 3. For $j, p, k \in \mathbb{N}^3$, define

$$\mathcal{A}_{(j,p,k)}(\alpha) = \left\{\varphi \in C^{\beta+d}(\mathbb{T}^d, \mathbb{R}) : \forall \lambda \in \left[\frac{1}{p}, p\right], \inf_{m\in\mathbb{N}} \left| \int_{\mathbb{T}^d} e^{i\lambda\varphi_m(\theta)} \chi_j(\theta)\, d\theta \right| < \frac{1}{k}\right\}.$$

We have

$$\mathcal{A}(\alpha) = \bigcap_{j\in\mathbb{N}} \bigcap_{p\in\mathbb{N}} \bigcap_{k\in\mathbb{N}} \mathcal{A}_{(j,p,k)}(\alpha).$$

The set $\mathcal{A}_{(j,p,k)}(\alpha)$ is obviously open (we took $\lambda \in [1/p, p]$ for this purpose), and we just have to prove that it is dense, which will be the concern of the rest of the section.

First we write the fact that α is not β-Diophantine. There exist a sequence $k_n \in \mathbb{Z}^d$ with $\lim_{n\to\infty} \|k_n\| = +\infty$ and a sequence d_n of positive real numbers with $\lim_{n\to\infty} d_n = +\infty$, such that

$$\|\langle k_n, \alpha\rangle\| < \frac{1}{d_n} \frac{1}{\|k_n\|^{d+\beta}}. \tag{6}$$

(If $\beta = +\infty$, i.e. if α is a Liouville vector, we put $\|k_n\|^n$ in (6) instead of $\|k_n\|^{d+\beta}$.)

We introduce next a sequence of real functions on $\mathbb{T}^d$:

$$\psi^{(n)}(\theta) = \frac{\cos(2\pi\langle k_n, \theta\rangle)}{\sqrt{d_n}\, \|k_n\|^{d+\beta}}.$$

(If $\beta = +\infty$, we take n instead of $d + \beta$.)

Finally, we denote by m_n the integer part of $d_n^{3/4}\|k_n\|^{d+\beta}$ (in the following, the crucial fact about m_n will be that $\sqrt{d_n}\, \|k_n\|^{d+\beta} \ll m_n \ll d_n \|k_n\|^{d+\beta} < \|\langle k_n, \alpha\rangle\|^{-1}$).

We will need the following lemmas (the first one is direct from the definition of $\psi^{(n)}$).

LEMMA 3. *If $\|\cdot\|_{\beta+d}$ denotes the $C^{\beta+d}$ norm, then we have*

$$\|\psi^{(n)}\|_{d+\beta} \xrightarrow[n\to\infty]{} 0.$$

(If $\beta = +\infty$, we take n instead of $d + \beta$.)

From our choice of m_n we have the following.

LEMMA 4. *We have*

$$\psi^{(n)}_{m_n}(\theta) = X_n \cos(2\pi\langle k_n, \theta\rangle + \phi_n),$$

where X_n is a real positive number satisfying $X_n \geq \pi^{-1} d_n^{1/4}$.

Proof of Lemma 4. We have

$$\begin{aligned}
\psi^{(n)}_{m_n}(\theta) &= \mathrm{Re}\left(\frac{1 - e^{i2\pi m_n\langle k_n,\alpha\rangle}}{1 - e^{i2\pi\langle k_n,\alpha\rangle}} \frac{e^{i2\pi\langle k_n,\theta\rangle}}{\sqrt{d_n}\, \|k_n\|^{d+\beta}}\right), \\
&= \mathrm{Re}\left(e^{i2\pi(m_n-1)\langle k_n,\alpha\rangle} \frac{\sin(\pi m_n\langle k_n, \alpha\rangle)}{\sin(\pi\langle k_n, \alpha\rangle)} \frac{e^{i2\pi\langle k_n,\theta\rangle}}{\sqrt{d_n}\, \|k_n\|^{d+\beta}}\right), \\
&= X_n \cos(2\pi\langle k_n, \theta\rangle + \phi_n),
\end{aligned}$$

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

if we let

$$X_n = \frac{\sin(\pi m_n \langle k_n, \alpha\rangle)}{\sin(\pi \langle k_n, \alpha\rangle)} \frac{1}{\sqrt{d_n}\,\|k_n\|^{d+\beta}}.$$

Since $m_n \|\!| \langle k_n, \alpha\rangle |\!\| \leq d_n^{-1/4}$, we have

$$\frac{\sin(\pi m_n \langle k_n, \alpha\rangle)}{\sin(\pi \langle k_n, \alpha\rangle)} \geq \frac{2}{\pi} m_n,$$

hence

$$X_n \geq \frac{2}{\pi} \frac{m_n}{\sqrt{d_n}\,\|k_n\|^{d+\beta}} \geq \frac{1}{\pi} d_n^{1/4}. \qquad \square$$

Comment. The functions $\psi^{(n)}$ we introduced are 'α' periodic (in the sense that $\psi^{(n)}(x + j\alpha) \sim \psi^{(n)}(x)$) as long as the number of iterations of R_α is such that $j \ll \|\!| \langle k_n, \alpha\rangle |\!\|^{-1}$. Consequently, the Birkhoff sums of $\psi^{(n)}$ up to m_n will look like $m_n \psi^{(n)}$. If m_n is moreover, as it is the case here, such that $m_n D_n^{-1} \to \infty$, where D_n is the denominator in the expression of $\psi^{(n)}$, then $\psi^{(n)}_{m_n}$ will have large oscillations. This essential phenomenon will allow us to check the criterion of Lemma 2 by means of the stationary phase method. The idea is to perturb a given function by adding to it a function $\psi^{(n)}$ that will 'produce mixing' at time m_n. It becomes clear, now, that the less α is Diophantine, the larger we can take D_n, and the smaller (i.e. more differentiable) will be the perturbation $\psi^{(n)}$ that we have to add to the given ceiling function to produce some mixing features for the special flow.

From the estimation in Lemma 4 and a basic fact on the stationary phase, we can derive the following fundamental result.

LEMMA 5. *For every $g \in C^\infty(\mathbb{T}, \mathbb{R})$, and for every $\lambda \neq 0$, we have, as n goes to infinity,*

$$\int_{\mathbb{T}^d} e^{i\lambda \psi^{(n)}_{m_n}(\theta)} g(\theta)\, d\theta = O\left(\frac{1}{\sqrt{|\lambda| X_n}}\right).$$

Proof of Lemma 5. From a lemma on a stationary phase† that can be found in [**4**, Chapter IV, paragraph (4.8)], we know that for every g in $C^\infty(\mathbb{T}, \mathbb{R})$ we have for large X_n

$$\int_{\mathbb{T}^1} e^{i\lambda X_n \cos 2\pi x} g(x)\, dx \leq \frac{C(g)}{\sqrt{|\lambda| X_n}}$$

where the constant $C(g)$ only depends (as n goes to infinity) on the C^0 norm of g, $\|g\|_0$. In the expression of $\psi^{(n)}$, we can assume that the first coordinate k_n^1 of the vector k_n is different from zero. From the inequality above we have when n is large enough

$$\int_{\mathbb{T}^1} e^{i\lambda X_n \cos(2\pi k_n^1 \theta_1 + u_n)} g(\theta_1, \tilde{\theta})\, d\theta_1 \leq \frac{C(g)}{\sqrt{|\lambda| X_n}},$$

† The exact statement of Dieudonné is the following. Assume g and h are two functions in $C^\infty([a, b], \mathbb{R})$, such that h' has only one zero $c \in]a, b[$ with $g(c) \neq 0$ and $h''(c) \neq 0$. Then for any t large enough, if $h''(c) > 0$

$$\int_a^b g(x) e^{ith(x)}\, dx = \left(\frac{\pi}{2th''(c)}\right)^{1/2} g(c) e^{ith(c) + i\pi/4} + O\left(\frac{1}{t}\right).$$

If $h''(c) < 0$ the same is true if we change $i\pi/4$ to $-i\pi/4$ and $h''(c)$ to $-h''(c)$ in the above expression. If $g(c) = 0$ the same equality holds.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

where $\tilde{\theta} = (\theta_2, \ldots, \theta_d)$ and $u_n = 2\pi(\langle k_n, \theta\rangle - k_n^1\theta_1) + \phi_n$. Integrating along the other variables we obtain the result. □

Remark. If g is a C^∞ function, and g_n is a sequence converging to g in norm L^1, then we have from the above lemma

$$\lim_{n\to\infty} \int_{\mathbb{T}^d} e^{i\lambda\psi_{m_n}^{(n)}(\theta)} g_n(\theta)\, d\theta \xrightarrow[n\to\infty]{} 0.$$

Now, we are ready to prove that $\mathcal{A}_{(j,p,k)}(\alpha)$ is a residual set in $C^{\beta+d}(\mathbb{T}^d, \mathbb{R})$.

Fix a function $\varphi \in C^{\beta+d}(\mathbb{T}^d, \mathbb{R})$ and take $\epsilon > 0$ arbitrarily small. We can assume the integral of φ is zero and arbitrarily close to it find a C^∞ coboundary, i.e. a function of the type $\hat{\varphi} = \eta \circ R_\alpha - \eta$, where η is of class C^∞ such that

$$\|\varphi - \hat{\varphi}\|_{d+\beta} < \epsilon.$$

Indeed, any trigonometric polynomial with zero integral is a coboundary. Then, for $n \in \mathbb{N}$ we define

$$\hat{\varphi}^{(n)} = \hat{\varphi} + \psi^{(n)}.$$

Now, if we take n large enough, we will have, from Lemma 3:

$$\|\varphi - \hat{\varphi}^{(n)}\|_{d+\beta} < 2\epsilon. \tag{7}$$

On the other hand, considering a subsequence if necessary, we can assume that $\hat{\varphi}_{m_n}$ converges to a fixed cocycle $\eta \circ R_{\tilde{\alpha}} - \eta$, then, applying Lemma 5 and the remark that follows, we have

$$\int_{\mathbb{T}^d} e^{i\lambda\hat{\varphi}_{m_n}^{(n)}(\theta)} \chi_j(\theta)\, d\theta = \int_{\mathbb{T}^d} e^{i\lambda\hat{\varphi}_{m_n}^{(n)}(\theta)} e^{i\lambda\hat{\varphi}_{m_n}^{(n)}(\theta)} \chi_j(\theta)\, d\theta \xrightarrow[n\to\infty]{} 0.$$

(We used Lemma 5 for $g(\theta) = e^{i\lambda(\eta\circ R_{\tilde{\alpha}}(\theta) - \eta(\theta))}\chi_j(\theta)$, and the fact that $e^{i\lambda\hat{\varphi}_{m_n}(\theta)}\chi_j(\theta) \to e^{i\lambda(\eta\circ R_{\tilde{\alpha}}(\theta)-\eta(\theta))}\chi_j(\theta)$ uniformly when n goes to infinity.)

So when n is large enough, $\hat{\varphi}^{(n)}$ is in $\mathcal{A}_{(j,p,k)}(\alpha)$. The real number ϵ being arbitrarily small this completes the proof of density of $\mathcal{A}_{(j,p,k)}(\alpha)$. For the C^∞ case, i.e. β is infinite, we take $\varphi \in C^\infty$ and for an arbitrary $l \in \mathbb{N}$, with the same arguments used above, we find a function C^l close to φ and in $\mathcal{A}_{(j,p,k)}(\alpha)$. Proposition 3 is proved. □

Remark 1. In Lemma 5, we could establish a uniform bound on the integral that only depends on the C^1 norm of g. This would enable us to check the validity of Lemma 2 to the function $\psi = \sum \psi^{(n)}$ and obtain thus an explicit ceiling function under which the special flow is weak mixing.

Before we close this section, we want to show how the proof applies to the analytic case.

Proof of Theorem 4. Take for $\psi^{(n)}$ the functions

$$\psi^{(n)}(\theta) = \mathrm{Re}\left(\frac{e^{i2\pi\langle k_n,\theta\rangle}}{n^{-2}e^{\delta\|k_n\|}}\right).$$

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

We have

$$\|\psi^{(n)}\| \xrightarrow[n\to\infty]{} 0,$$

for the norm on $C^w_{\delta/2\pi}$ of uniform convergence on compact sets in the open band $B_{\delta/2\pi}$.

Now, for m_n we take the integer part of $n^{-1}e^{\delta\|k_n\|}$, which obviously satisfies

$$m_n \|\!| \langle k_n, \alpha\rangle |\!\| \xrightarrow[n\to\infty]{} 0,$$

while

$$\frac{m_n}{n^{-2}e^{\delta\|k_n\|}} \xrightarrow[n\to\infty]{} \infty.$$

This leads to the proof, as in Theorem 3. □

3. *Reparameterization of translation flows*

In this section we consider reparameterizations of irrational flows, with speed $1/\phi$:

$$\frac{d\theta}{dt} = \frac{\alpha}{\phi(\theta, s)}, \quad \frac{ds}{dt} = \frac{1}{\phi(\theta, s)},$$

where $\phi > 0$ is of class at least C^1, R_α is a minimal translation on $\mathbb{T}^d$; $\theta \in \mathbb{T}^d$ and $s \in \mathbb{R}$.

Considering a Poincaré section, this flow can be viewed as a special flow constructed from R_α on $\mathbb{T}^d$ and from the ceiling function given by

$$R(\phi)(\theta) = \varphi(\theta) = \int_0^1 \phi(\theta + \xi\alpha, \xi)\, d\xi. \tag{8}$$

Remark. If we consider the section $(\theta, 0)$, this formula just translates the fact that the return time to that section is the integral of one over the speed along the fibres.

From this correspondence between reparameterizations and special flow, we can give a proof to our main theorems using the results of §2.

Proof of Theorem 1. Fix the vector α not β-Diophantine and consider the irrational flow $R_{t(1,\alpha)}$. It is a classical fact, as in Proposition 2 of §2, that the set of $\phi > 0$ for which the reparameterization of $R_{t(1,\alpha)}$ with speed $1/\phi$ is weak mixing is a G_δ, in particular for the topology $C^{\beta+d}$. We want to prove density. We introduce

$$\mathcal{B}(\alpha) = \{\phi \in C^{\beta+d}(\mathbb{T}^{d+1}, \mathbb{R}^*_+) : R(\phi) \in \mathcal{A}(\alpha)\},$$

and

$$\mathcal{B}_{(j,p,k)}(\alpha) = \{\phi \in C^{\beta+d}(\mathbb{T}^{d+1}, \mathbb{R}^*_+) : R(\phi) \in \mathcal{A}_{(j,p,k)}(\alpha)\}.$$

The function $R(\phi)$ being the ceiling function obtained from ϕ by (8).

As in the proof of Theorem 3, we need only to prove that $\mathcal{B}_{(j,p,k)}(\alpha)$ is $C^{\beta+d}$ dense in $C^{\beta+d}(\mathbb{T}^{d+1}, \mathbb{R}^*_+)$. We fix ϕ and we take $\hat{\phi}$ a trigonometrical polynomial close to ϕ. Obviously, $\hat{\varphi} = R(\hat{\phi})$ (given by (8)) will be a trigonometrical polynomial and, by the proof of Theorem 2, for n sufficiently large, $\hat{\varphi} + \psi^{(n)} \in \mathcal{A}_{(j,p,k)}(\alpha)$. Remember that

$$\psi^{(n)}(\theta) = \operatorname{Re}\left(\frac{e^{i2\pi\langle k_n, \theta\rangle}}{\sqrt{d_n}\,\|k_n\|^{d+\beta}}\right).$$

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

Also define

$$\Psi^{(n)}(\theta, s) = \frac{i2\pi(\langle k_n, \alpha\rangle + l_n)}{e^{i2\pi(\langle k_n,\alpha\rangle + l_n)} - 1} \operatorname{Re}\left(\frac{e^{i2\pi\langle k_n,\theta\rangle} e^{i2\pi l_n s}}{\sqrt{d_n}\,\|k_n\|^{d+\beta}}\right),$$

where l_n is chosen to be the closest integer to $-\langle k_n, \alpha\rangle$. A straightforward computation implies

$$R(\Psi^{(n)}) = \psi^{(n)}.$$

We still have to check one lemma.

LEMMA 6. *If $\|\cdot\|_{d+\beta}$ designates the $C^{\beta+d}$ norm, one has*

$$\|\Psi^{(n)}\|_{d+\beta} \xrightarrow[n\to\infty]{} 0.$$

Proof of Lemma 6. The choice of l_n such that $|\langle k_n, \alpha\rangle + l_n| < \frac{1}{2}$ implies

$$\left|\frac{i2\pi(\langle k_n, \alpha\rangle + l_n)}{e^{i2\pi(\langle k_n,\alpha\rangle + l_n)} - 1}\right| < \frac{\pi}{2}.$$

Since $|l_n| \le \|k_n\|(\sum |\alpha_j|) + 1$, $\Psi^{(n)}$, just like $\psi^{(n)}$, goes to 0 in the $C^{d+\beta}$ topology when n goes to infinity. □

Now, because the correspondence in (8), $\phi \to R(\phi)$, is linear we will have $R(\hat{\phi} + \Psi^{(n)}) = \hat{\varphi} + \psi^{(n)} \in \mathcal{A}_{(j,p,k)}(\alpha)$, or equivalently $\hat{\phi} + \Psi^{(n)} \in \mathcal{B}_{(j,p,k)}(\alpha)$. By Lemma 6 and the choice of $\hat{\phi}$ this last function is close to ϕ, which proves density of $\mathcal{B}_{(j,p,k)}(\alpha)$. Theorem 1 on reparameterization is thus proved (β finite). For the C^∞ and analytic case we make the same modifications as in the preceding section in our choice of the $\Psi^{(n)}$'s. □

4. *Aperiodic flows on the two torus*

In this section we consider smooth flows on the two torus given by the system of differential equations

$$\frac{du}{dt} = A(u, v), \quad \frac{dv}{dt} = B(u, v), \tag{9}$$

where A and B are of class C^∞.

When the flow in (9) has no singularities and no closed orbits we will say it is *aperiodic* (although this is not the standard definition of aperiodic flows). We can define the corresponding set of smooth vector fields:

$$\mathcal{E} = \{(A, B) \in C^\infty(\mathbb{T}^2, \mathbb{R}) \times C^\infty(\mathbb{T}^2, \mathbb{R}) \text{ such that the flow in (9) is aperiodic.}\}$$

It follows from standard arguments that the set $\mathcal{E}$ is a G_δ in $C^\infty(\mathbb{T}^2, \mathbb{R}) \times C^\infty(\mathbb{T}^2, \mathbb{R})$, therefore it is a Baire space for the C^∞ topology, and it makes sense to state a 'general', or more precisely a generic property, on it with regard to the Baire category. The theorem we want to prove is the following.

THEOREM 5. *For a dense G_δ of (A, B) in $\mathcal{E}$, the flow given by (9) is weak mixing for its unique invariant measure.*

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

Proof of Theorem 5. Unique ergodicity. For aperiodic flows, a result of Kneser and Siegel (see the paper of Siegel [**19**]) states that there exists on the torus, a closed non-self-intersecting curve Γ of class C^∞ transversal to the flow such that any point of Γ comes back to Γ in a finite time and all trajectories of the flow intersect Γ. Therefore the flow can be viewed as a special flow over Γ. It follows from the smoothness of Γ and of the flow that the return map T on the basis and the return time function f (the ceiling function) associated with this special flow are of class C^∞. We call α the rotation number of T on Γ. By our assumption on the flow, α has to be irrational, therefore T is uniquely ergodic and so is the flow.

Weak mixing. Reasoning as in the previous sections we only need to prove that the couples (A, B) for which the flow (9) is weak mixing are dense in $\mathcal{E}$; the G_δ property being obtained from standard remarks on weak mixing. Pick $(A_0, B_0) \in \mathcal{E}$, and let Γ_0 be a Siegel curve for the corresponding flow that we view as a special flow over Γ_0, denoted by $\{T_0, f_0\}$. To this end, we will need to consider special flows above Γ_0 built with a diffeomorphism T and a ceiling function f that are C^∞ perturbations of T_0 and f_0. These special flows are clearly flows on the torus that are C^∞ close to the special flow $\{T_0, f_0\}$, i.e. to the flow given by (A_0, B_0). Therefore the flows $\{T, f\}$ correspond to vector fields (A, B) that are C^∞ perturbations of (A_0, B_0).

We are ready now to prove density of the weak mixing flows. First of all we perturb T_0 in the C^∞ topology into a diffeomorphism T on Γ_0 that has a Diophantine rotation number α. By Herman's theorem, the map T_0 is C^∞-conjugated to R_α. After conjugacy, we obtain a special flow over R_α with a smooth ceiling function. We can then perturb R_α into a Liouvillian rotation and, finally, use Theorem 3 to obtain a weak mixing flow after an arbitrarily small smooth perturbation of the ceiling function. □

Appendix. A remark on the existence of rank one and weak mixing analytic diffeomorphisms of the torus

Let (X, T, μ) be a dynamical system. We say that T is of *rank one* if there is a sequence of Rokhlin towers whose levels generate the σ-algebra of all measurable sets. For instance, any minimal translation on the torus is of rank one [**3**]. Later on, we will be using this fact.

It is known from [**1**] that weak mixing rank one transformations are abundant in the neighborhood of translations in the C^∞ topology. What about the analytic case? We state here the following fact communicated to us by Herman [**8**].

THEOREM A.1. *Assume φ is a given real analytic strictly positive function on the circle that is not a polynomial. Fix $t_0 \in \mathbb{R}$ such that $t_0 / \int_{\mathbb{T}} \varphi(x)\,dx \in \mathbb{R} - \mathbb{Q}$. Then, there exists a G_δ dense subset of $\mathbb{R}$, $\mathcal{W}_\varphi$, such that, for any $\alpha \in \mathcal{W}_\varphi$, the time map T^{t_0} of the special flow constructed over R_α with the ceiling function φ is of rank one and weak mixing.*

Proof of Theorem A.1. The function φ is fixed. Let $W_\varphi \subset \mathbb{R}$ be the set of numbers α for which the time T^{t_0} of the special flow constructed over R_α and under the ceiling function φ is weak mixing. We claim that W_φ is a dense G_δ of $\mathbb{R}$. As in the previous sections, we only need to prove that W_φ is dense. By the result of Shklover [**18**], we know that since φ is not a polynomial, as close as we want from any α there is a real number $\tilde{\alpha}$ such that

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

the special flow constructed over $R_{\tilde{\alpha}}$ with the ceiling function φ is weak mixing. But then, any time t_0 map of this flow is weak mixing (we use for this the spectral characterization of weak mixing and the fact that the maximal spectral measure of a time t_0 map of a flow is continuous if and only if the maximal spectral measure of the flow is continuous). The density of W_φ is thus proved.

On the other hand we denote by V_φ the set of numbers α for which the time T^{t_0} of the special flow constructed over R_α and under the ceiling function φ is of rank one. We claim that V_φ is a dense G_δ of $\mathbb{R}$. Here, again, it is easy to see that V_φ is a G_δ in $\mathbb{R}$. For this, one uses the definition of the rank one property for a dynamical system (X, T, μ): let $\{\mathcal{P}_i\}_{i\in\mathbb{N}}$ be a generating sequence of finite measurable partitions. The transformation T is of rank one if and only if for any $i \in \mathbb{N}$, and for every $n \in \mathbb{N}$, there exists a Rokhlin tower for T that refines $\mathcal{P}_i$ up to $1/n$; that is, if any atom of $\mathcal{P}_i$ is up to a precision $1/n$ a union of levels of the tower (when we say up to $1/n$ we mean that the measure of the symmetrical difference is smaller than $1/n$). For a fixed partition $\mathcal{P}_i$ and a fixed precision $1/n$, the property of $(1/n)$-refining $\mathcal{P}_i$ is clearly an open condition on T. So if we denote by $V_{\varphi,i,n}$ the set of α such that the time T^{t_0} of the special flow constructed over R_α and under the ceiling function φ satisfies the above condition for the integers i and n; $V_{\varphi,i,n}$ will be open in $\mathbb{R}$. Hence, $V_\varphi = \bigcap_n \bigcap_i V_{\varphi,i,n}$ is a G_δ. We still need to show it is dense. However, by the result of Kolmogorov, we know in particular that if α is Diophantine, then the special flow over R_α with the ceiling function φ is analytically conjugate to the minimal translation flow on the torus of vector $(\alpha/\int_{\mathbb{T}}^1 \varphi(x)\,dx, 1/\int_{\mathbb{T}}^1 \varphi(x)\,dx)$. From now on we assume α is Diophantine so that this conjugacy holds. Because the rank one property is invariant by conjugacy, the time t_0 map of the special flow will be of rank one if the time t_0 map of the translation flow, $R_{t_0(\alpha/\int\varphi, 1/\int\varphi)}$, is of rank one. As we mentioned above, this will be true when the latter translation is minimal. With the assumption on t_0 made in the hypothesis of the theorem, we will just have to discard the countable set of α consisting of

$$\frac{\int_{\mathbb{T}} \varphi(x)\,dx}{t_0}\mathbb{Q} + \mathbb{Q}.$$

Of course, we will still remain with a dense set of Diophantine numbers α. Hence, we proved that V_φ is a G_δ dense in $\mathbb{R}$.

The subset of $\mathbb{R}$, $\mathcal{W}_\varphi$, intersection of V_φ and W_φ, both G_δ dense, is a G_δ dense.

Acknowledgements. I wish to thank Michael Herman, for supervising this work, and Patrice Le Calvez and Raphaël Krikorian, for useful comments and advice. I also want to thank the referee for interesting historical comments and a critical review of the manuscript.

References

[1] D. V. Anosov and A. B. Katok. New examples of ergodic diffeomorphisms of smooth manifolds. *Uspehi Mat. Nauk* **25** (1970), 173–174.

[2] I. P. Cornfeld, S. V. Fomin and Ya. G. Sinai. *Ergodic Theory*. Springer, New York, 1982, ch. 16.

[3] A. del Junco. Transformations with discrete spectrum are stacking transformations. *Canad. J. Math.* **28** (1976), 836–839.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press

[4] J. Dieudonné. *Calcul infintésimal*. Hermann, Paris, 1968.
[5] A. Fathi and M. R. Herman. Existence de difféomorphismes minimaux. *Dynamical Systems*, vol. I.—Warsaw. *Astérisque* **49** (1977), 37–59.
[6] B. R. Fayad. Analytic mixing reparametrizations for irrational flows on the torus $\mathbb{T}^n$, $n \geq 3$. *Ergod. Th. & Dynam. Sys.* **22** (2002), to appear.
[7] P. R. Halmos. *Lectures on Ergodic Theory*. Chelsea, New York, 1960.
[8] M. R. Herman. Personal communication, 1999.
[9] M. R. Herman. On the dynamics of Lagrangian tori invariant by symplectic diffeomorphisms. *Progress in Variational Methods in Hamiltonian Systems and Elliptic Equations*. Longman, Harlow, 1990, pp. 92–112.
[10] M. R. Herman. Sur la conjugaison différentiable des difféomorphismes du cercle à des rotations. *Inst. Hautes Études Sci. Publ. Math.* **49** (1979), 5–233.
[11] M. R. Herman. Exemples de flots hamiltoniens dont aucune perturbation en topologie C^∞ n'a d'orbites périodiques sur un ouvert de surfaces d'énergies. *C. R. Acad. Sci. Paris* **312** (1991), 989–994.
[12] A. Iwanik. Generic smooth cocycles of degree zero over irrational rotations. *Studia Math.* **115** (1995), 241–250.
[13] A. Iwanik. Cyclic approximation of analytic cocycles over irrational rotations. *Colloq. Math.* **70** (1996), 73–78.
[14] A. B. Katok. Constructions in ergodic theory. Unpublished lecture notes.
[15] A. Katok and B. Hasselblatt. *Introduction to the Modern Theory of Dynamical Systems*. Cambridge University Press, Cambridge, 1995, ch. 4.
[16] A. N. Kolmogorov. On dynamical systems with an integral invariant on the torus. *Doklady Akad. Nauk SSSR* **93** (1953), 763–766.
[17] W. Parry. *Topics in Ergodic Theory*. Cambridge University Press, Cambridge, 1981.
[18] M. D. Shklover. On dynamical systems on the torus with continuous spectrum. *Izv. Vuzov* **10** (1967), 113–124.
[19] C. L. Siegel. Note on the differential equations on the torus. *Ann. Math.* **46** (1945), 423–428.
[20] J.-C. Yoccoz. Travaux de Herman sur les tores invariants. *Astérisque, Séminaire Bourbaki* **206** (1992), 332–344.

https://doi.org/10.1017/S0143385702000081 Published online by Cambridge University Press