Canad. J. Math. 2025, pp. 1–24
http://dx.doi.org/10.4153/S0008414X25101739
© The Author(s), 2025. Published by Cambridge University Press on behalf of Canadian Mathematical Society. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

On the Hilbert space derived from the Weil distribution

Masatoshi Suzuki

Abstract. We study the Hilbert space obtained by completing the space of all smooth and compactly supported functions on the real line with respect to the Hermitian form arising from the Weil distribution under the Riemann hypothesis. It turns out that this Hilbert space is isomorphic to a de Branges space by a composition of the Fourier transform and a simple map. This result is applied to state new equivalence conditions for the Riemann hypothesis in a series of equalities.

1 Introduction

The Weil distribution is a distribution associated with the Riemann zeta-function $\zeta(s)$. Let

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma\left(\frac{s}{2}\right)\zeta(s)$$

be the Riemann xi-function, where $\Gamma(s)$ is the gamma-function. Let Γ be the set of all zeros of $\xi(1/2 - iz)$ without multiplicity and let m_γ denote the multiplicity of $\gamma \in \Gamma$. The Riemann hypothesis (RH, for short) claims that all nontrivial zeros of $\zeta(s)$ lie on the critical line $\Re(s) = 1/2$. It is equivalent to the assertion that all $\gamma \in \Gamma$ are real.

The Weil distribution is the linear functional $W : C_c^\infty(\mathbb{R}) \to \mathbb{C}$ defined by

$$C_c^\infty(\mathbb{R}) \ni \psi \mapsto W(\psi) := \sum_{\gamma\in\Gamma} m_\gamma \widehat{\psi}(-\gamma),$$

where $C_c^\infty(\mathbb{R})$ is the space of all smooth and compactly supported functions on $\mathbb{R}$ and

$$\widehat{\psi}(z) := (\mathsf{F}\psi)(z) := \int_{-\infty}^{\infty} \psi(x)\, e^{izx}\, dx \tag{1.1}$$

is the Fourier transform. We omit the description of the topology of $C_c^\infty(\mathbb{R})$, since we do not need it later. Weil [17] (see also the note in [14, Section 3.2]) discovered that the RH is true if and only if the Weil distribution W is nonnegative definite, that is,

$$W(\psi * \widetilde{\psi}) \geq 0 \quad \text{for every } \psi \in C_c^\infty(\mathbb{R}),$$

Received by the editors September 26, 2024; revised October 22, 2025; accepted October 27, 2025.
Published online on Cambridge Core November 3, 2025.
This work was supported by JSPS KAKENHI Grant no. JP23K03050. This work was also supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.
AMS subject classification: **11M26**, 42A82, 46E22.
Keywords: Riemann zeta-function, Riemann hypothesis, Weil distribution, de Branges spaces, screw function.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

where

$$(\phi * \psi)(x) := \int_{-\infty}^{\infty} \phi(y)\psi(x-y)\,dy \quad \text{and} \quad \widetilde{\psi}(x) := \overline{\psi(-x)}.$$

Further, if the RH is true, the Weil distribution is positive definite, that is, $W(\psi * \widetilde{\psi}) > 0$ for every nonzero $\psi \in C_c^\infty(\mathbb{R})$.

Using the Weil distribution, we define the Hermitian form $\langle \cdot, \cdot \rangle_W$ on $C_c^\infty(\mathbb{R})$ by

$$\langle \psi_1, \psi_2 \rangle_W = W(\psi_1 * \widetilde{\psi_2}) = \sum_{\gamma \in \Gamma} m_\gamma \widehat{\psi_1}(-\gamma)(\widehat{\psi_2})^\sharp(-\gamma), \quad \psi_1, \psi_2 \in C_c^\infty(\mathbb{R}), \tag{1.2}$$

where

$$F^\sharp(z) := \overline{F(\bar{z})}$$

for complex-valued functions of a complex variable. We often use this $\sharp$ notation. We call this Hermitian form the *Weil Hermitian form*. Yoshida [19] has studied the Weil Hermitian form in detail by restricting it to a function space on a finite interval $[-a, a]$ $(a > 0)$. The subject of the present article is the behavior of the Weil Hermitian form on the whole line $\mathbb{R}$. Yoshida proposed a method to complete a function space on a finite interval with respect to the Weil Hermitian form without assuming the RH, but it does not extend to the whole line.

Suppose that the RH is true. Then, the Weil Hermitian form $\langle \cdot, \cdot \rangle_W$ is positive definite on $C_c^\infty(\mathbb{R})$. Therefore, the completion $\mathcal{H}_W$ of the pre-Hilbert space $C_c^\infty(\mathbb{R})$ with respect to $\langle \cdot, \cdot \rangle_W$ is defined. The first main result is an explicit description of the Hilbert space $\mathcal{H}_W$. The elements of $\mathcal{H}_W$ are equivalence classes of Cauchy sequences with respect to $\langle \cdot, \cdot \rangle_W$, where two Cauchy sequences are equivalent if their difference converges to zero with respect to $\langle \cdot, \cdot \rangle_W$. The representative of each class can be chosen from $L^2(\mathbb{R})$ (Theorem 5.5 below). Such a result is expected from Lemmas 2 and 3 in [19]. Therefore, we denote the class represented by $\psi \in L^2(\mathbb{R})$ as $[\psi]$ and often identify ψ with $[\psi]$.

For the concrete description of $\mathcal{H}_W$, we use a de Branges space and a model space. The entire function E_ξ defined by

$$E_\xi(z) := \xi(1/2 - iz) + \xi'(1/2 - iz) \tag{1.3}$$

belongs to the Hermite–Biehler class under the RH [8, Theorem 1] and hence it defines the de Branges space $\mathcal{H}(E_\xi)$, where the dash on the right-hand side of (1.3) means differentiation of $\xi(s)$ with respect to s. Furthermore, the meromorphic function

$$\Theta_\xi(z) := E_\xi^\sharp(z)/E_\xi(z) \tag{1.4}$$

in $\mathbb{C}$ is a meromorphic inner function in the upper-half plane $\mathbb{C}_+ = \{z \,|\, \Im(z) > 0\}$ under the RH, and therefore it defines the model space $\mathcal{K}(\Theta_\xi)$. These two Hilbert spaces $\mathcal{H}(E_\xi)$ and $\mathcal{K}(\Theta_\xi)$ are isomorphic with $\|E_\xi F\|_{\mathcal{H}(E_\xi)} = \|F\|_{\mathcal{K}(\Theta_\xi)}$ for every $F \in \mathcal{K}(\Theta_\xi)$ (see Section 2 for details on the Hermite–Biehler class, de Branges spaces, and model spaces). Then, the first result is stated as follows.

Theorem 1.1 *Assume that the RH holds. Let $\mathcal{H}_W$, $\mathcal{H}(E_\xi)$, and $\mathcal{K}(\Theta_\xi)$ be Hilbert spaces as above. Then, the map $\mathcal{K}(\Theta_\xi) \to \mathcal{H}_W$ defined by*

$$F \mapsto [\psi_F], \quad \psi_F := \mathsf{F}^{-1}(F)$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

is an isomorphism between Hilbert spaces and satisfies

$$\|E_\xi F\|^2_{\mathcal{H}(E_\xi)} = \|F\|^2_{\mathcal{K}(\Theta_\xi)} = \pi\langle \psi_F, \psi_F\rangle_W = \pi\langle [\psi_F], [\psi_F]\rangle_W$$

for $F \in \mathcal{K}(\Theta_\xi)$*, where* F^{-1} *is the Fourier inversion on* $L^2(\mathbb{R})$.

This result is proved in Section 5. Note that Theorem 1.1 provides an isomorphism as a Hilbert space, not as a reproducing kernel Hilbert space. The space $\mathcal{H}_W$ is a space of equivalence classes of functions, not a space of functions.

Lagarias suggested after Theorem 1 of [8] that the norm of the de Branges space $\mathcal{H}(E_\xi)$ and the Weil Hermitian form (the spectral side of the "explicit formula" of prime number theory) are similar. Theorem 1.1 shows that they are naturally coincident. Hence, $\mathcal{H}_W$ and $\mathcal{H}(E_\xi)$ must have an "arithmetic structure" through the Weil explicit formula (3.3) below, but we will not discuss this further.

Connes, Consani, and Moscovici [4, Section 4.8] also describe the relation between the theory of de Branges spaces and the Weil Hermitian form, but their de Branges spaces $\mathcal{B}^S_\lambda$ and $\mathcal{H}(E_\xi)$ have completely different properties. Due to the difference in the generators of the de Branges spaces, they are not isomorphic, and a more obvious difference is that they have different spectral properties (see the second half of Section 6).

One of the remarkable properties of de Branges spaces is the structure of subspaces. The set of all de Branges subspaces of a given de Branges space is totally ordered by set-theoretical inclusion (see [18, pp. 500–506] for details). Such a structure also comes to $\mathcal{H}_W$ through the isomorphism of Theorem 1.1 as stated in Theorem 5.7 below.

Another notable property of de Branges spaces is the explicit description of the family of self-adjoint extensions of the multiplication operator by an independent variable $F(z) \mapsto zF(z)$. It enables us to interpret the set of zeros Γ as the set of eigenvalues of a self-adjoint operator on $\mathcal{H}_W$. This means that one of the Hilbert–Pólya spaces is the Hilbert space $\mathcal{H}_W$ naturally obtained from the Weil distribution (see Sections 2.3 and 6 for details).

As stated in Theorem 1.1, the Hilbert space $\mathcal{H}_W$ is isomorphic to a de Branges space under the RH. Moreover, representatives of classes in $\mathcal{H}_W$ can be chosen from the concrete subspace $V(0)$ of $L^2(\mathbb{R})$ defined in (5.2) below. It is surprising that such an explicit description of $\mathcal{H}_W$ is possible, and interesting in itself. However, it is a matter of concern that it is not even possible to define $\mathcal{H}_W$, $\mathcal{H}(E_\xi)$, and $\mathcal{K}(\Theta_\xi)$ without assuming the RH. Fortunately, by considering a screw line of the screw function attached to $\zeta(s)$, which will be explained in Sections 2.1 and 4.2, we can *unconditionally* construct two special Hilbert spaces $\mathcal{H}_0$ and $\mathcal{K}_0$ (in Section 3.3) to be isomorphic to $\mathcal{H}_W$ and $\mathcal{K}(\Theta_\xi)$, respectively, under the RH (Theorem 5.6). The construction of such spaces leads to an equivalence condition for the RH stated below. That is the second main result.

In Selberg's answer to the second question in [1, p. 632], he states that the construction of a space assuming the RH will not be useful for attacking the RH. However, $\mathcal{H}_0$ and $\mathcal{K}_0$ may be useful in future research on the RH, since they are defined *without* the RH.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Let $L^2(\mathbb{R})$ be the usual L^2-space on the real line with respect to the Lebesgue measure. We define

$$\mathfrak{S}_t(z) := \frac{i(1+\Theta_\xi^\sharp(z))}{2}\,\mathfrak{P}_t(z) \tag{1.5}$$

with

$$\begin{aligned}\mathfrak{P}_t(z) := {} & \frac{4(e^{t/2}-1)}{1+2iz} + \frac{4(e^{-t/2}-1)}{1-2iz} \\ & + \frac{e^{-izt}-1}{iz}\frac{\zeta'}{\zeta}\left(\frac{1}{2}-iz\right) + \sum_{n\le e^t}\frac{\Lambda(n)}{\sqrt{n}}\frac{e^{-iz(t-\log n)}-1}{iz} \\ & - \frac{1}{2iz}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}-\frac{iz}{2}\right) - \frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)\right] \\ & - \frac{1}{2iz}e^{-t/2}\left[\Phi\left(e^{-2t},1,\tfrac{1}{2}\left(\tfrac{1}{2}-iz\right)\right) - \Phi\left(e^{-2t},1,\tfrac{1}{4}\right)\right]\end{aligned} \tag{1.6}$$

for a nonnegative real number t and a complex number z, where $\Lambda(n)$ is the von Mangoldt function defined by $\Lambda(n) = \log p$ if $n = p^k$ with $k \in \mathbb{Z}_{>0}$ and $\Lambda(n) = 0$ otherwise, and

$$\Phi(z,s,a) = \sum_{n=0}^{\infty}\frac{z^n}{(n+a)^s}$$

is the Hurwitz–Lerch zeta-function. For negative t, we set $\mathfrak{S}_t(z) := \mathfrak{S}_{-t}(z)$. The definition of $\mathfrak{P}_t(z)$ is quite complicated. However, using the set Γ of zeros of $\xi(1/2-iz)$, it can be expressed in the simple form (3.2) (see Proposition 3.1 below). Nevertheless, as a tool for stating an equivalent condition for the RH, it seems preferable to have a representation that does not involve Γ. Thus, here we adopt a version of (3.2) rewritten without Γ using Weil's explicit formula (3.3). In Weil's explicit formula, the first, second, and the third–fourth lines on the right-hand side of (1.6) correspond to the poles of the completed zeta-function $\pi^{-s/2}\Gamma(s/2)\zeta(s)$, the non-archimedean part (Euler product), and the archimedean part (gamma factor), respectively.

For this $\mathfrak{S}_t$, we first obtain the following.

Proposition 1.2 *For any fixed $t \in \mathbb{R}$, $\mathfrak{S}_t(z)$ belongs to $L^2(\mathbb{R})$ as a function of z.*

Proof See Section 3.2. ∎

From this result, the mapping $t \mapsto \mathfrak{S}_t(z)$ from $\mathbb{R}$ to $L^2(\mathbb{R})$ is defined. By the uniformity of the L^2-norm of $\mathfrak{S}_t(z)$ on a compact set of t obtained in the proof of Proposition 1.2 and Minkowski's integral inequality, the following holds.

Proposition 1.3 *For $\phi \in C_c^\infty(\mathbb{R})$, we define*

$$\widehat{\mathcal{P}_\phi}(z) := \int_{-\infty}^{\infty}\mathfrak{S}_t^\sharp(z)\phi(t)\,dt \left(= \int_{-\infty}^{\infty}\overline{\mathfrak{S}_t(\bar{z})}\,\phi(t)\,dt\right) \tag{1.7}$$

using (1.5). Then, $\widehat{\mathcal{P}_\phi}(z)$ belongs to $L^2(\mathbb{R})$.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Using the image of the composition $\widehat{\mathcal{P}_D} := \widehat{\mathcal{P}} \circ D$ of the integral operator $\widehat{\mathcal{P}}$ and the differential operator

$$(D\psi)(t) := i\psi'(t), \tag{1.8}$$

we obtain the following equivalence condition for the RH.

Theorem 1.4 *The RH is true if and only if the equality*

$$\|\widehat{\mathcal{P}_{D\psi}}\|^2_{L^2(\mathbb{R})} = \pi\langle\psi,\psi\rangle_W \tag{1.9}$$

holds for all $\psi \in C_c^\infty(\mathbb{R})$. Furthermore, by choosing the test functions appropriately, if (1.9) holds for countably many choices of ψ's, then the RH follows.

Proof See Section 4.3. ∎

Equation (1.9) is reformulated to the following simpler form.

Corollary 1.5 *Define the subspace $V^\circ(0)$ of $L^2(\mathbb{R})$ by*

$$V^\circ(0) := \left\{ \mathsf{F}^{-1}\widehat{\mathcal{P}_{D\psi}} \,\middle|\, \psi \in C_c^\infty(\mathbb{R}) \right\}.$$

Then, the RH is true if and only if the equality

$$2\|\psi\|^2_{L^2(\mathbb{R})} = \langle\psi,\psi\rangle_W \tag{1.10}$$

holds for all $\psi \in V^\circ(0)$.

Proof See Section 4.3 and Theorem 5.6. ∎

The advantage of Theorem 1.4 and Corollary 1.5 is that it has turned the criterion of the RH from a set of inequalities like Weil's criterion into a set of equalities. It should also be noted that equations (1.9) and (1.10) can be expressed without zeros of $\xi(1/2 - iz)$ by (1.5) and (1.6). Furthermore, equations (1.9) and (1.10) claim that the nonnegativity of Weil's Hermitian form is explained by the nonnegativity of the L^2-norm.

In the following sections, first, in Section 2, we briefly review necessary notions, such as screw functions, screw lines, the Hermite–Biehler class, de Branges spaces, and model spaces. Then, in Section 3, we state and prove unconditional results that we need to prove the main results. Moreover, we unconditionally define two Hilbert spaces $\mathcal{H}_0$ and $\mathcal{K}_0$ that agree with the Hilbert spaces $\mathcal{H}_W$ and $\mathcal{K}(\Theta)$, respectively, under the RH.

In Section 4, we show that $\mathfrak{S}_t(z)$ in (1.5) gives a screw line of the screw function corresponding to the Riemann zeta-function under the RH (Theorem 4.2). Furthermore, we prove Theorem 1.4 and Corollary 1.5. The strategy of the proof of Theorem 4.2 is basically the same as that of [15, Theorem 1.1], with Proposition 4.1 playing an essential role in both cases. To carry this out, the rewriting of (1.5) into (3.6), prepared in Section 3 using Weil's explicit formula, corresponds to the transformation from (1.7) to (3.6) in [15], although the technical details of the calculations differ considerably. On the other hand, the analytic or geometric meaning of the functions giving the norms was unclear in [15], whereas in the present article, these functions have a clear

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

interpretation as a screw line. Furthermore, as an advantage of employing the screw line $\mathfrak{S}_t(z)$, we obtain Theorem 1.4, for which no analog was obtained in [15].

In Section 5, we prove Theorem 1.1 in a more detailed form. In addition, we prove that $\mathcal{H}_0 = \mathcal{H}_W$ and $\mathcal{K}_0 = \mathcal{K}(\Theta_\xi)$ under the RH. Afterward, we explain that the Hilbert space $\mathcal{H}_W$ is one of the Hilbert–Pólya spaces in Section 6. Finally, we mention two special values of $\mathfrak{S}_t(z)$ in Section 7 as an appendix.

2 Review on necessary notions

2.1 Screw functions and screw lines

In this and the next part, we refer to [7, Sections 5 and 12]. See also its references for details. Following Kreĭn, we denote by $\mathcal{G}_\infty$ the space of all continuous functions $g(t)$ on $\mathbb{R}$ such that $g(-t) = \overline{g(t)}$ and the kernel

$$G_g(t,u) := g(t-u) - g(t) - g(-u) + g(0) \tag{2.1}$$

is nonnegative definite on $\mathbb{R}$, that is, $\sum_{i,j=1}^n G_g(t_i,t_j)\,\xi_i\overline{\xi_j} \geq 0$ for all $n \in \mathbb{N}$, $t_i \in \mathbb{R}$, and $\xi_i \in \mathbb{C}(i = 1,2,\dots,n)$. Functions belonging to $\mathcal{G}_\infty$ are called *screw functions* on $\mathbb{R}$.

If an (even) real-valued function $g(t)$ is a screw function, then there exists a Hilbert space $\mathcal{H}$ and a continuous mapping $t \mapsto x(t)$ from $\mathbb{R}$ into $\mathcal{H}$ such that

$$\langle x(t+v) - x(v), x(u+v) - x(v)\rangle_{\mathcal{H}}$$

is independent of $v \in \mathbb{R}$ for all $t,u \in \mathbb{R}$ and the equality $\langle x(t) - x(0), x(u) - x(0)\rangle_{\mathcal{H}} = G_g(t,u)$ holds. Therefore, $\|x(t) - x(0)\|_{\mathcal{H}}^2 = -2g(t)$ if $g(0) = 0$. A mapping $x : \mathbb{R} \to \mathcal{H}$ endowed with the translation-invariance described above is called a *screw line* for $g(t)$.

2.2 Hilbert spaces associated with screw functions

Each $g \in \mathcal{G}_\infty$ defines a nonnegative definite Hermitian form on $\mathbb{R}$ by

$$\langle \phi_1, \phi_2\rangle_{G_g} := \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} G_g(t,u)\phi_1(u)\overline{\phi_2(t)}\,dudt. \tag{2.2}$$

According to [7, Section 5], we denote by $\mathcal{L}(G_g)$ the space $C_0(\mathbb{R})$ of all continuous and compactly supported functions ϕ on $\mathbb{R}$ such that $\widehat{\phi}(0) = 0$ equipped with the Hermitian inner product $\langle\cdot,\cdot\rangle_{G_g}$. We also denote by $\mathcal{H}(G_g)$ the completion of the factor space $\mathcal{L}(G_g)/\mathcal{L}^\circ(G_g)$, where $\mathcal{L}^\circ(G_g) = \{\phi \in \mathcal{L}(G_g) \mid \langle\phi,\phi\rangle_{G_g} = 0\}$. Note that even if $\langle\cdot,\cdot\rangle_{G_g}$ is positive definite on $\mathcal{L}(G_g)$, that is, $\mathcal{L}^\circ(G_g) = \{0\}$, there possibly exists a sequence $(\phi_n)_n$ of $\mathcal{L}(G_g)$ such that $\phi_n \to 0$ as $n \to \infty$ with respect to $\langle\cdot,\cdot\rangle_{G_g}$. The completion $\mathcal{H}(G_g)$ is a space of equivalence classes of Cauchy sequences with respect to $\langle\cdot,\cdot\rangle_{G_g}$. Two Cauchy sequences are equivalent if their difference converges to zero with respect to $\langle\cdot,\cdot\rangle_{G_g}$. We denote by $[\phi] \in \mathcal{H}(G_g)$ the equivalence class represented by ϕ. In general, elements of $\mathcal{H}(G_g)$ are not necessarily represented by functions unlike $\mathcal{H}_W$ (cf. [7, Section 4.3]).

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Every $g \in \mathcal{G}_\infty$ admits a representation

$$g(t) = g(0) + ibt + \int_{-\infty}^{\infty} \left(e^{i\lambda t} - 1 - \frac{i\lambda t}{1+\lambda^2} \right) \frac{d\tau(\lambda)}{\lambda^2} \tag{2.3}$$

with $b \in \mathbb{R}$ and a measure τ on $\mathbb{R}$ such that $\int_{-\infty}^{\infty} d\tau(\lambda)/(1+\lambda^2) < \infty$ and vice versa. If $g(t)$ is real-valued, $b = 0$. Without loss of generality, we suppose that $g(0) = 0$.

We define

$$\Phi_1(\phi, \lambda) := \int_{-\infty}^{\infty} \frac{e^{i\lambda x} - 1}{\lambda} \phi(x)\, dx = \frac{\widehat{\phi}(\lambda) - \widehat{\phi}(0)}{\lambda} = \frac{\widehat{\phi}(\lambda)}{\lambda}$$

for $\phi \in \mathcal{L}(G_g)$. Then, $\langle \phi_1, \phi_2 \rangle_{G_g} = \langle \Phi_1(\phi_1), \Phi_1(\phi_2) \rangle_{L^2(\tau)}$ for $\phi_1, \phi_2 \in \mathcal{L}(G_g)$ and Φ_1 establishes an isomorphism between $\mathcal{H}(G_g)$ and $L^2(\tau)$.

2.3 De Branges spaces

In this part, we refer to [12, 18]. See also those references for details. Let $H^2 := H^2(\mathbb{C}_+) = \mathsf{F}(L^2(0,\infty))$ be the Hardy space in the upper half-plane. As usual, we identify H^2 with a closed subspace of $L^2(\mathbb{R})$ via boundary values. Then, the inner product of H^2 coincides with the standard inner product of $L^2(\mathbb{R})$.

The Hermite–Biehler class consists of entire functions E satisfying $|E^\sharp(z)| < |E(z)|$ for all $z \in \mathbb{C}_+$. For each entire function E belonging to the Hermite–Biehler class, the de Branges space $\mathcal{H}(E)$ is defined as a Hilbert space consisting of entire functions $F(z)$ such that both $F(z)/E(z)$ and $F^\sharp(z)/E(z)$ belong to H^2 and have the norm

$$\|F\|_{\mathcal{H}(E)} := \|F/E\|_{L^2(\mathbb{R})}. \tag{2.4}$$

The multiplication operator M by an independent variable is defined by $\mathfrak{D}(\mathsf{M}) = \{F(z) \in \mathcal{H}(E) \,|\, zF(z) \in \mathcal{H}(E)\}$ and $(\mathsf{M}F)(z) = zF(z)$ for $F \in \mathfrak{D}(\mathsf{M})$. The domain $\mathfrak{D}(\mathsf{M})$ is dense in $\mathcal{H}(E)$ if and only if

$$S_\theta(z) := \frac{i}{2}(e^{i\theta}E(z) - e^{-i\theta}E^\sharp(z))$$

does not belong to $\mathcal{H}(E)$ for all $\theta \in [0, \pi)$ [12, Theorem 11]. The particular two θ cases are often written as $A(z) := -S_{\pi/2}(z)$ and $B(z) := S_0(z)$.

If $\mathfrak{D}(\mathsf{M})$ is dense in $\mathcal{H}(E)$, all self-adjoint extensions of M are parametrized by $\theta \in [0, \pi)$ and are described as follows. The domain of M_θ is

$$\mathfrak{D}(\mathsf{M}_\theta) = \left\{ G(z) = \frac{S_\theta(w_0)F(z) - S_\theta(z)F(w_0)}{z - w_0} \;\middle|\; F(z) \in \mathcal{H}(E) \right\}, \tag{2.5}$$

and the operation is defined by

$$\mathsf{M}_\theta G(z) = z\, G(z) + F(w_0)S_\theta(z), \tag{2.6}$$

where w_0 is a fixed complex number with $S_\theta(w_0) \neq 0$ [6, Propositions 4.6 and 6.1]. The domain $\mathfrak{D}(\mathsf{M}_\theta)$ is independent of the choice of the number w_0. For a fixed $\theta \in [0, \pi)$, we confirm that $G(z) = S_\theta(z)/(z - \gamma)$ belongs to $\mathfrak{D}(\mathsf{M}_\theta)$ by taking

$$F(z) = \frac{S_\theta(z)}{S_\theta(w_0)} \frac{\gamma - w_0}{z - \gamma}$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

for every zero γ of $S_\theta(z)$ and is an eigenfunction of M_θ with the eigenvalue γ. Further, $\{S_\theta(z)/(z-\gamma) \,|\, S_\theta(\gamma) = 0\}$ forms an orthogonal basis of $\mathcal{H}(E)$ [5, Theorem 22].

2.4 Model subspaces

In this part, we refer to [9, Section 2], [13, Section 3.5], and [15, Section 3.1]. See also those references for details.

Let $H^\infty = H^\infty(\mathbb{C}_+)$ be the space of all bounded analytic functions in $\mathbb{C}_+$. A function $\Theta \in H^\infty$ is called an inner function in $\mathbb{C}_+$ if $\lim_{y\to 0+} |\Theta(x+iy)| = 1$ for almost all $x \in \mathbb{R}$. For an inner function Θ, a model space $\mathcal{K}(\Theta)$ is defined as the orthogonal complement $\mathcal{K}(\Theta) = H^2 \ominus \Theta H^2$ and has the alternative representation

$$\mathcal{K}(\Theta) = H^2 \cap \Theta \bar{H}^2, \tag{2.7}$$

where $\Theta H^2 = \{\Theta(z)F(z) \,|\, F \in H^2\}$ and $\bar{H}^2 = H^2(\mathbb{C}_-)$ is the Hardy space in the lower half-plane. The model space $\mathcal{K}(\Theta)$ is a subspace of $L^2(\mathbb{R})$ as a Hilbert space. In particular, the inner product of $\mathcal{K}(\Theta)$ matches that of $L^2(\mathbb{R})$ on the real line.

If an inner function Θ in $\mathbb{C}_+$ extends to a meromorphic function in $\mathbb{C}$, then it is called a meromorphic inner function in $\mathbb{C}_+$. For any meromorphic inner function Θ, there exists E of the Hermite–Biehler class such that $\Theta = E^\sharp/E$. The de Branges space $\mathcal{H}(E)$ is isometrically isomorphic to $\mathcal{K}(\Theta)$ by $F(z) \mapsto E(z)F(z)$. In particular, $\mathcal{H}(E) = E\,H^2 \cap E^\sharp\,\bar{H}^2$.

For a meromorphic inner function Θ, let μ_Θ be the positive discrete measure on $\mathbb{R}$ supported on $\sigma(\Theta) = \{x \in \mathbb{R} \,|\, \Theta(x) = -1\}$ and

$$\mu_\Theta(x) = \frac{2\pi}{|\Theta'(x)|}. \tag{2.8}$$

Then, the restriction map $F \mapsto F|_{\sigma(\Theta)}$ is an isometric operator from $\mathcal{K}(\Theta)$ to $L^2(\mu_\Theta)$ [9, Theorem 2.1]. The isometric property of the map implies that the family of functions

$$f_\gamma(z) = \sqrt{\frac{2}{\pi|\Theta'(\gamma)|}}\,\frac{1+\Theta(z)}{2(z-\gamma)} = \sqrt{\frac{2}{\pi|\Theta'(\gamma)|}}\,\frac{A(z)}{(z-\gamma)E(z)} \tag{2.9}$$

parametrized by all zeros γ of $A(z) = -S_{\pi/2}(z)$ forms an orthonormal basis of $\mathcal{K}(\Theta)$ if $\mathfrak{D}(\mathsf{M})$ is dense in $\mathcal{H}(E)$.

3 Unconditional results

Throughout this and later sections, we denote $E = E_\xi$ and $\Theta = \Theta_\xi = E_\xi^\sharp/E_\xi$ for functions defined in (1.3) and (1.4), respectively. Otherwise, it is mentioned.

3.1 Expansion of $\mathfrak{P}_t(z)$ over the zeros

For the basic properties of the Riemann zeta-function, we refer to [16]. By the two functional equations $\xi(s) = \xi(1-s)$ and $\xi(s) = \xi^\sharp(s)$, if γ belongs to the set of zeros Γ, then both $-\gamma$ and $\overline{\gamma}$ also belong to Γ with the same multiplicity. On the other hand, $|\Im(\gamma)| < 1/2$ for every $\gamma \in \Gamma$, since all zeros of $\xi(s)$ lie in the strip $0 < \Re(s) < 1$. For $E(z)$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

of (1.3), we define

$$A(z) := (E(z) + E^\sharp(z))/2 \tag{3.1}$$

as in Section 2.3. Then, $A(z) = \xi(1/2 - iz)$, because $E^\sharp(z) = \overline{E(\bar{z})} = \xi(1/2 - iz) - \xi'(1/2 - iz)$ by functional equations of $\xi(s)$. Therefore, the set Γ coincides with the set of all zeros of both $A(z)$ and $1 + \Theta(z)$. We define

$$P_t(z) := \sum_{\gamma\in\Gamma} m_\gamma \frac{e^{-i\gamma t} - 1}{\gamma} \cdot \frac{1}{z - \gamma} \tag{3.2}$$

for nonnegative t. For negative t, we set $P_t(z) := P_{-t}(z)$. The series on the right-hand side of (3.2) converges absolutely and uniformly on every compact subset of $\mathbb{C}\backslash\Gamma$, since $\sum_{\gamma\in\Gamma} m_\gamma|\gamma|^{-1-\delta} < \infty$ for any $\delta > 0$, because $A(z)$ is an entire function of order one. Therefore, $P_t(z)$ is a meromorphic function on $\mathbb{C}$ with Γ as the set of all poles.

Proposition 3.1 *Let $\mathfrak{P}_t(z)$ and $P_t(z)$ be meromorphic functions defined by (1.6) and (3.2), respectively. Then, both coincide.*

Proof For $t \geq 0$ and $z \in \mathbb{C}_+$, we define

$$\phi_{z,t}(x) = \begin{cases} (iz)^{-1} e^{izx}(e^{-izt} - 1), & t < x, \\ (iz)^{-1} e^{izx}(e^{-izx} - 1), & 0 \leq x \leq t, \\ 0, & x < 0. \end{cases}$$

The main tool for the proof is the Weil explicit formula

$$\begin{aligned} &\lim_{X\to\infty} \sum_{\substack{\gamma\in\Gamma \\ |\gamma|\leq X}} m_\gamma \int_{-\infty}^{\infty} \phi(x)\, e^{-i\gamma x}\, dx \\ &= \int_{-\infty}^{\infty} \phi(x)(e^{x/2} + e^{-x/2})dx - \sum_{n=1}^{\infty} \frac{\Lambda(n)}{\sqrt{n}} \phi(\log n) - \sum_{n=1}^{\infty} \frac{\Lambda(n)}{\sqrt{n}} \phi(-\log n) \\ &\quad - (\log 4\pi + \gamma_0)\phi(0) - \int_0^{\infty} \left\{\phi(x) + \phi(-x) - 2e^{-x/2}\phi(0)\right\} \frac{e^{x/2}dx}{e^x - e^{-x}}, \end{aligned} \tag{3.3}$$

which is obtained from the explicit formula in [3, p. 186] by taking $\phi(x) = e^{x/2} f(e^x)$ for test functions $f(t)$ in that formula with the conditions for $f(t)$ in [2, Section 3], where γ_0 is the Euler–Mascheroni constant. (Note that the formula in [2] has two typographical errors in the second line of the right-hand side.)

As is easily seen, Weil's explicit formula can be applied to $\phi(x) = \phi_{z,t}(x)$. We have

$$\int_{-\infty}^{\infty} \phi_{z,t}(x)\, e^{-i\gamma x}\, dx = \frac{e^{-i\gamma t} - 1}{\gamma} \cdot \frac{1}{z - \gamma} \quad \text{when } \Im(z) > \Im(\gamma).$$

Therefore, the left-hand side of Weil's explicit formula for $\phi_{z,t}(x)$ gives $P_t(z)$ of (3.2) when $\Im(z) > 1/2$. Hence, if it is shown that the right-hand side is equal to $\mathfrak{P}_t(z)$ for $\Im(z) > 1/2$, then the conclusion of the proposition follows by analytic continuation.

It is easy to verify

$$\int_{-\infty}^{\infty} \phi_{z,t}(x)(e^{x/2} + e^{-x/2})dx = \frac{4(e^{t/2} - 1)}{1 + 2iz} + \frac{4(e^{-t/2} - 1)}{1 - 2iz}$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

and

$$\sum_{n=1}^{\infty}\frac{\Lambda(n)}{\sqrt{n}}\phi_{z,t}(\log n)=\frac{1}{iz}\sum_{n\le e^t}\frac{\Lambda(n)}{\sqrt{n}}(1-n^{iz})+\frac{e^{-izt}-1}{iz}\sum_{t<\log n}\frac{\Lambda(n)}{n^{1/2-iz}}$$
$$=-\sum_{n\le e^t}\frac{\Lambda(n)}{\sqrt{n}}\frac{e^{-iz(t-\log n)}-1}{iz}-\frac{e^{-izt}-1}{iz}\frac{\zeta'}{\zeta}\left(\frac{1}{2}-iz\right),$$
$$\sum_{n=1}^{\infty}\frac{\Lambda(n)}{\sqrt{n}}\phi_{z,t}(-\log n)=0,\qquad \phi_{z,t}(0)=0$$

for $\Im(z)>1/2$ by direct calculation.

Therefore, the remaining task is to calculate the fifth term on the right-hand side. We split it into $\int_t^{\infty}$ and $\int_0^t$. For the first integral,

$$\int_t^{\infty}\left\{\phi_{z,t}(x)+\phi_{z,t}(-x)-2e^{-x/2}\phi_{z,t}(0)\right\}\frac{e^{x/2}dx}{e^x-e^{-x}}$$
$$=\frac{e^{-izt}-1}{iz}\int_t^{\infty}e^{izx}\frac{e^{x/2}dx}{e^x-e^{-x}}=\frac{e^{-izt}-1}{iz}\int_t^{\infty}e^{izx}\,e^{-x/2}\sum_{n=0}^{\infty}e^{-2nx}\,dx$$
$$=\frac{e^{-izt}-1}{2iz}e^{-t(\frac{1}{2}-iz)}\sum_{n=0}^{\infty}\frac{e^{-2nt}}{n+\frac{1}{2}(\frac{1}{2}-iz)}$$
$$=\frac{e^{-izt}-1}{2iz}e^{-t(\frac{1}{2}-iz)}\Phi(e^{-2t},1,\tfrac{1}{2}(\tfrac{1}{2}-iz)).$$

For the second integral,

$$\int_0^t\left\{\phi_{z,t}(x)+\phi_{z,t}(-x)-2e^{-x/2}\phi_{z,t}(0)\right\}\frac{e^{x/2}dx}{e^x-e^{-x}}$$
$$=-\frac{1}{iz}\int_0^t(e^{izx}-1)\frac{e^{x/2}dx}{e^x-e^{-x}}=-\frac{1}{iz}\int_0^t(e^{izx}-1)\,e^{-x/2}\sum_{n=0}^{\infty}e^{-2nx}\,dx.$$

To handle the right-hand side, we calculate as

$$\int_0^t(e^{izx}-1)\,e^{-x/2}\sum_{n=0}^{N}e^{-2nx}\,dx$$
$$=\frac{1}{2}\sum_{n=0}^{N}\left[\frac{1-e^{-2t(n+\frac{1}{2}(\frac{1}{2}-iz))}}{n+\frac{1}{2}(\frac{1}{2}-iz)}-\frac{1-e^{-2t(n+\frac{1}{4})}}{n+\frac{1}{4}}\right]$$
$$=-\frac{1}{2}e^{-t(\frac{1}{2}-iz)}\sum_{n=0}^{N}\frac{e^{-2tn}}{n+\frac{1}{2}(\frac{1}{2}-iz)}+\frac{1}{2}e^{-t/2}\sum_{n=0}^{N}\frac{e^{-2tn}}{n+\frac{1}{4}}$$
$$+\frac{1}{2}\sum_{n=0}^{N}\left[\frac{1}{n+\frac{1}{2}(\frac{1}{2}-iz)}-\frac{1}{n+\frac{1}{4}}\right]$$
$$=-\frac{1}{2}e^{-t(\frac{1}{2}-iz)}\Phi(e^{-2t},1,\tfrac{1}{2}(\tfrac{1}{2}-iz))+\frac{1}{2}e^{-t/2}\Phi(e^{-2t},1,\tfrac{1}{4})$$
$$-\frac{1}{2}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}-\frac{iz}{2}\right)-\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)\right]+O(e^{-2Nt})+O(N^{-1})$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

using the well-known series expansion

$$\frac{\Gamma'}{\Gamma}(w) = -\gamma_0 - \sum_{n=0}^{\infty}\left(\frac{1}{w+n} - \frac{1}{n+1}\right), \tag{3.4}$$

where the implied constant depends on t and z. Therefore, we obtain

$$\begin{aligned}\int_0^t &\left\{\phi_{z,t}(x) + \phi_{z,t}(-x) - 2e^{-x/2}\phi_{z,t}(0)\right\}\frac{e^{x/2}dx}{e^x - e^{-x}} \\ &= \frac{1}{2iz}e^{-t(\frac{1}{2}-iz)}\Phi(e^{-2t}, 1, \tfrac{1}{2}(\tfrac{1}{2} - iz)) - \frac{1}{2iz}e^{-t/2}\Phi(e^{-2t}, 1, \tfrac{1}{4}) \\ &\quad + \frac{1}{2iz}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4} - \frac{iz}{2}\right) - \frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)\right].\end{aligned}$$

Combining the results for $\int_t^\infty$ and $\int_0^t$,

$$\begin{aligned}\int_0^\infty &\left\{\phi_{z,t}(x) + \phi_{z,t}(-x) - 2e^{-x/2}\phi_{z,t}(0)\right\}\frac{e^{x/2}dx}{e^x - e^{-x}} \\ &= \frac{1}{2iz}e^{-t/2}\left[\Phi(e^{-2t}, 1, \tfrac{1}{2}(\tfrac{1}{2} - iz)) - \Phi(e^{-2t}, 1, \tfrac{1}{4})\right] \\ &\quad + \frac{1}{2iz}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4} - \frac{iz}{2}\right) - \frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)\right].\end{aligned}$$

From the calculation of the five terms on the right-hand side above, we conclude that the right-hand side of the Weil explicit formula for $\phi_{z,t}(x)$ equals (1.6). ∎

3.2 Proof of Proposition 1.2

We have $|\Theta(z)| = 1$ for every $z \in \mathbb{R}$ by definition. In fact, zeros of $E(z)$ in the denominator cancel out in the numerator $E^\sharp(z)$, even if they exist. Further, $\mathfrak{P}_t(z)$ has poles of order one at $\gamma \in \Gamma$, but $\mathfrak{S}_t(z)$ is holomorphic there, since $(1 + \Theta(z))/2 = A(z)/E(z) = A(z)/(A(z) + iA'(z)) = (z - \gamma)(-i/m_\gamma + o(1))$ near $z = \gamma$ by direct calculation. Hence, $\mathfrak{S}_t(z)$ is bounded and holomorphic on the real line by (1.5), (3.2), and Proposition 3.1. On the other hand, in the horizontal strip $|\Im(z)| \le 1/2$, we have the well-known estimate $(\Gamma'/\Gamma)(1/4 + iz/2) \ll \log|z|$ and

$$\frac{\zeta'}{\zeta}\left(\frac{1}{2} - iz\right) = \sum_{|\Re(z)-\gamma|\le 1}\frac{i}{z-\gamma} + O(\log|z|)$$

by [16, Theorem 9.6(A)]. In both estimates, implied constants are uniform in $|\Im(z)| \le 1/2$. The number of zeros $\gamma \in \Gamma$ satisfying $|\Re(z) - \gamma| \le 1$ is $O(\log|z|)$ counting with multiplicity by [16, Theorem 9.2]. Therefore, $\mathfrak{S}_t(z) \ll |z|^{-1}\log|z|$ as $|z| \to \infty$ with an implied constant depending on a compact set of t by (1.6). Hence, $\mathfrak{S}_t(z)$ belongs to $L^2(\mathbb{R})$ and the norm is uniformly bounded on a compact set of t.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

3.3 Two special Hilbert spaces

We first introduce the set of meromorphic functions

$$F_\gamma(z) := \sqrt{\frac{m_\gamma}{\pi}}\,\frac{i(1+\Theta(z))}{2(z-\gamma)}, \quad \gamma \in \Gamma. \tag{3.5}$$

Then, we have

$$\mathfrak{S}_t(z) = \sum_{\gamma\in\Gamma} \sqrt{\pi m_\gamma}\,\frac{e^{-i\gamma t}-1}{\gamma}\,F_\gamma^\sharp(z) \tag{3.6}$$

by Proposition 3.1. Therefore,

$$\widehat{\mathcal{P}_\phi}(z) = \sum_{\gamma\in\Gamma} \sqrt{\pi m_\gamma}\,\frac{\widehat{\phi}(\gamma)-\widehat{\phi}(0)}{\gamma}\,F_\gamma(z) \tag{3.7}$$

for any $\phi \in C_c^\infty(\mathbb{R})$ by definition (1.7) and the symmetry $\gamma \mapsto \bar{\gamma}$ of Γ with $m_\gamma = m_{\bar{\gamma}}$. This implies

$$\widehat{\mathcal{P}_{D\psi}}(z) = \sum_{\gamma\in\Gamma} \sqrt{\pi m_\gamma}\,\widehat{\psi}(\gamma)\,F_\gamma(z) \tag{3.8}$$

for any $\psi \in C_c^\infty(\mathbb{R})$, since $(\widehat{D\psi}(z) - \widehat{D\psi}(0))/z = \widehat{D\psi}(z)/z = \widehat{\psi}(z)$ for D in (1.8).

On the other hand, we define the norm $\|\ \|_0$ on $C_c^\infty(\mathbb{R})$ by

$$\|\psi\|_0 := \frac{1}{\sqrt{\pi}}\,\|\widehat{\mathcal{P}_{D\psi}}\|_{L^2(\mathbb{R})}, \quad \psi \in C_c^\infty(\mathbb{R}) \tag{3.9}$$

based on Proposition 1.3. Then, we have the following.

Lemma 3.2 *Equation* (3.9) *defines a norm on* $C_c^\infty(\mathbb{R})$.

Proof We obtain $\|\psi_1+\psi_2\|_0 \le \|\psi_1\|_0 + \|\psi_2\|_0$ and $\|k\psi\|_0 = |k|\|\psi\|_0$ for $\psi_1, \psi_2, \psi \in C_c^\infty(\mathbb{R})$ and $k \in \mathbb{C}$ by the obvious linearity of $\widehat{\mathcal{P}_D}$. Therefore, the proof is completed if it is shown that $\|\psi\|_0 = 0$ implies $\psi = 0$. If $\|\psi\|_0 = 0$, the image $\widehat{\mathcal{P}_{D\psi}}(z)$ is identically zero. The latter means that $\widehat{\psi}(\gamma) = 0$ for all $\gamma \in \Gamma$, because, if not, there must exist a sequence $(c_\gamma)_{\gamma\in\Gamma}$ such that $\sum_{\gamma\in\Gamma} c_\gamma (z-\gamma)^{-1}$ is identically zero on $\mathbb{C}$ by (3.5) and (3.8), but it is impossible. If $\widehat{\psi}(\gamma) = 0$ for all $\gamma \in \Gamma$, it implies that ψ is identically zero by [14, Lemma 2.1]. ∎

By Lemma 3.2, we can complete the space $C_c^\infty(\mathbb{R})$ with respect to $\|\ \|_0$. We denote the completion by $\mathcal{H}_0$. On the other hand, we denote the L^2-closure of the image $\widehat{\mathcal{P}_D}(C_c^\infty(\mathbb{R}))$ in $L^2(\mathbb{R})$ by $\mathcal{K}_0$. Then, two Hilbert spaces $\mathcal{H}_0$ and $\mathcal{K}_0$ are isometrically isomorphic up to a constant multiple. The map $\widehat{\mathcal{P}_D}$ from $C_c^\infty(\mathbb{R})$ to $\widehat{\mathcal{P}_D}(C_c^\infty(\mathbb{R})) \subset L^2(\mathbb{R})$ extends to the map from $\mathcal{H}_0$ to $\mathcal{K}_0$ by (3.9). As proved in Theorem 5.6 below, $\mathcal{H}_0 = \mathcal{H}_W$ and $\mathcal{K}_0 = \mathcal{K}(\Theta)$ under the RH.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

4 A screw line of the Riemann zeta-function

4.1 A special orthonormal basis

Assuming the RH is true, $E = E_\xi$ belongs to the Hermite–Biehler class [8, Theorem 1], and thus $\Theta = \Theta_\xi$ is a meromorphic inner function. Therefore, they define the de Branges space $\mathcal{H}(E)$ and the model space $\mathcal{K}(\Theta)$, respectively. We need the following result for the later discussion.

Proposition 4.1 *Assume that the RH is true. Then, the family* (3.5) *forms an orthonormal basis of the Hilbert space* $\mathcal{K}(\Theta)$. *Furthermore,*

$$\frac{\Theta'(\gamma)}{2} = -\frac{i}{m_\gamma} \tag{4.1}$$

and

$$F_\gamma(\gamma) = \frac{1}{\sqrt{m_\gamma \pi}}, \qquad F_\gamma(\gamma') = 0 \quad \textit{for every} \quad \gamma \in \Gamma,\ \gamma' \in \Gamma\backslash\{\gamma\}. \tag{4.2}$$

Proof See [15, Proposition 3.2] and its proof. ∎

4.2 Screw line of the Riemann zeta-function

We define the even real-valued function $g_\xi(t)$ on the real line by

$$\begin{aligned} g_\xi(t) := {} & -4\left(e^{t/2} + e^{-t/2} - 2\right) + \sum_{n \le e^t} \frac{\Lambda(n)}{\sqrt{n}}\left(t - \log n\right) \\ & - \frac{t}{2}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right) - \log \pi\right] - \frac{1}{4}\left(\Phi(1,2,1/4) - e^{-t/2}\Phi(e^{-2t},2,1/4)\right) \end{aligned} \tag{4.3}$$

for nonnegative t. We easily obtain $g_\xi(0) = 0$. Then, $g_\xi(t)$ is a screw function on $\mathbb{R}$ under the RH as stated in [14, Theorem 1.2]. One of the screw lines corresponding to $g_\xi(t)$ can be constructed as follows.

Let τ_ξ be the nonnegative measure representing $g_\xi(t)$ as in (2.3) under the RH. Then, the Hilbert space $\mathcal{H} = L^2(\tau_\xi)$ and the mapping $t \mapsto x(t) := (e^{it\gamma} - 1)/\gamma$ provide a screw line satisfying $\|x(t) - x(0)\|_{\mathcal{H}}^2 = -2g_\xi(t)$ [7, Section 12]. This spectral construction of the screw line is important and useful in analysis, but it is of limited use for studying the nontrivial zeros of $\zeta(s)$ without assuming the RH. In the following, we show that $\mathfrak{S}_t$ gives a screw line of $g_\xi(t)$. In contrast to the spectral screw line above, this screw line can be used to study $\mathcal{H}_W$, as will be done later.

Theorem 4.2 *Assume the RH is true and let* $g(t) = g_\xi(t)$. *Then, the mapping* $t \mapsto \pi^{-1/2}\mathfrak{S}_t(z)$ *from* $\mathbb{R}$ *to* $L^2(\mathbb{R})$ *is a screw line of* $g(t)$. *That is,*

$$\frac{1}{\pi}\langle \mathfrak{S}_t, \mathfrak{S}_u \rangle_{L^2(\mathbb{R})} = G_g(t,u) \tag{4.4}$$

holds for $t, u \in \mathbb{R}$.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Proof The sum of coefficients on the right-hand side of (3.6) is convergent in L^2-sense:

$$\sum_{\gamma\in\Gamma}\left|\sqrt{\pi m_\gamma}\frac{e^{-i\gamma t}-1}{\gamma}\right|^2 \le \pi\sum_{\gamma\in\Gamma}\frac{m_\gamma}{|\gamma|^2}<\infty.$$

Therefore, applying Proposition 4.1 to $\mathfrak{S}_t(z)$ via formula (3.6), we find that it belongs to the subspace $\mathcal{K}(\Theta)$ of $L^2(\mathbb{R})$ and

$$\frac{1}{\pi}\langle \mathfrak{S}_{t+v}-\mathfrak{S}_v, \mathfrak{S}_{u+v}-\mathfrak{S}_v\rangle_{L^2(\mathbb{R})} = \sum_{\gamma\in\Gamma} m_\gamma \frac{e^{-i\gamma t}-1}{\gamma}\cdot\frac{e^{i\gamma u}-1}{\gamma} \tag{4.5}$$

holds. The right-hand side is equal to $G_g(t,u)$ by the formula

$$G_g(t,u) = \sum_{\gamma\in\Gamma} m_\gamma \frac{(e^{i\gamma t}-1)(e^{-i\gamma u}-1)}{\gamma^2} \tag{4.6}$$

given in [14, Equation (1.9)] and the symmetry $\gamma\mapsto-\gamma$ of Γ with $m_\gamma = m_{-\gamma}$. Hence, $\pi^{-1/2}\mathfrak{S}_t : \mathbb{R}\to L^2(\mathbb{R})$ is a screw line of $g(t)$ under the RH.

We find that $\mathfrak{S}_0(z)$ is identically zero by (1.5) and (1.6), since

$$\lim_{t\to0}\left(\Phi(e^{-2t},1,\tfrac14)-\Phi(e^{-2t},1,\tfrac12(\tfrac12-iz))\right) = -\frac{\Gamma'}{\Gamma}\left(\frac14\right)+\frac{\Gamma'}{\Gamma}\left(\frac12\left(\frac12-iz\right)\right)$$

by (2.8). Therefore, by taking $v=0$ in (4.5), we obtain (4.4). ∎

The following immediately follows from Theorem 4.2.

Corollary 4.3 *The RH is true if and only if the equality*

$$\frac{1}{2\pi}\|\mathfrak{S}_t\|^2_{L^2(\mathbb{R})} = -g(t) \tag{4.7}$$

holds for all $t\ge t_0$ for some $t_0\ge0$.

Proof Assuming the RH, we obtain (4.7) by taking $u=t$ in (4.4), since $G_g(t,t) = -2g(t)$ by (2.1) and $g(0)=0$. Conversely, we suppose that equality (4.7) holds for all $t\ge t_0$. Then, $-g(t)$ is nonnegative on $[t_0,\infty)$, which implies that the RH is true by [14, Theorems 1.7 and 11.1]. ∎

4.3 Proof of Theorem 1.4

Theorem 1.4 is a corollary of the following result.

Theorem 4.4 *Let $g(t) = g_\xi(t)$. The RH is true if and only if the equality*

$$\|\widehat{\mathcal{P}_\phi}\|^2_{L^2(\mathbb{R})} = \pi\langle\phi,\phi\rangle_{G_g} \tag{4.8}$$

holds for all $\phi\in C_c^\infty(\mathbb{R})$ satisfying $\widehat{\phi}(0)=0$. If the RH is true, equality (4.8) holds for all $\phi\in C_c^\infty(\mathbb{R})$.

Proof First, we prove (4.8) assuming the RH holds. We have

$$\|\widehat{\mathcal{P}_\phi}\|^2_{L^2(\mathbb{R})} = \pi\sum_{\gamma\in\Gamma} m_\gamma\left|\frac{\widehat{\phi}(\gamma)-\widehat{\phi}(0)}{\gamma}\right|^2 \tag{4.9}$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

by (3.7) and Proposition 4.1. Applying (4.6) to (2.2) and noting the symmetry $\gamma \mapsto -\gamma$ of Γ with $m_\gamma = m_{-\gamma}$, we find that the right-hand side of (4.9) equals $\pi\langle\phi,\phi\rangle_{G_g}$.

Conversely, we prove that the RH is true assuming equality (4.8). We show that a contradiction arises if the RH is false. We take a nonreal $\gamma_0 \in \Gamma$. For any $\varepsilon > 0$, there exists $\psi_1,\ \psi_2 \in C_c^\infty(\mathbb{R})$ such that $\widehat{\psi_1}(-\gamma_0) = i$, $\widehat{\psi_2}(-\overline{\gamma_0}) = -i$, $|\widehat{\psi_1}(-\gamma)| \le \varepsilon|\gamma_0 - \gamma|^{-1-\delta}$ for every $\gamma \in \Gamma\backslash\{\gamma_0\}$, and $|\widehat{\psi_2}(-\gamma)| \le \varepsilon|\overline{\gamma_0} - \gamma|^{-1-\delta}$ for every $\gamma \in \Gamma\backslash\{\overline{\gamma_0}\}$ by [19, Lemma 1]. We define $\psi := \psi_1 + \psi_2\ (\neq 0)$ and set $\phi := D\psi$. Then, $\widehat{\phi}(0) = 0$ by definition, and $\langle\phi,\phi\rangle_{G_g} = \langle\psi,\psi\rangle_W$ holds by the relation

$$\langle D\psi_1, D\psi_2\rangle_{G_g} = \langle\psi_1,\psi_2\rangle_W \tag{4.10}$$

in [14, Proposition 3.1]. The right-hand side equals $\sum_{\gamma\in\Gamma} m_\gamma\widehat{\psi}(-\gamma)(\widehat{\psi})^\sharp(-\gamma) = -m_{\gamma_0} + O(\varepsilon)$, since $\sum_{\gamma\in\Gamma} m_\gamma|\gamma|^{-1-\delta} < \infty$. Therefore, $\langle\phi,\phi\rangle_{G_g}$ is negative for a sufficiently small $\varepsilon > 0$, but it contradicts the nonnegativity that follows from (4.8). ■

Proof of Theorem 1.4 The conclusion follows from Theorem 4.4 and the relation (4.10) of Hermitian forms, since the differential operator D in (1.8) gives a bijection from $C_c^\infty(\mathbb{R})$ to the subspace $C_0^\infty(\mathbb{R}) \subset C_c^\infty(\mathbb{R})$ consisting of functions ϕ with $\widehat{\phi}(0) = 0$. ■

Proof of Corollary 1.5 The RH is true if (1.10) holds by the same argument as the second half of the proof of Theorem 4.4. Therefore, we prove (1.10) assuming the RH.

Let $\psi \in V^\circ(0)$. Then, $\widehat{\psi}(z) = \widehat{\mathcal{P}_{D\psi_0}}(z)$ for some $\psi_0 \in C_c^\infty(\mathbb{R})$ by definition. Therefore, $\widehat{\psi}(z) = \sum_{\gamma\in\Gamma}\sqrt{\pi m_\gamma}\,\widehat{\psi_0}(\gamma)F_\gamma(z)$ by (3.8). The equality shows that $\widehat{\psi}(z)$ is a continuous function of $z \in \mathbb{R}$ by the uniform convergence of the right-hand side on a compact set of z. Taking $z = \gamma$ in this equality, we have $\widehat{\psi}(\gamma) = \widehat{\psi_0}(\gamma)$ by (4.2). Therefore, $\langle\psi,\psi\rangle_W$ is defined and satisfies $\langle\psi,\psi\rangle_W = \langle\psi_0,\psi_0\rangle_W$. The right-hand side is equal to $\|\widehat{\psi_0}\|^2_{L^2(\mathbb{R})} = 2\pi\|\psi_0\|^2_{L^2(\mathbb{R})}$ by (1.9) and Plancherel's identity. The same argument works if we start with $\psi_0 \in C_c^\infty(\mathbb{R})$. Hence, we obtain (1.10). ■

Using (3.9), Theorem 1.4 is stated as follows.

Theorem 4.5 *The RH is true if and only if the equality*

$$\|\psi\|_0^2 = \langle\psi,\psi\rangle_W \tag{4.11}$$

holds for all $\psi \in C_c^\infty(\mathbb{R})$.

Equality (4.11) leads to Theorem 5.6 below.

For $n \in \mathbb{Z}_{>0}$, we define

$$g_n(x) := e^{-x/2}\sum_{j=1}^{n}\binom{n}{j}\frac{(-x)^{j-1}}{(j-1)!} \quad (x > 0), \quad g_n(0) := \frac{n}{2}, \quad g_n(x) := 0 \quad (x < 0).$$

Then, the RH holds if $\langle g_n, g_n\rangle_W \ge 0$ for all $n \in \mathbb{Z}_{>0}$ by Bombieri and Lagarias [2, Section 4]. Therefore, we obtain the following.

Corollary 4.6 *The RH holds if* (4.11) *holds for all* g_n $(n \in \mathbb{Z}_{>0})$.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

5 Proof of Theorem 1.1 and its refinement

Throughout this section, we assume that the RH is true and denote $E = E_\xi$, $\Theta = \Theta_\xi = E_\xi^\sharp/E_\xi$ as before, and denote $g = g_\xi$. Therefore, E belongs to the Hermite–Biehler class, Θ is a meromorphic inner function in $\mathbb{C}_+$, and g belongs to the class of screw functions $\mathcal{G}_\infty$.

For use in the proof of Theorem 1.1 and its refinement, we introduce the operator K acting on $L^2(\mathbb{R})$ by

$$\mathsf{K} := \mathsf{F}^{-1}\mathsf{M}_\Theta \mathsf{J}\mathsf{F} \tag{5.1}$$

with

$$(\mathsf{M}_\Theta F)(z) := \Theta(z)F(z) \quad \text{and} \quad (\mathsf{J}F)(z) := F^\sharp(z).$$

The Fourier transform F, the multiplication operator M_Θ, and the involution J are defined for functions of a complex variable, and the latter two are isometries on $L^2(\mathbb{R})$. The Fourier transform F is an isometry up to a constant factor. Therefore, K is isometric on $L^2(\mathbb{R})$. Further, K is invertible by $\mathsf{K}^2 = \mathrm{id}$. By definition, K is *not* $\mathbb{C}$-linear but $\mathbb{R}$-linear and conjugate linear. Using the isometric involution K, we define

$$V(t) := L^2(t,\infty) \cap \mathsf{K}L^2(t,\infty) \tag{5.2}$$

and

$$\mathcal{H}_W(t) := \{\, [\psi] \mid \psi \in V(t) \,\}$$

for $t \geq 0$. The set of subspaces $V(t)$ of $L^2(\mathbb{R})$ are clearly totally ordered by the set-theoretical inclusion.

First, Theorem 1.1 is shown using $V(t)$ for $t = 0$, and it is refined using general $t \geq 0$.

Lemma 5.1 *Let $V(0) = L^2(0,\infty) \cap \mathsf{K}L^2(0,\infty)$. Then, we have $\mathcal{K}(\Theta) = \mathsf{F}(V(0))$, and hence $\mathcal{H}(E) = E\mathsf{F}(V(0)) = \{E(z)\widehat{\psi}(z) \mid \psi \in V(0)\}$.*

Proof It is sufficient to prove that $\mathcal{K}(\Theta) = \mathsf{F}(V(0))$, since $\mathcal{H}(E) = E\mathcal{K}(\Theta)$. The proof below is essentially the same as the proof in [13, Lemma 4.1].

If $\psi \in V(0)$, both $\mathsf{F}\psi$ and $\mathsf{F}\mathsf{K}\psi$ belong to the Hardy space H^2 by definition (5.1) and $H^2 = \mathsf{F}(L^2(0,\infty))$. On the other hand, we have $(\mathsf{F}\mathsf{K}\psi)(z) = \Theta(z)(\mathsf{F}\psi)^\sharp(z)$ by definition (5.1) again. This implies $(\mathsf{F}\psi)(z) = \Theta(z)(\mathsf{F}\mathsf{K}\psi)^\sharp(z)$, since $\Theta(z)\Theta^\sharp(z) = 1$ by definition (1.4). Therefore, $\mathsf{F}\psi$ belongs to $\mathcal{K}(\Theta)$ by (2.7).

Conversely, if $F \in \mathcal{K}(\Theta)$, there exists $f \in L^2(0,\infty)$ and $g \in L^2(-\infty,0)$ such that

$$F(z) = (\mathsf{F}f)(z) = \Theta(z)(\mathsf{F}g)(z).$$

We have $(\mathsf{F}g)^\sharp(z) = \Theta(z)(\mathsf{F}f)^\sharp(z)$ by using $\Theta(z)\Theta^\sharp(z) = 1$ again. Here, $(\mathsf{F}g)^\sharp(z) = (\mathsf{F}\tilde{g})(z)$ for $\tilde{g}(x) = \overline{g(-x)} \in L^2(0,\infty)$, and $\Theta(z)(\mathsf{F}f)^\sharp(z) = (\mathsf{F}\mathsf{K}f)(z)$ as above. Hence, $\mathsf{K}f$ belongs to $L^2(0,\infty)$, and thus $f \in V(0)$. ∎

Remark 5.2 By Lemma 5.1, it follows that the RH would be false if $V(0) = 0$, since $A(z)/(z-\gamma) = \xi(1/2 - iz)/(z-\gamma)$ belongs to $\mathcal{H}(E)$ for all $\gamma \in \Gamma$ under the assumption of the RH. Therefore, it is an interesting problem to prove or disprove $V(0) \neq 0$ unconditionally. Since $V(0)$ is K-invariant, if $V(0) \neq 0$, then for any nonzero $f \in$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

$V(0)$, the functions $(1 \pm \mathsf{K})f$ are eigenfunctions of K with eigenvalues ± 1. Hence, the problem reduces to determining whether the isometric involution K admits an eigenfunction in $L^2(0,\infty)$, which appears to be extremely difficult. We therefore do not pursue this issue further in the present article.

Let $\tau = \tau_\xi$ be the measure on $\mathbb{R}$ determined from the screw function $g = g_\xi$ by (2.3). Then, we have $g(0) = 0$, $b = 0$, and

$$d\tau(\lambda) = \sum_{\gamma\in\Gamma} m_\gamma \delta(\lambda - \gamma)\, d\lambda, \quad \lambda \in \mathbb{R}, \tag{5.3}$$

since

$$g(t) = \sum_{\gamma\in\Gamma} m_\gamma \frac{e^{i\gamma t} - 1}{\gamma^2}$$

by [14, Theorem 1.1(2)], where δ is the Dirac mass at $\lambda = 0$, We understand that the Hilbert space $L^2(\tau)$ is the space of sequences $S = (S(\gamma))_{\gamma\in\Gamma}$ with

$$\|S\|^2_{L^2(\tau)} = \sum_{\gamma\in\Gamma} m_\gamma |S(\gamma)|^2. \tag{5.4}$$

Then, we prove two isomorphisms for $L^2(\tau)$ necessary for the proof of Theorem 1.1.

Lemma 5.3 *Hilbert spaces $V(0)$ and $L^2(\tau)$ are isomorphic by the linear map*

$$V(0) \ni \psi \mapsto S_\psi := \left(\widehat{\psi}(\gamma)\right)_{\gamma\in\Gamma} \in L^2(\tau)$$

with

$$2\|\psi\|^2_{L^2(\mathbb{R})} = \|S_\psi\|^2_{L^2(\tau)}. \tag{5.5}$$

Proof Let μ_Θ be the measure on $\mathbb{R}$ determined from $\Theta = \Theta_\xi$ by (2.8). Then, the linear map $\mathcal{K}(\Theta) \to L^2(\mu_\Theta)$ given by $\widehat{\psi} \mapsto S_\psi$ is an isometric isomorphism as reviewed in Section 2.4. On the other hand, $L^2(\mu_\Theta) = L^2(\tau)$ with $\|S\|^2_{L^2(\mu_\Theta)} = \pi\|S\|^2_{L^2(\tau)}$ by (2.8), (4.1), and (5.3). Therefore, by composing the maps $V(0) \to \mathcal{K}(\Theta) = \mathcal{F}(V(0))$ and $\mathcal{K}(\Theta) \to L^2(\mu_\Theta)$, we obtain the conclusion of the lemma, since $2\pi\|\psi\|^2_{L^2(\mathbb{R})} = \|\widehat{\psi}\|^2_{L^2(\mathbb{R})}$. ∎

Lemma 5.4 *For $\psi = \lim_{n\to\infty} \psi_n \in \mathcal{H}_W$ with $\{\psi_n\}_{n\geq 1} \subset C_c^\infty(\mathbb{R})$, we define $S_\psi \in L^2(\tau)$ by*

$$S_\psi := \lim_{n\to\infty} \left(\widehat{\psi_n}(\gamma)\right)_{\gamma\in\Gamma} \quad \textit{in} \quad L^2(\tau).$$

Then, it is well-defined and provides an isomorphism between $\mathcal{H}_W$ and $L^2(\tau)$ through the mapping

$$\mathcal{H}_W \ni \psi \mapsto S_\psi \in L^2(\tau)$$

with

$$\langle \psi, \psi \rangle_W = \|S_\psi\|^2_{L^2(\tau)}. \tag{5.6}$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Proof We consider $C_0^\infty(\mathbb{R}) = \{\phi \in C_c^\infty(\mathbb{R}) \mid \widehat{\phi}(0) = 0\}$, since we obtain the same completion $\mathcal{H}(G_g)$ even if we start from this space instead of $C_0(\mathbb{R})$. Then differentiation $\psi \mapsto \psi'$ gives a bijection from $C_c^\infty(\mathbb{R})$ to $C_0^\infty(\mathbb{R})$. The inverse map is $\phi \mapsto \int_{-\infty}^x \phi(y)\,dy$. The Weil Hermitian form and the Hermitian form $\langle\cdot,\cdot\rangle_{G_g}$ defined by (2.2) for the screw function g are related as in (4.10), which is written as

$$\langle\phi,\phi\rangle_{G_g} = \langle\psi,\psi\rangle_W, \quad \psi(x) = \int_{-\infty}^x \phi(y)\,dy, \quad \psi \in C_c^\infty(\mathbb{R}). \tag{5.7}$$

(Although not necessary for the proof, $\langle\phi,\phi\rangle_{G_g}$ and $\langle\psi,\psi\rangle_W$ are positive definite on $C_0^\infty(\mathbb{R})$ and $C_c^\infty(\mathbb{R})$, respectively, by [14, Lemma 2.1].) Relation (5.7) extends to the completed Hilbert spaces. Therefore, $\mathcal{H}_W$ is isometrically isomorphic to the Hilbert space $\mathcal{H}(G_g)$ by $\mathcal{H}(G_g) \to \mathcal{H}_W : [\phi] \mapsto [\psi]$ with $\psi = \lim_{n\to\infty} \psi_n$ and $\psi_n(x) = \int_{-\infty}^x \phi_n(y)\,dy$ for $\phi = \lim_{n\to\infty} \phi_n$ ($\phi_n \in C_c^\infty(\mathbb{R})$).

We define $\mathcal{H}(G_g) \to L^2(\tau)$ as follows. For $[\phi] \in \mathcal{H}(G_g)$, we define $S_\phi = (S_\phi(\gamma))_{\gamma\in\Gamma} \in L^2(\tau)$ by

$$\lim_{n\to\infty} \left(\widehat{\phi_n}(\gamma)/\gamma\right)_{\gamma\in\Gamma} \quad \text{in} \quad L^2(\tau)$$

using a sequence $(\phi_n)_n$ in $C_0^\infty(\mathbb{R})$ satisfying $\phi = \lim_{n\to\infty} \phi_n$. Then, the map is well-defined and $\langle[\phi],[\phi]\rangle_{G_g} = \langle\phi,\phi\rangle_{G_g} = \|S_\phi\|_{L^2(\tau)}$ by (2.2), (4.6), and (5.4). Therefore, it establishes the isomorphic isomorphism $\mathcal{H}(G_g) \to L^2(\tau) : [\phi] \mapsto S_\phi$ [7, Sections 5.3 and 12.5]. Using $\mathcal{H}(G_g) \to \mathcal{H}_W$ and noting $\widehat{\phi}(\lambda)/\lambda = i\widehat{\psi}(\lambda)$ for $\phi \in C_0^\infty(\mathbb{R})$, we define $\mathcal{H}_W \to L^2(\tau)$ by $[\psi] \mapsto S_\psi$ with

$$S_\psi = (S_\psi(\gamma))_{\gamma\in\Gamma} = \lim_{n\to\infty} \left(\widehat{\psi_n}(\gamma)\right)_{\gamma\in\Gamma} = \lim_{n\to\infty} \left(-i\widehat{\phi_n}(\gamma)/\gamma\right)_{\gamma\in\Gamma} \quad \text{in} \quad L^2(\tau),$$

where $(\phi_n)_n$ is a sequence in $C_0^\infty(\mathbb{R})$ such that $\psi = \lim_{n\to\infty} \psi_n$ with $\phi_n = \psi_n'$. Then, the map is well-defined and

$$\langle[\psi],[\psi]\rangle_W = \langle\psi,\psi\rangle_W = \|S_\psi\|_{L^2(\tau)} = \|S_\phi\|_{L^2(\tau)} = \langle\phi,\phi\rangle_{G_g} = \langle[\phi],[\phi]\rangle_{G_g}$$

holds, where $\phi = \lim_{n\to\infty} \phi_n$ and the second equality follows from (1.2) and (5.4). Hence, it establishes an isometric isomorphism $\mathcal{H}_W \to L^2(\tau)$ by $[\psi] \mapsto S_\psi$. As a result, the mapping $\mathcal{H}_W \to L^2(\tau)$ is directly defined by $S_\psi = \lim_{n\to\infty} \left(\widehat{\psi_n}(\gamma)\right)_{\gamma\in\Gamma}$ and $[\psi] \mapsto S_\psi$ for $\psi = \lim_{n\to\infty} \psi_n$ with the desired equality for norms. ∎

Theorem 5.5 *Assume that the RH is true. Let $\mathcal{H}_W$, $\mathcal{H}(E)$, and $\mathcal{K}(\Theta)$ be as above. Let $V(t)$ be the spaces defined in (5.2). Then, the following hold:*

(1) $\|E\widehat{\psi}\|_{\mathcal{H}(E)}^2 = \|\widehat{\psi}\|_{L^2(\mathbb{R})}^2 = 2\pi\|\psi\|_{L^2(\mathbb{R})}^2 = \pi\langle\psi,\psi\rangle_W$ *for* $\psi \in V(0)$.

(2) *The map from $\mathcal{K}(\Theta)$ to $\mathcal{H}_W$ obtained by the composition of the inverse of*

$$V(0) \to \mathcal{K}(\Theta) : \psi \mapsto \widehat{\psi}(z), \quad 2\pi\|\psi\|_{L^2(\mathbb{R})}^2 = \|\widehat{\psi}\|_{L^2(\mathbb{R})}^2 \tag{5.8}$$

and

$$V(0) \to \mathcal{H}_W : \psi \mapsto [\psi], \quad 2\|\psi\|_{L^2(\mathbb{R})}^2 = \langle[\psi],[\psi]\rangle_W = \langle\psi,\psi\rangle_W \tag{5.9}$$

agrees with the isomorphism $F \mapsto \psi_F$ in Theorem 1.1. In particular, (5.9) is an isometric isomorphism up to a constant multiple.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Proof (1) It suffices to show that the equality

$$\|\psi\|^2_{L^2(\mathbb{R})} = \frac{1}{2}\langle \psi, \psi \rangle_W \tag{5.10}$$

holds, since $\|E\widehat{\psi}\|_{\mathcal{H}(E)} = \|\widehat{\psi}\|_{L^2(\mathbb{R})}$ by (2.4) and $\|\widehat{\psi}\|^2_{L^2(\mathbb{R})} = 2\pi\|\psi\|^2_{L^2(\mathbb{R})}$ by (1.1). For each $\gamma \in \Gamma$, we define $\psi_\gamma \in L^2(\mathbb{R})$ by

$$F_\gamma = \widehat{\psi_\gamma}. \tag{5.11}$$

Then, each ψ_γ belongs to $V(0)$, and $\{\psi_\gamma\}_{\gamma\in\Gamma}$ forms an orthogonal basis satisfying $2\pi\|\psi_\gamma\|^2_{L^2(\mathbb{R})} = \|\widehat{\psi_\gamma}\|^2_{\mathcal{K}(\Theta)} = \|F_\gamma\|^2_{\mathcal{K}(\Theta)} = 1$ by Proposition 4.1 and Lemma 5.1, since the orthogonality of F_γ's is preserved under the Fourier transform. For $\psi = \sum_\gamma c_\gamma \psi_\gamma \in V(0)$, we have

$$\|\psi\|^2_{L^2(\mathbb{R})} = \frac{1}{2\pi}\sum_{\gamma\in\Gamma} |c_\gamma|^2$$

by the orthogonality and

$$\widehat{\psi}(\gamma) = \frac{1}{\sqrt{m_\gamma \pi}}\, c_\gamma$$

by applying (4.2) to $\widehat{\psi} = \sum_\gamma c_\gamma F_\gamma$. From these two and (1.2), we get (5.10).

(2) It is clear that the composition of the inverse of (5.8) and (5.9) agrees with the map $F \mapsto \psi_F$ of Theorem 1.1 including the equality for norms, and we observed in the proof of Lemma 5.3 that the map (5.8) is an isometric isomorphism up to the multiple $\sqrt{2\pi}$. Therefore, it suffices to show that the map (5.9) gives an isometric isomorphism up to the multiple $\sqrt{2}$.

For $\psi \in V(0)$, the function $S_\psi \in L^2(\tau)$ is defined and satisfies $2\|\psi\|^2_{L^2(\mathbb{R})} = \|S_\psi\|^2_{L^2(\tau)}$ by Lemma 5.3. Then, there exists a sequence $(\psi_n^*)_n \subset C_c^\infty(\mathbb{R})$ that converges to ψ^* with respect to $\langle\cdot,\cdot\rangle_W$ and $S_\psi = S_{\psi^*}$ by Lemma 5.4. The latter implies $\langle \psi - \psi_n^*, \psi - \psi_n^*\rangle_W = \langle \psi^* - \psi_n^*, \psi^* - \psi_n^*\rangle_W \to 0$ $(n \to \infty)$. Therefore, $\psi = \psi^*$, and hence $V(0) \to \mathcal{H}_W$ is directly defined by $\psi \mapsto [\psi]$. Furthermore, we obtain $2\|\psi\|^2_{L^2(\mathbb{R})} = \langle \psi, \psi\rangle_W$ from (5.5) and (5.6). Hence, this map is precisely the one given in (5.9). ∎

The equality $\|\psi\|^2_{L^2(\mathbb{R})} = 2^{-1}\langle \psi, \psi\rangle_W$ in Theorem 5.5 (1) shows that the L^2-structure induced from $L^2(\mathbb{R})$ and "arithmetic structure" (or "local structure") arising from the geometric side of the Weil explicit formula (3.3) coincide on a dense subspace of $V(0)$ consisting of functions for which the Weil explicit formula holds.

Theorem 5.6 *Let $\mathcal{H}_0$ and $\mathcal{K}_0$ are Hilbert spaces defined unconditionally in Section 3.3. Assume that the RH is true. Then, $\mathcal{H}_0 = \mathcal{H}_W$ and $\mathcal{K}_0 = \mathcal{K}(\Theta)$, and the extended map $\widehat{\mathcal{P}_D} : \mathcal{H}_W \to \mathcal{K}(\Theta)$ provides the inverse of the map in Theorem 5.5 (2). In particular, $V(0)$ is the L^2-closure of $V^\circ(0)$ in Corollary 1.5.*

Proof For $\psi \in C_c^\infty(\mathbb{R})$, we have

$$\|\widehat{\mathcal{P}_{D\psi}}\|^2_{L^2(\mathbb{R})} = \pi\sum_{\gamma\in\Gamma} m_\gamma |\widehat{\psi}(\gamma)|^2 = \pi\langle \psi, \psi\rangle_W$$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

by (1.2), (3.8), and Proposition 4.1. Hence, $\mathcal{H}_0$ coincides with $\mathcal{H}_W$ by definition (3.9). Formula (3.8) shows that the image $\widehat{\mathcal{P}_{D\psi}}$ is defined independent of the representatives of $[\psi]$ in $\mathcal{H}_W$. On the other hand, $\mathcal{K}_0$ is a subspace of $\mathcal{K}(\Theta)$ by Proposition 4.1 again.

We denote $F = \widehat{\mathcal{P}_{D\psi}}$ for $[\psi] \in \mathcal{H}_W$ and set $\psi_F = \mathsf{F}^{-1}(F)$ as in Theorem 1.1. Then, $F(\gamma) = \widehat{\psi}(\gamma)$ by (3.8) and (4.2). Therefore, $\widehat{\psi_F}(\gamma) = \psi(\gamma)$ for all $\gamma \in \Gamma$, and hence $[\psi] = [\psi_F]$ in $\mathcal{H}_W$. On the other hand, $\widehat{\mathcal{P}_{D\psi_F}}(z) = F$ by (3.8), since $\widehat{\psi_F} = F$ by definition and $F(\gamma) = \widehat{\psi}(\gamma)$. Hence, we obtain the desired conclusion. ∎

The totally ordered structure of the subspaces of the de Branges space $\mathcal{H}(E)$ is described by $V(t)$ as follows.

Theorem 5.7 *Assume that the RH is true. Then, $E\,\mathsf{F}(V(t))$ is a de Branges subspace of $\mathcal{H}(E)$ for every $t \geq 0$ and is isometrically isomorphic to $\mathcal{H}_W(t)$ up to a constant multiple by the map of Theorem 1.1.*

Proof It is sufficient to prove the first half of the theorem, since the second half follows from Theorem 5.5 (2). We prove the claim for positive t such that $V(t) \neq \{0\}$, since the case of $t = 0$ was proved in Lemma 5.1 and the claim is trivial if $V(t) = \{0\}$. The following is essentially the same as the proof of [13, Lemma 4.3].

We show that $\mathcal{H} := E(z)\mathsf{F}(V(t))$ is a Hilbert space consisting of entire functions and satisfies the axiom of the de Branges spaces:

(dB1) For each $z \in \mathbb{C}\backslash\mathbb{R}$, the point evaluation $\Phi \mapsto \Phi(z)$ is a continuous linear functional on $\mathcal{H}$.
(dB2) If $\Phi \in \mathcal{H}$, $\Phi^\sharp$ belongs to $\mathcal{H}$ and $\|\Phi\|_{\mathcal{H}} = \|\Phi^\sharp\|_{\mathcal{H}}$.
(dB3) If $w \in \mathbb{C}\backslash\mathbb{R}$, $\Phi \in \mathcal{H}$ and $\Phi(w) = 0$,

$$\frac{z-\bar{w}}{z-w}\Phi(z) \in \mathcal{H} \quad \text{and} \quad \left\|\frac{z-\bar{w}}{z-w}\Phi(z)\right\|_{\mathcal{H}} = \|\Phi\|_{\mathcal{H}},$$

where the Hilbert space structure is the one induced from $V(t)$ that is equivalent to $\langle F, G\rangle_{\mathcal{H}} = \int_{\mathbb{R}} F(z)\overline{G(z)}|E(z)|^{-2}dz$ for $F, G \in \mathcal{H}$.

Let $\Phi(z) = E(z)(\mathsf{F}f)(z) \in \mathcal{H}$ with $f \in V(t)$. First, we prove that $\mathcal{H}$ consists of entire functions. We see that $\Phi(z)$ is holomorphic in $\mathbb{C}_+$ by $f \in L^2(t, \infty)$. If we write $(\mathsf{J}_\sharp f)(x) := \overline{f(-x)}$, the commutative relation $\mathsf{JF} = \mathsf{FJ}_\sharp$ holds. Therefore, using (5.1) and $\mathsf{K}^2 = 1$, we have $\Phi(z) = E(z)(\mathsf{F}f)(z) = E^\sharp(z)(\mathsf{FJ}_\sharp\mathsf{K}f)(z)$. This shows that $\Phi(z)$ is also holomorphic in $\mathbb{C}_-$. Furthermore, $\mathsf{J}_\sharp\mathsf{K}f \in L^2(-\infty, -t)$, because the tempered distribution kernel $k := \mathsf{F}^{-1}\Theta$ of K has support in $[0, \infty)$ by [10, Theorems 1.1 and 1.2]. On the real line, $\lim_{z\to x}(\mathsf{F}f)(z) = (\mathsf{F}f)(x)$ and $\lim_{z\to x}(\mathsf{FJ}_\sharp\mathsf{K}f)(z) = \lim_{z\to x}(\mathsf{FK}f)^\sharp(z) = \Theta^\sharp(x)(\mathsf{F}f)(x)$ for almost all $x \in \mathbb{R}$, where z is allowed to tend to x nontangentially from $\mathbb{C}_+$ and $\mathbb{C}_-$, respectively. Hence, $\Phi(z)$ is also holomorphic in a neighborhood of each point of $\mathbb{R}$. By the above, $\Phi(z)$ is an entire function.

We confirm (dB1). For $z \in \mathbb{C}_+$, $\Phi \mapsto \Phi(z) = E(z)\int_t^\infty f(x)e^{izx}dx$ is a continuous linear form. On the other hand, for $z \in \mathbb{C}_-$, $\Phi \mapsto \Phi(z) = E^\sharp(z)\int_{-\infty}^{-t}\overline{(\mathsf{K}f)(-x)}e^{izx}\,dx$ is a continuous linear functional.

We confirm (dB2). We have $\Phi^\sharp(z) = E(z)(\mathsf{FK}f)(z)$. Since $\mathsf{K}f \in V(t)$, the function $\Phi^\sharp$ belongs to $\mathcal{H}$. Since K is isometric, the equality of norms in (dB2) holds.

We confirm (dB3). The equality of norms in (dB3) is trivial by the definition of the norm of $\mathcal{H}$. From (dB2), it is sufficient to show only the case of $w \in \mathbb{C}_+$.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

Suppose that $\Phi(w) = 0$ for some $w \in \mathbb{C}_+$. Then, $(\mathsf{F}f)(w) = 0$, since $E(z)$ has no zeros on $\mathbb{C}_+$. We put $f_w(x) := f(x) - i(w - \bar{w}) \int_0^{x-t} f(x-y)e^{-iwy}dy$. Then, we easily find that $f_w \in L^2(t, \infty)$ and $(\mathsf{F}f_w)(z) = ((z-\bar{w})/(z-w))(\mathsf{F}f)(z)$ for $z \in \mathbb{C}_+$. Hence, we complete the proof if it is shown that $\mathsf{K}f_w$ has support in $[t, \infty)$, since $\mathsf{K}f_w \in L^2(\mathbb{R})$ by $f_w \in L^2(t, \infty)$. We put $g_w(x) := (\mathsf{K}f)(x) - i(\bar{w} - w) \int_0^{x-t} (\mathsf{K}f)(x-y)e^{-i\bar{w}y}dy$. Then, g_w has support in $[t, \infty)$ by $\mathsf{K}f \in L^2(t, \infty)$ and $(\mathsf{F}g_w)(z) = ((z-w)/(z-\bar{w}))(\mathsf{FK}f)(z) = (\mathsf{FK}f_w)(z)$ for $z \in \mathbb{C}_+$. Hence, $g_w = \mathsf{K}f_w$ and the proof is completed. ∎

We expect that $V(t) \neq 0$ for some $t > 0$, or rather that $V(t) \neq 0$ holds for all $t \geq 0$, but we do not discuss this in the present article, since it seems to be a nontrivial problem related to the eigenfunctions of K, as mentioned in Remark 5.2.

5.1 A weaker variant of Corollary 1.5

Since the space $V(0)$ can be constructed unconditionally as well as $V^\circ(0)$ in Corollary 1.5, it can be used to state an equivalence condition for the RH. However, since the construction of $V(0)$ is simpler than that of $V^\circ(0)$, more conditions are required for the equivalence condition.

Proposition 5.8 *Let $V(0) = L^2(0, \infty) \cap \mathsf{K}L^2(0, \infty)$ be as in (5.2). Then, the RH is true if and only if the following two conditions hold:*

(1) $\|\psi\|^2_{L^2(\mathbb{R})} = 2^{-1}\langle \psi, \psi \rangle_W$ *for every* $\psi \in V(0)$.

(2) *For a given* $\gamma \in \Gamma$ *and any* $\varepsilon > 0$, *there exists* $\psi \in V(0)$ *such that*

$$\widehat{\psi}(-\gamma) = 1, \qquad |\widehat{\psi}(-\gamma')| \leq \frac{\varepsilon}{|\gamma - \gamma'|^{1+\delta}} \quad \text{for every } \gamma' \in \Gamma \backslash \{\gamma\}$$

for some $\delta > 0$ *independent of* γ, ε, *and* ψ.

Proof Assuming the RH, (1) follows from Theorem 5.5 (1). Also, (2) holds, since $\psi_\gamma = \mathsf{F}^{-1}(F_\gamma)$ in $V(0)$ satisfies $\widehat{\psi_\gamma}(\gamma) \neq 0$ and $\widehat{\psi_\gamma}(\gamma') = 0$ for $\gamma' \in \Gamma \backslash \{\gamma\}$.

Conversely, we assume that (1) and (2) are satisfied. Then, we show that a contradiction arises if the RH is false. We take a nonreal $\gamma_0 \in \Gamma$. For any $\varepsilon > 0$, there exists $\psi_1, \psi_2 \in V(0)$ such that $\widehat{\psi_1}(-\gamma_0) = i$, $\widehat{\psi_2}(-\overline{\gamma_0}) = -i$, $|\widehat{\psi_1}(-\gamma)| \leq \varepsilon |\gamma_0 - \gamma|^{-1-\delta}$ for every $\gamma \in \Gamma \backslash \{\gamma_0\}$, and $|\widehat{\psi_2}(-\gamma)| \leq \varepsilon |\overline{\gamma_0} - \gamma|^{-1-\delta}$ for every $\gamma \in \Gamma \backslash \{\overline{\gamma_0}\}$ by (2). Then, for $\psi := \psi_1 + \psi_2 \ (\neq 0)$, we have $\langle \psi, \psi \rangle_W = \sum_{\gamma \in \Gamma} m_\gamma \widehat{\psi}(-\gamma)(\widehat{\psi})^\sharp(-\gamma) = -m_{\gamma_0} + O(\varepsilon)$, since $\sum_{\gamma \in \Gamma} m_\gamma |\gamma|^{-1-\delta} < \infty$. Therefore, $\langle \psi, \psi \rangle_W$ is negative for a sufficiently small $\varepsilon > 0$, but it contradicts (1). Hence, the RH holds. ∎

6 Hilbert–Pólya space

One of attractive strategies for proving the RH is the construction of a Hilbert–Pólya space, which is a pair of a Hilbert space and a self-adjoint operator acting on it such that all nontrivial zeros of the Riemann zeta-function are eigenvalues of the self-adjoint operator. In this section, we state that $\mathcal{H}_W$ is one of Hilbert–Pólya spaces *under the RH*. Note that $\mathcal{H}_W$ is unconditionally defined as $\mathcal{H}_0$ by Theorem 5.6.

We assume the RH and denote $E = E_\xi$ as in Section 5. In this case, the domain $\mathfrak{D}(\mathsf{M})$ of the multiplication operator M on $\mathcal{H}(E)$ is dense in $\mathcal{H}(E)$, because $S_\theta(z)$

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

does not belongs to $\mathcal{H}(E)$ for all $\theta \in [0, \pi)$ by the estimate $|S_\theta(iy)/E(iy)| \gg (\log y)^{-1}$ $(y \to +\infty)$ obtained by the Stirling formula for the gamma-function and [11, Proposition 2.1]. Using M, we define the operator $\mathsf{A} := \mathsf{F}^{-1}\mathsf{M}\mathsf{F}$ on $V(0)$ with the domain $\mathfrak{D}(\mathsf{A}) = \mathsf{F}^{-1}(\mathfrak{D}(\mathsf{M}))$. If $\psi \in V(0)$ is differentiable and ψ' also belongs to $V(0)$, then $\mathsf{A}\psi = i\psi'$. Further, we define the operator A_W on $\mathcal{H}_W$ as follows.

By Theorem 5.6, the inverse of (5.9) from $\mathcal{H}_W$ to $V(0)$ is given by $[\psi] \mapsto \mathsf{F}^{-1}\widehat{\mathcal{P}_{D\psi}}$. Further, if we choose the representative of ψ from $V(0)$, it is possible and uniquely determined by Theorem 5.5 (2), and therefore $\psi = \mathsf{F}^{-1}\widehat{\mathcal{P}_{D\psi}}$. By choosing representatives in this way, we define A_W on $\mathcal{H}_W$ by $\mathsf{A}_W[\psi] = [\mathsf{A}\psi]$. By the same procedure as above, the family of self-adjoint extensions M_θ of M determines the corresponding families of self-adjoint extensions of A and A_W (see (2.5) and (2.6)). By this correspondence, the orthogonal basis $\{[\psi_\gamma]\}_{\gamma\in\Gamma}$ of $\mathcal{H}_W$ consists of eigenvectors $[\psi_\gamma]$ of $\mathsf{A}_{W,\pi/2}$ with eigenvalues $\gamma \in \Gamma$, since $\{EF_\gamma\}_{\gamma\in\Gamma}$ with (3.5) is an orthogonal basis of $\mathcal{H}(E)$ consists of eigenfunctions of $\mathsf{M}_{\pi/2}$ with eigenvalues Γ (see Seciton 2.3). Therefore, the pair $(\mathcal{H}_W, \mathsf{A}_{W,\pi/2})$ is a Hilbert–Pólya space.

It is important to note that the multiplicity of $\gamma \in \Gamma$ as an eigenvalue of $\mathsf{A}_{W,\pi/2}$ (and $\mathsf{M}_{\pi/2}$) is one. In other words, the multiplicity of $\gamma \in \Gamma$ as a zero of $\xi(1/2 - iz)$ is not reflected in the multiplicity of $\mathsf{A}_{W,\pi/2}$ (and $\mathsf{M}_{\pi/2}$). In particular, it shows the explicit difference between the de Branges space $\mathcal{H}(E_\xi)$ and the de Branges space $\mathcal{B}_\lambda^S$ in [4, Section 4.8].

In the above discussion, we assumed the RH, but (2.5) and (2.6) allow us to define the operator M_θ without the RH. However, its properties as an operator become unclear.

7 Special values of the screw line $\mathfrak{S}_t(z)$

The screw line $\mathfrak{S}_t(z)$ has the following unconditional relations with the screw function $g(t)$. It is interesting that they are not a special case of equations obtained from the general theory of screw functions.

Theorem 7.1 *Let $g_\xi(t)$ and $\mathfrak{P}_t(z)$ be functions of (4.3) and (1.6), respectively. Then, the following equations hold independently of the truth of the RH:*

$$\mathfrak{P}_t(0) = -g_\xi(t), \tag{7.1}$$

$$\lim_{y\to+\infty}\left[y\,\mathfrak{B}_t(-iy) - \frac{1}{2}\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}+\frac{y}{2}\right) + \frac{1}{2}\log\pi\right] = -g'_\xi(t), \tag{7.2}$$

where we assume $t \neq \log n$ for any $n \in \mathbb{N}$ in (7.3).

Proof Equality (7.1) follows from (3.2), Proposition 3.1, and [14, Theorem 1.1 (2)], but it follows directly from (4.3) and (1.6) as follows. By $\Phi(z, s, a) = \sum_{n=0}^{\infty} z^n (n+a)^{-s}$ and (2.8),

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

$$\lim_{z\to 0}\frac{1}{iz}\Big[\Phi(e^{-2t},1,\tfrac{1}{2}(\tfrac{1}{2}-iz))-\Phi(e^{-2t},1,1/4)\Big]=-\frac{1}{2}\Phi(e^{-2t},2,1/4),$$
$$\lim_{z\to 0}\frac{1}{iz}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}-\frac{iz}{2}\right)-\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)\right]=\frac{1}{2}\psi_1\left(\frac{1}{4}\right),$$

where $\psi_1(z)$ is the polygamma function of order one. The expansion $\psi_1(w)=\sum_{n=0}^{\infty}(w+n)^{-2}$ gives $\psi_1(1/4)=\Phi(1,2,1/4)$. Taking $s=1/2$ in the logarithmic derivative of $\xi(s)=\xi(1-s)$ and using

$$\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)=-\gamma_0-3\log 2-\frac{\pi}{2},$$

we have

$$\frac{\zeta'}{\zeta}\left(\frac{1}{2}\right)=\frac{1}{2}\left(\gamma_0+3\log 2+\log\pi+\frac{\pi}{2}\right).$$

Hence, by taking the limit $z\to 0$ in (1.6), we obtain the minus of (4.3).

To show (7.3), we multiply (1.6) by y and substitute $-iy$ for z:

$$\begin{aligned}y\mathfrak{P}_t(-iy):=&\frac{4y(e^{t/2}-1)}{1+2y}+\frac{4y(e^{-t/2}-1)}{1-2y}\\&+(e^{-yt}-1)\frac{\zeta'}{\zeta}\left(\frac{1}{2}-y\right)+\sum_{n\le e^t}\frac{\Lambda(n)}{\sqrt{n}}(e^{-y(t-\log n)}-1)\\&+\frac{1}{2}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)-\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}-\frac{y}{2}\right)\right]\\&+\frac{1}{2}e^{-t/2}\left[\Phi(e^{-2t},1,1/4)-\Phi(e^{-2t},1,\tfrac{1}{2}(\tfrac{1}{2}-y))\right].\end{aligned}$$

Therefore, for positive $t>0$,

$$\begin{aligned}&\lim_{y\to+\infty}\left[y\mathfrak{B}_t(-iy)-\frac{1}{2}\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}+\frac{y}{2}\right)+\frac{1}{2}\log\pi\right]\\&=2(e^{t/2}-e^{-t/2})-\sum_{n\le e^t}\frac{\Lambda(n)}{\sqrt{n}}+\frac{1}{2}\left[\frac{\Gamma'}{\Gamma}\left(\frac{1}{4}\right)-\log\pi\right]+\frac{1}{2}e^{-t/2}\Phi(e^{-2t},1,1/4)\end{aligned}$$

by using the logarithmic derivative of $\xi(s)=\xi(1-s)$ at $s=1/2-y$. The right-hand side equals $-g'_\xi(t)$ if $t\neq\log n$ by (4.3), and $(d/dt)(e^{-t/2}\Phi(e^{-2t},2,1/4))=-2e^{-t/2}\Phi(e^{-2t},2,1/4)$ follows from $\Phi(z,s,a)=\sum_{n=0}^{\infty}z^n(n+a)^{-s}$. ∎

References

[1] N. A. Baas and C. F. Skau, *The lord of the numbers, Atle Selberg. On his life and mathematics*. Bull. Amer. Math. Soc. (N.S.) **45**(2008), no. 4, 617–649.

[2] E. Bombieri and J. C. Lagarias, *Complements to Li's criterion for the Riemann hypothesis*. J. Number Theory **77**(1999), no. 2, 274–287.

[3] E. Bombieri, *Remarks on Weil's quadratic functional in the theory of prime numbers. I*. Atti Accad Naz. Lincei Cl. Sci. Fis. Mat. Natur. Rend. Lincei (9) Mat. Appl. **11**(2001), no. 3, 183–233.

[4] A. Connes, C. Consani, and H. Moscovici, *Zeta zeros and prolate wave operators*. Ann. Funct. Anal. **15**(2024), 87.

[5] L. de Branges, *Hilbert spaces of entire functions*. Prentice-Hall, Inc., Englewood Cliffs, NJ, 1968.

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press

[6] M. Kaltenbäck and H. Woracek, *Pontryagin spaces of entire functions I*. Integral Equ. Oper. Theory **33**(1999), no. 1, 34–97.
[7] M. G. Kreĭn and H. Langer, *Continuation of hermitian positive definite functions and related questions*. Integr. Equ. Oper. Theory **78**(2014), no. 1, 1–69.
[8] J. C. Lagarias, *Hilbert spaces of entire functions and Dirichlet L-functions*. In: P. Cartier, B. Julia, P. Moussa, and P. Vanhove (eds.), Frontiers in number theory, physics, and geometry. I, Springer, Berlin, 2006, pp. 365–377.
[9] N. Makarov and A. Poltoratski, *Meromorphic inner functions, Toeplitz kernels and the uncertainty principle*. In: M. Benedicks, P. W. Jones, and S. Smirnov (eds.), Perspectives in analysis, Mathematical Physics Studies, 27, Springer, Berlin, 2005, pp. 185–252.
[10] T. Qian, Y. Xu, D. Yan, L. Yan, and B. Yu, *Fourier spectrum characterization of hardy spaces and applications*. Proc. Am. Math. Soc. **137**(2009), no. 3, 971–980.
[11] C. Remling, *Schrödinger operators and de Branges spaces*. J. Funct. Anal. **196**(2002), no. 2, 323–394.
[12] L. O. Silva and J. H. Toloza, *de Branges spaces and Kreĭn's theory of entire operators*. In: D. Alpay (ed.), Operator theory, Springer, Basel, 2015, pp. 549–580.
[13] M. Suzuki, *An inverse problem for a class of canonical systems having Hamiltonians of determinant one*. J. Funct. Anal. **279**(2020), no. 12, 108699.
[14] M. Suzuki, *Aspects of the screw function corresponding to the Riemann zeta function*. J. Lond. Math. Soc. **108**(2023), no. 4, 1448–1487.
[15] M. Suzuki, *Li coefficients as norms of functions in a model space*. J. Number Theory **252**(2023), 177–194.
[16] E. C. Titchmarsh and D. R. Heath-Brown (ed.), *The theory of the Riemann zeta-function*. 2nd ed., Edited and with a preface by, The Clarendon Press, Oxford University Press, New York, NY, 1986.
[17] A. Weil, *Sur les "formules explicites" de la théorie des nombres premiers*. Commun. Sém. Math. Univ. Lund [Medd. Lunds Univ. Mat. Sem.] **1952**(1952), 252–265.
[18] H. Woracek, *De Branges spaces and growth aspects*. In: D. Alpay (ed.), Operator theory, Springer, Basel, 2015, pp. 489–523.
[19] H. Yoshida, *On Hermitian forms attached to zeta functions*. In: N. Kurokawa and T. Sunada (eds.), Zeta functions in geometry (Tokyo, 1990), Advanced Studies in Pure Mathematics, 21, Kinokuniya, Tokyo, 1992, pp. 281–325.

Department of Mathematics, School of Science, Institute of Science Tokyo, 2-12-1 Ookayama, 152-8551, Japan
e-mail: msuzuki@math.sci.isct.ac.jp

https://doi.org/10.4153/S0008414X25101739 Published online by Cambridge University Press